

Correct and incorrect knowledge of the risk factors concerning the development of arterial hypertension. Part 2. Population of patients with identified arterial hypertension

Andrzej Kleinrok^{1,2}, Michał Kamiński¹, Tomasz Domański¹, Grażyna Prokop-Lewicka¹

¹The Pope John Paul II Province Hospital, Zamość, Poland

²University of Management and Administration in Zamość

Head of department: Andrzej Kleinrok

Abstract

Introduction. Hypertension (HT) and its complications significantly shorten the life expectancy in the world population. In addition to pharmacotherapy, life-style changes play a key role in treatment, which requires appropriate knowledge among patients.

Material and methods. The study was conducted in a group of 613 adults with HT. After obtaining sociodemographic data, performing anthropometric measurements, measuring resting blood pressure twice, the knowledge was assessed using a personally conducted questionnaire. The questions evaluated the effects of smoking, alcohol, coffee, salt, fat, irregular diet, low fibre and vegetable intake, as well as skimmed dairy products in diet, overweight and obesity, cholesterol, physical activity and stressful lifestyle on the development of HT.

Results. False knowledge of the influence on the development of HT in the case of alcohol drinking was found in 16% of the respondents, coffee — 56.4%, large amounts of salt — 7.2%, large amounts of fats — 88%, small amounts of fibre and vegetables and skimmed dairy products — 27.8%, irregular diet — 46.4%, cigarette smoking — 73.4%, overweight and obesity — 4.4%, high cholesterol — 86.9%, low physical activity — 11.2% %, stressful lifestyle — 5.4%

Conclusions. Incorrect knowledge of HT risk factors was most likely to be associated with excessive intake of fat, elevated cholesterol, smoking and coffee drinking. There were no common features of the people who answered the questions incorrectly. The results show the need to educate HT patients about the factors that aggravate this disease.

key words: hypertension, risk factors, knowledge

Arterial Hypertens. 2017, vol. 21, no. 2, pages: 83–92

DOI: 10.5603/AH.2017.0011

Introduction

Hypertension (HT) is a risk factor of many diseases, including renal failure, aortic dissection, stroke, ischaemic heart disease, significantly shortening life in the general population [24–26].

Raised blood pressure (RR), by increasing post-load, causes left ventricular hypertrophy [27]. This results in a progressive restriction of wall motion leading to diastolic dysfunction. In addition, as a result of the compression of the hypopharynx capillaries

Address for correspondence: lek. Michał Kamiński

The Pope John Paul II Province Hospital

22–400 Zamość, Poprzeczna 17/18

tel.: 663 358 271

e-mail: m.kaminski@onet.pl

and the reduction of their density, the features of ischaemia appear. Left ventricular hypertrophy leads to left atrial enlargement, its remodelling and consequent supraventricular arrhythmias, including atrial fibrillation. In this type of arrhythmia, HT not only contributes to its formation but also raises the risk of ischaemic stroke in patients affected by this disorder. Reconstruction of the myocardium significantly increases the risk of ventricular arrhythmias, and consequently may lead to sudden cardiac death [28, 29].

HT leads to thickening of the intima and media of peripheral arteries, contributing to increased stiffness and atherothrombosis. This leads to the formation of ischaemic stroke in case of damage to carotid arteries and to acute coronary syndrome in case of coronary artery occlusion. Raised RR is also a major risk factor of aortic dissection [25, 29].

Kidneys are other organs that are damaged in the course of HT. Damage to renal artery endothelial cells leads to local inflammation, which results in gradual remodelling and fibrosis of the renal interstitium. Initially no clinical symptoms are observed, but over time the symptoms of chronic kidney disease become apparent [27, 29].

Raised RR is considered one of the main risk factors for both ischaemic stroke and haemorrhagic stroke. It is worth pointing out, however, that it also leads to clinically silent episodes of ischaemia resulting in dementia. In the case of the eye organ, arterial remodelling and bleeding that damage the retina occur, resulting in lower quality of life [27–29].

In order to prevent these complications, it is necessary to treat HT effectively, which, apart from pharmacotherapy, is based on lifestyle modification [27]. Therefore, it is so important for a patient to cooperate, which is inextricably linked to their knowledge of the factors that aggravate the disease.

The causes of HT development and, consequently, the increase in the demand for antihypertensive drugs include the consumption of alcohol, high amounts of sodium, small amounts of fibre, vegetables and skimmed dairy products, overweight and obesity, low physical activity and stressful lifestyle.

There were theoretical foundations to believe that some factors could affect the severity of HT, but no correlation was found in the study. This group includes cigarette smoking, coffee consumption, high fat intake, irregular diet, and lipid disorders.

With the aim of increasing the effectiveness of prevention it is helpful to know the group of patients with incorrect knowledge. Thus, it seemed interesting to assess the knowledge that HT patients have, and to find common characteristics of people with

misconceptions. The aim of the study is estimate the HT patient's knowledge and extracting groups with incorrect information.

Material and methods

Data for the study were collected for the bachelor's degree thesis in the period from June to October 2012 in the Lublin Region by the students* of the Nursing Department of the Faculty of Physiotherapy and Pedagogy of the University of Management and Administration in Zamość. The study included an incidental group of 613 people, including 268 men, who reported to the GP practice due to HT. Each participant delivered a written consent to participate in the study. The scope and method of collecting data have been shown in Part 1.

To assess the knowledge the authorial survey questionnaire, approved by the Authorities and the Bioethics Committee of the University of Management and Administration in Zamość, was used. The data were collected personally by interviewers who asked about the effects of alcohol, coffee, large amounts of salt, large amounts of fat, small amounts of fibre and vegetables, and skimmed dairy products in the diet, irregular meals, smoking cigarettes and obesity, high cholesterol, insufficient physical activity, stressful lifestyle on development of HT.

The respondents could assess the impact of factors on HT by marking: high/moderate/low/no impact. They could also answer that they have no opinion. Responses were divided into two groups: high impact i.e. the combination of moderate and high impact responses and no impact, i.e. the combination of no and low impact responses.

The method of statistical analysis has been presented in Part 1.

Results

In the study group, the average age was 60.2 years, and women accounted for over 56%. More than half of respondents lived in the city, more than 40% had basic or vocational education and almost three quarters were married. The average heart rate in the study group was 77.5 beats per minute, mean systolic blood pressure was 141 mmHg and diastolic blood pressure was 84.7 mmHg. The remaining data have been shown in Table I.

Nearly 95% of the study population had either a poor knowledge or were unaware of the effect of

*Data collected by: Iwona Kobel, Barbara Kuszykiewicz, Marzena Tyś, Agata Gardyńska, Violetta Mazur.

Table I. General population characteristics (n = 613)

Sociodemographic data	Respondents
Male gender	268 (43.7)
Age (years)	60.2 (12.7)
Residence — city	316 (51.5)
Married	457 (74.5)
Education secondary or higher	350 (57.1)
Professionally active including:	253 (41.3)
— physical work	110 (43.5)
— intellectual work	143(56.5)
Number of people living in the household	2.7 (± 1.6)
Monthly income/person (PLN) n = 363*	1202.4 (± 993.7)
Medical data	
Current body weight [kg]	81.8 (± 15.1)
Current height [cm]	167.5 (± 9.6)
Waist measurement [cm]	98.1 (± 15.4)
— women [cm]	96.1 (± 16.4)
— men [cm]	100.7 (± 13.6)
Mean systolic RR [mmHg]	141.0 (± 17.0)
Mean diastolic RR [mmHg]	84.7 (± 9.9)
Average pulse rate (bpm)	77.5 (± 23.1)
BMI	29.2 (± 6.6)
— women	28.8 (± 4.8)
— men	29.7 (± 8.5)
Current smokers	96 (15.7)
Current smokers and ex-smokers	224 (36.5)
Average weekly consumption of pure alcohol [g]	34.4 (± 100.3)
— women	9.3 (± 58.5)
— men	65.4 (± 130.1)

Data presented as n (%) or mean (± SD)
RR — blood pressure; BMI — Body Mass Index; *number of respondents

high fat intake and high cholesterol on HT development. About 70% of the respondents had none or incorrect information about the impact of coffee consumption and irregular diet on HT, and as much as 86% about the influence of cigarette smoking on this disease. More than half of the respondents reported that eating small amounts of fibre, vegetables and skimmed dairy products did not affect HT, which is incompatible with current knowledge, or did not know about such a relationship, and less than one third gave a similar answer in case of alcohol. Of the respondents, 19% did not have a proper knowledge of the relationship between low physical activity and HT and almost 12% of the connection between high salt intake and HT. The respondents had

best knowledge in the case of the influence of stressful lifestyle and overweight and obesity on the abovementioned disease. Here 9% and 7.3% respectively had no opinion or gave an incorrect answer (Table II).

Misconceptions about the effects of alcohol on the development of HT were reported in 16% of respondents. More often, people who did not receive information from the Internet (26% vs. 16.6%; $p = 0.019$) had incorrect information (Tables II and III).

Coffee, as a substance affecting HT, was indicated by 56.4% of respondents. This view was more frequently presented by people who received information from family and friends (70.9% vs. 61.4%; $p = 0.031$) (Tables II and IV).

In the study group, 73.4% of people believed that cigarette smoking influences the development of HT, which is incompatible with current knowledge. Particularly, people who received information from health professionals (86.5% vs. 76.6%; $p = 0.007$) and non-smokers (85.5 vs. 77.2 $p = 0.045$) had false knowledge (Tables II and V).

In the case of salt intake, 7.2% of the respondents believed that it did not influence the development of HT, which is incompatible with the current state of knowledge. Such opinion was slightly more frequent, at the edge of statistical significance, for those who did not receive information from health professionals (11.1% vs. 6.3%; $p = 0.059$) (Tables II and VI).

As many as 88% of respondents mistakenly answered the question about the effect of high intake of fat on the development of HT. Especially women (96.1% vs. 92.1%; $p = 0.043$) and people who achieved information from health professionals (96.1% vs. 89%; $p = 0.002$) had incorrect information (Tables II and VII).

Incorrect information about the influence of irregularly consumed meals on HT development was reported by 46.4% of the respondents. In the study group people acquiring information from health professionals (64.3% vs. 50.0%; $p = 0.009$), family and friends (68.7% vs. 57.5%; $p = 0.022$) and the Internet (4% vs. 58.5%; $p = 0.042$) were predominant. (Tables II and VIII).

In the study group, 27.8% of people had incorrect information about the effect of low fibre intake and vegetable consumption and skimmed dairy products on the development of HT. More often, people who did not achieve information from health professionals (46.0% vs. 37.1%; $p = 0.099$) had incorrect information. The above relationship was at the edge of statistical significance (Tables II and IX).

Overweight and obesity as a factor not affecting HT were indicated by 5.1% of respondents. More

Table II. Patients' knowledge of factors affecting HT development (n = 613)

Patients' knowledge of factors affecting HT development	Impact				No opinion
	High	Moderate	Low	No	
Alcohol consumption	266 (43.4)	157 (25.6)	84 (13.7)	14 (2.3)	92 (15.0)
Coffee consumption	145 (23.6)	201 (32.8)	144 (23.5)	47 (7.7)	76 (12.4)
Smoking	318 (51.9)	132 (21.5)	76 (12.4)	9 (1.5)	78 (12.7)
Consuming large amount of salt	426 (69.5)	116 (18.9)	35 (5.7)	9 (1.5)	27 (4.4)
Consuming large amount of fat	402 (65.6)	137 (22.4)	22 (3.6)	10 (1.6)	42 (6.8)
Irregular diet	124 (20.3)	160 (26.1)	141 (23.0)	40 (6.5)	148 (24.1)
Consuming a small amount of fibre, vegetables and skimmed dairy products	128 (20.9)	134 (21.8)	112 (18.3)	58 (9.5)	181 (29.5)
Overweight and obesity	499 (81.4)	69 (11.3)	23 (3.8)	4 (0.6)	18 (2.9)
High level of cholesterol	444 (72.4)	89 (14.5)	29 (4.8)	5 (0.8)	46 (7.5)
Low physical activity	343 (56.0)	153 (25.0)	53 (8.6)	16 (2.6)	48 (7.8)
Stressful lifestyle	457 (74.5)	101 (16.5)	25 (4.1)	8 (1.3)	22 (3.6)

Data presented as n (%)

Table III. Factors affecting knowledge of the influence of alcohol on the development of HT

n = 521*	Significant impact**	No impact***	p
Female/male	245 (81.4)/178 (80.9)**	56 (18.6)/42 (19.1)	p = 0.888
Age < 49 years/> 49 years	93 (86.1)/330 (79.9)	15 (13.9)/83 (20.1)	P = 0.142
Married/single	321 (81.3)/102 (80.9)	74 (18.7)/24 (19.1)	p = 0.940
Rural/urban residence	199 (82.9)/224 (79.7)	41 (17.1)/57 (20.3)	p = 0.351
Primary education or vocational/secondary or higher	181 (84.6)/242 (78.8)	33 (15.4)/65 (21.2)	p = 0.098
Professionally active yes/no	182 (80.5)/241 (81.7)	44 (19.5)/54 (18.3)	p = 0.736
Knowledge from health professionals yes/no	325 (81.7)/98 (79.7)	73 (18.3)/25 (20.3)	p = 0.623
Knowledge from the press, TV and newspapers, books yes/no	273 (79.8)/150 (83.8)	69 (20.2)/29 (16.2)	p = 0.270
Knowledge from family and friends yes/no	143 (84.6)/280 (79.5)	26 (15.4)/72 (20.5)	p = 0.166
Knowledge from the Internet yes/no	91 (74.0)/332 (83.4)	32 (26.0)/66 (16.6)	p = 0.019
Obesity no/yes	268 (80.1)/155 (81.6)	63 (19.0)/35 (18.4)	p = 0.860
Current smoker yes/no	66 (79.5)/357 (81.5)	17 (20.5)/81 (18.5)	p = 0.670

Data is presented as n (%); *not including people who replied "I have no opinion"; **combination of high and moderate impact; ***combination of low and no impact

Table IV. Factors affecting knowledge of the influence of coffee on the development of HT

n = 537*	Significant impact **	No impact***	p
Female/male	200 (64.1)/146 (64.9)	112 (35.9)/79 (35.1)	p = 0.850
Age < 49 years/> 49 years	78 (68.4)/268 (63.4)	36 (31.6)/155 (36.6)	p = 0.320
Married/single	261 (65.6)/85 (61.1)	137 (34.4)/54 (38.9)	p = 0.348
Rural/urban residence	164 (65.3)/182 (63.6)	87 (34.7)/104 (36.4)	p = 0.681
Primary education or vocational/secondary or higher	146 (65.8)/200 (63.5)	76 (34.2)/115 (36.5)	p = 0.588
Professionally active yes/no	155 (68.0)/191 (61.8)	73 (32.0)/118 (38.2)	p = 0.140
Knowledge from health professionals yes/no	273 (66.6)/73 (57.5)	137 (33.4)/54 (43.5)	p = 0.061
Knowledge from the press, TV and newspapers, books yes/no	220 (62.7)/126 (67.7)	131 (37.3)/60 (32.3)	p = 0.243
Knowledge from family and friends yes/no	122 (70.9)/224 (61.4)	50 (29.1)/141 (38.6)	p = 0.031
Knowledge from the Internet yes/no	80 (64.0)/266 (64.6)	45 (36.0)/146 (35.4)	p = 0.908
Obesity no/yes	220 (65.9)/126 (62.1)	114 (34.1)/77 (37.9)	p = 0.373
Current smoker yes/no	53 (60.9)/293 (65.1)	34 (39.1)/157 (34.9)	p = 0.455

Data is presented as n (%); *not including people who replied "I have no opinion"; **combination of high and moderate impact; ***combination of low and no impact

Table V. Factors affecting knowledge of the influence of cigarette smoking on the development of HT

n = 535*	Significant impact**	No impact***	p
Female/male	255 (84.7)/195 (83.3)	46 (15.3)/39 (16.7)	p = 0.664
Age < 49 years/> 49 years	94 (82.5)/356 (84.6)	20 (17.5)/65 (15.4)	p = 0.586
Married/single	336 (83.8)/114 (85.0)	65 (16.2)/20 (14.9)	p = 0.725
Rural/urban residence	212 (85.1)/238 (83.2)	37 (14.9)/48 (16.8)	P = 0.544
Primary education or vocational/secondary or higher	189 (85.5)/261 (83.1)	32 (14.5)/53 (16.9)	p = 0.455
Professionally active yes/no	188 (81.7)/262 (85.9)	42 (18.3)/43 (14.1)	p = 0.192
Knowledge from health professionals yes/no	352 (86.5)/98 (76.6)	55 (13.5)/30 (23.4)	p = 0.007
Knowledge from the press, TV and newspapers, books yes/no	292 (84.9)/158 (82.7)	52 (15.1)/33 (17.3)	p = 0.512
Knowledge from family and friends yes/no	148 (8.6)/302 (82.5)	21 (12.4)/64 (17.5)	p = 0.137
Knowledge from the Internet yes/no	101 (81.4)/349 (84.9)	23 (18.6)/62 (15.1)	p = 0.355
Obesity no/yes	274(82.5)/176 (86.7)	58 (17.5)/27 (13.3)	p = 0.201
Current smoker yes/no	71(77.2)/379 (85.5)	21 (22.8)/64 (14.5)	p = 0.045

Data is presented as n (%); *not including people who replied "I have no opinion"; **combination of high and moderate impact; ***combination of low and no impact

Table VI. Factors affecting knowledge of the influence of large amounts of salt on the development of HT

n = 586*	Significant impact**	No impact***	p
Female/male	314 (93.4)/228 (91.2)	22 (6.6)/22 (8.8)	p = 0.306
Age < 49 years/> 49 years	111 (94.1)/431 (92.1)	7 (5.9)/37 (7.9)	P = 0.467
Married/single	404 (92.9)/138 (91.4)	31 (7.1)/13 (8.6)	p = 0.551
Rural/urban residence	264 (93.3)/278 (91.7)	19 (6.7)/25 (8.3)	p = 0.480
Primary education or vocational/secondary or higher	232 (93.1)/310 (92.0)	17 (6.8)/27 (8.0)	p = 0.591
Professionally active yes/no	228 (93.1)/314 (92.1)	17 (6.9)/27 (7.9)	p = 0.657
Knowledge from health professionals yes/no	414 (93.7)/128 (88.9)	28 (6.3)/16 (11.1)	p = 0.059
Knowledge from the press, TV and newspapers, books yes/no	355 (93.7)/187 (90.3)	24 (6.3)/20 (9.7)	p = 0.144
Knowledge from family and friends yes/no	176 (92.1)/366 (92.7)	15 (7.9)/29 (7.3)	p = 0.826
Knowledge from the Internet yes/no	125 (94.7)/417 (91.8)	7 (5.3)/37 (8.2)	p = 0.275
Obesity no/yes	330 (92.2)/212 (93.0)	28 (7.8)/16 (7.0)	p = 0.719
Current smoker yes/no	82 (89.1)/460 (93.1)	10 (10.9)/34 (6.9)	p = 0.183

Data is presented as n (%); *not including people who replied "I have no opinion"; **combination of high and moderate impact; ***combination of low and no impact

Table VII. Factors affecting knowledge of the influence of large amounts of fat on the development of HT

n = 571*	Significant impact**	No impact***	p
Female/male	317 (96.1)/222 (92.1)	13 (3.9)/19 (7.9)	p = 0.043
Age < 49 years/> 49 years	112 (95.7)/427 (94.0)	5 (4.3)/27 (6.0)	p = 0.483
Married/single	403 (96.2)/136 (95.1)	25 (5.8)/7 (4.9)	p = 0.670
Rural/urban residence	255 (94.1)/284 (94.7)	16 (5.9)/16 (5.3)	p = 0.767
Primary education or vocational/secondary or higher	224 (93.3)/315 (95.2)	16 (6.7)/16 (4.8)	p = 0.347
Professionally active yes/no	229 (95.0)/310 (93.9)	12 (5.0)/20 (6.1)	p = 0.579
Knowledge from health professionals yes/no	418 (96.1)/121 (89.0)	17 (3.9)/15 (11.0)	p = 0.002
Knowledge from the press, TV and newspapers, books yes/no	352 (94.4)/187 (94.4)	21 (5.6)/11 (5.6)	p = 0.971
Knowledge from family and friends yes/no	174 (93.0)/365 (95.0)	13 (7.0)/19 (5.0)	p = 0.329
Knowledge from the Internet yes/no	128 (97.0)/411 (93.6)	4 (3.0)/28 (6.4)	p = 0.143
Obesity no/yes	326 (93.4)/213 (95.9)	23 (6.6)/9 (4.1)	p = 0.199
Current smoker yes/no	83 (96.5)/456 (94.0)	3 (3.5)/29 (6.0)	p = 0.355

Data is presented as n (%); *not including people who replied "I have no opinion"; **combination of high and moderate impact; ***combination of low and no impact

Table VIII. Factors affecting knowledge of the influence of irregular diet on the development of HT

n = 465*	Significant impact**	No impact***	p
Female/male	164 (62.8)/120 (58.8)	97 (37.2)/84 (41.2)	p = 0.379
Age < 49 years/> 49 years	71 (68.9)/213 (58.8)	32 (31.1)/149 (41.2)	p = 0.064
Married/single	210 (60.0)/74 (64.3)	140 (40.0)/41 (35.7)	p = 0.407
Rural/urban residence	124 (57.7)/160 (64.0)	91 (42.3)/90 (36.0)	p = 0.163
Primary education or vocational/secondary or higher	107 (56.3)/177 (64.4)	83 (43.7)/98 (35.6)	p = 0.080
Professionally active yes/no	134 (66.0)/150 (57.2)	69 (34.0)/112 (42.8)	p = 0.055
Knowledge from health professionals yes/no	232 (64.3)/52 (50.0)	129 (35.7)/52 (50.0)	p = 0.009
Knowledge from the press, TV and newspapers, books yes/no	177 (60.2)/107 (62.6)	117 (39.8)/64 (37.4)	p = 0.613
Knowledge from family and friends yes/no	101 (68.7)/183 (57.5)	46 (31.3)/135 (42.5)	p = 0.022
Knowledge from the Internet yes/no	75 (69.4)/209 (58.5)	33 (30.6)/148 (41.5)	p = 0.042
Obesity no/yes	170 (60.1)/114 (62.6)	113 (39.9)/68 (37.4)	p = 0.580
Current smoker yes/no	46 (58.2)/238 (61.7)	33 (41.8)/148 (38.3)	p = 0.569

Data is presented as n (%); *not including people who replied "I have no opinion"; **combination of high and moderate impact; ***combination of low and no impact

Table IX. Factors affecting knowledge of the influence of small amounts of fibre and vegetables and skimmed dairy products on the development of HT

n = 432*	Significant impact**	No impact***	p
Female/male	166 (63.1)/96 (56.8)	97 (36.9)/73 (43.2)	p = 0.190
Age < 49 years/> 49 years	55 (63.9)/207 (59.8)	31 (36.1)/139 (40.2)	p = 0.483
Married/single	193 (59.6)/69 (63.9)	131 (40.4)/39 (36.1)	p = 0.426
Rural/urban residence	120 (59.1)/142 (62.0)	83 (40.9)/87 (38.0)	p = 0.539
Primary education or vocational/secondary or higher	100 (60.6)/162 (60.7)	65 (39.4)/105 (39.3)	p = 0.989
Professionally active yes/no	121 (61.4)/141 (60.0)	76 (38.6)/94 (40.0)	p = 0.763
Knowledge from health professionals yes/no	202 (62.9)/60 (54.0)	119 (37.1)/51 (46.0)	p = 0.099
Knowledge from the press, TV and newspapers, books yes/no	184 (62.8)/78 (56.1)	109 (37.2)/61 (43.9)	p = 0.184
Knowledge from family and friends yes/no	87 (64.9)/175 (58.7)	47 (35.1)/123 (41.3)	p = 0.222
Knowledge from the Internet yes/no	64 (58.7)/198 (61.3)	45 (41.3)/125 (38.7)	p = 0.633
Obesity no/yes	148 (58.0)/114 (64.4)	107 (42.0)/63 (35.6)	p = 0.183
Current smoker yes/no	36 (52.9)/226 (62.1)	32 (47.1)/138 (37.9)	p = 0.156

Data is presented as n (%); *not including people who replied "I have no opinion"; **combination of high and moderate impact; ***combination of low and no impact

often, people living in rural areas (6.6% vs. 2.6%; p = 0.019) had incorrect information (Tables II and X).

In the study population, as many as 86.9% of the respondents considered high cholesterol to have an impact on the development of HT. People who received information from family and friends (97.2% vs. 92.5%; p = 0.028) were more likely to have incorrect information (Tables II and XI).

In the case of low physical activity, 11.2% of the respondents stated that it did not influence the development of HT. Especially the rural population (15.6% vs. 9.1%; p = 0.019), people with basic or vocational education (16.7% vs. 9.1%; p = 0.007), those not receiving information from health profes-

sionals (17.7 vs. 10.4%; p = 0.021) had incorrect information (Tables II and XII).

Among the respondents, 5.4% had a misconception about the impact of stressful lifestyle on the HT, and this situation was more common among men (7.8% vs. 3.9%; p = 0.041) and those who did not receive information from health professionals (10.6% vs. 3.9%; p = 0.002) (Tables II and XIII).

Discussion

Effective HT treatment includes not only pharmacotherapy, but also involving the patient in the elim-

Table X. Factors affecting knowledge of the influence of overweight and obesity on the development of HT

n = 595*	Significant impact**	No impact***	p
Female/male	326 (96.4)/242 (94.2)	12 (3.6)/15 (5.8)	p = 0.184
Age < 49 years/> 49 years	114 (95.8)/454 (95.4)	5 (4.2)/22 (4.6)	p = 0.844
Married/single	423 (95.1)/145 (96.7)	22 (4.9)/5 (3.3)	p = 0.412
Rural/urban residence	269 (93.4)/299 (97.4)	19 (6.6)/8 (2.6)	p = 0.019
Primary education or vocational/secondary or higher	239 (94.5)/329 (96.2)	14 (5.5)/13 (3.8)	p = 0.315
Professionally active yes/no	237 (95.6)/331 (95.4)	11 (4.4)/16 (4.6)	p = 0.919
Knowledge from health professionals yes/no	431 (96.4)/137 (92.6)	16 (3.6)/11 (7.4)	p = 0.051
Knowledge from the press, TV and newspapers, books yes/no	370 (96.1)/198 (94.3)	15 (3.9)/12 (5.7)	p = 0.309
Knowledge from family and friends yes/no	184 (95.3)/384 (95.5)	9 (4.7)/18 (4.5)	p = 0.919
Knowledge from the Internet yes/no	129(97.0)/439(95.0)	4(3.0)/23(5.0)	p = 0.336
Obesity no/yes	340(94.7)/228(96.6)	19(5.3)/8(3.4)	p = 0.275
Current smoker yes/no	86(93.5)/482(95.8)	6(6.5)/21(4.2)	p = 0.320

Data is presented as n (%); *not including people who replied "I have no opinion"; **combination of high and moderate impact; ***combination of low and no impact

Table XI. Factors affecting knowledge of the influence of high cholesterol on the development of HT

n = 567*	Significant impact**	No impact***	p
Female/male	311 (94.5)/222 (93.3)	18 (5.5)/16 (6.7)	p = 0.536
Age < 49 years/> 49 years	109 (93.2)/424 (94.2)	8 (6.8)/26 (5.8)	p = 0.667
Married/single	402 (93.9)/131 (94.2)	26 (6.1)/8 (5.8)	p = 0.890
Rural/urban residence	255 (94.1)/278 (93.9)	16 (5.9)/18 (6.1)	P = 0.929
Primary education or vocational/secondary or higher	219 (92.8)/314 (94.9)	17 (7.2)/17 (5.1)	p = 0.307
Professionally active yes/no	220 (93.2)/313 (94.6)	16 (6.8)/18 (5.4)	p = 0.507
Knowledge from health professionals yes/no	408 (94.7)/125 (91.9)	23 (5.3)/11 (8.1)	p = 0.239
Knowledge from the press, TV and newspapers, books yes/no	345 (94.0)/188 (94.0)	22 (6.0)/12 (6.0)	p = 0.998
Knowledge from family and friends yes/no	175 (97.2)/358 (92.5)	5 (2.8)/29 (7.5)	p = 0.028
Knowledge from the Internet yes/no	122 (94.6)/411 (93.8)	7 (5.4)/27 (6.2)	p = 0.756
Obesity no/yes	325 (93.7)/208 (94.5)	22 (6.3)/12 (5.5)	p = 0.665
Current smoker yes/no	80(89.9)/453(94.8)	9(10.1)/25(5.2)	p = 0.075

Data is presented as n (%); *not including people who replied "I have no opinion"; **combination of high and moderate impact; ***combination of low and no impact

Table XII. Factors affecting knowledge of the influence of low physical activity on the development of HT

n = 565*	Significant impact**	No impact***	p
Female/male	283 (87.9)/213 (87.6)	39 (12.1)/30 (12.4)	p = 0.933
Age < 49 years/> 49 years	104 (89.7)/392 (87.3)	12 (10.3)/57 (12.7)	p = 0.491
Married/single	381 (89.2)/115 (83.3)	46 (10.8)/23 (16.7)	p = 0.066
Rural/urban residence	227 (84.4)/269 (90.9)	42 (15.6)/27 (9.1)	p = 0.019
Primary education or vocational/secondary or higher	195 (83.3)/301 (90.9)	39 (16.7)/30 (9.1)	p = 0.007
Professionally active yes/no	211 (87.5)/285 (88.0)	30 (12.5)/39 (12.0)	p = 0.883
Knowledge from health professionals yes/no	380 (89.6)/116 (82.3)	44 (10.4)/25 (17.7)	p = 0.021
Knowledge from the press, TV and newspapers, books yes/no	327 (88.4)/169 (86.7)	43 (11.6)/26 (13.3)	p = 0.555
Knowledge from family and friends yes/no	162 (88.5)/334 (87.4)	21 (11.5)/48 (12.6)	p = 0.711
Knowledge from the Internet yes/no	121 (91.7)/375 (86.6)	11 (8.3)/58 (13.4)	P = 0.120
Obesity no/yes	307 (86.7)/189 (89.6)	47 (13.3)/22 (10.4)	p = 0.317
Current smoker yes/no	74 (82.2)/422 (88.8)	16 (17.8)/53 (11.2)	p = 0.079

Data is presented as n (%); *not including people who replied "I have no opinion"; **combination of high and moderate impact; ***combination of low and no impact

Table XIII. Factors affecting knowledge of the influence of stressful lifestyle on the development of HT

n = 591*	Significant impact**	No impact***	p
Female/male	321 (96.1)/237 (92.2)	13 (3.9)/20 (7.8)	p = 0.041
Age <49 years/> 49 years	109 (94.0)/449 (94.5)	7 (6.0)/26 (5.5)	p = 0.814
Married/single	419 (94.6)/139 (93.9)	24 (5.4)/9 (6.1)	P = 0.761
Rural/urban residence	263 (92.9)/295 (95.8)	20 (7.1)/13 (4.2)	p = 0.132
Primary education or vocational/secondary or higher	230 (92.7)/328 (95.6)	18 (7.3)/15 (4.4)	p = 0.132
Professionally active yes/no	233 (94.3)/325 (94.5)	14 (5.7)/19 (5.5)	p = 0.940
Knowledge from health professionals yes/no	423 (96.1)/135 (89.4)	17 (3.9)/16 (10.6)	p = 0.002
Knowledge from the press, TV and newspapers, books yes/no	361 (93.5)/197 (96.1)	25 (6.5)/8 (3.9)	p = 0.195
Knowledge from family and friends yes/no	185 (95.4)/373 (93.9)	9 (4.6)/24 (6.1)	p = 0.484
Knowledge from the Internet yes/no	125 (94.0)/433 (94.5)	8 (6.0)/25 (5.5)	p = 0.806
Obesity no/yes	339 (94.2)/219 (94.8)	21 (5.8)/12 (5.2)	p = 0.741
Current smoker yes/no	85 (92.4)/473 (94.8)	7 (7.6)/26 (5.2)	p = 0.357

Data is presented as n (%); *not including people who replied "I have no opinion"; **combination of high and moderate impact; ***combination of low and no impact

ination of the exacerbating factors, which in turn requires them to have the appropriate knowledge.

Several decades ago, the adverse effects of alcohol on the RR in patients treated with HT were confirmed. Limiting alcohol intake is associated with reduced need for antihypertensive drugs and better control of RR [4, 5]. Unfortunately, false information or lack of knowledge in this area was observed in about 1/3 of respondents, 16% and 15%, respectively. In fact, people who got it from the Internet were more likely to have false information. Therefore, it seems appropriate to spread knowledge more widely, especially on websites.

Drinking coffee systematically does not affect the development of HT, but it can contribute to reducing mortality due to cardiovascular disease, lung diseases, diabetes, traumas and infections [17, 21]. Unfortunately, more than 55% of people in the study had incorrect knowledge of the subject. This phenomenon especially affected patients who received information from family and friends, the group that seems to be the most difficult to reach in social campaigns.

So far no direct connection has been proven between smoking and HT development, despite its association with mortality due to cardiovascular disease [15, 16]. In the study group, nearly three quarters believed that nicotine was associated with hypertension. This situation was more frequent in people who currently smoke cigarettes and seek information from health professionals. Perhaps HT patients are informed in health care facilities about the effects of smoking on the increased risk of cardiovascular disease and therefore they answered incorrectly more often about its effect on the RR.

The effect of consuming excessive amounts of salt on HT development has been frequently proven. Even the value of the decrease in RR when using a low sodium diet was estimated [5]. The respondents presented relatively good knowledge about the effect of salt intake on HT. A little over 7% gave a wrong answer and less than 4.5% did not have any opinion about this connection. Patients who did not receive information from health professionals were reported to have incorrect information slightly more often.

So far, many studies have examined the effects of individual dietary components on cardiovascular morbidity. Recent studies have shown that fat reduction alone did not affect HT development [20]. Of the respondents, 88% had false knowledge about the effect of high intake of fat on HT development, and nearly 7% did not have opinion about this effect. This was the worst result in the study. Women and patients who were informed by health professionals responded incorrectly more often. Likewise, obese patients demonstrated a lower level of knowledge [23].

The general opinion is that irregular eating can lead to obesity and many diseases associated with it, i.e. HT. No evidence has yet been found in clinical trials of the direct effect of irregular meals on the development of HT. In this study, incorrect knowledge about the effects of irregular eating on HT development was reported by over 46% of respondents, and more than 24% did not comment on this. More often, patients who received information from health professionals, family and friends, and the Internet were misinformed.

Changing diet can significantly reduce HT development. In the studies published in 1997 and 2000, the effects of a diet containing fibre, vegetables and

skimmed dairy products on the reduction of RR have been confirmed. Nearly 30% of the respondents did not comment on the impact of the aforementioned diet on HT development and nearly 28% had misleading information. Slightly more often, people who did not receive information from health professionals answered incorrectly. The above data show poor knowledge of the correct eating habits irrespective of sociodemographic characteristics of patients.

One element of a healthy lifestyle that can reduce RR and therefore antihypertensive drugs intake is to lose weight in overweight or obese patients [9]. Only 4.4% of respondents mistakenly answered the question about the connection between HT and excess weight. This was the best result in the poll. People living in rural areas, as compared to urban dwellers, had misleading information more often.

Until now, there is no convincing evidence for the connection between higher cholesterol levels and other lipid disorders on the development of HT. Despite this, almost 88% of respondents gave an incorrect answer about the connection between the two disorders. This was one of the two worst results in the study. Especially those who obtained information from family and friends had incorrect knowledge.

Regular exercise exerts a positive influence on cardiovascular risk factors including RR, and in patients with HT it allows the reduction of antihypertensives [5, 10]. In the surveyed population, more than 11% of the respondents had a false belief about the insignificance of low physical activity on the development of HT, and almost 8% had no opinion on it. Especially those living in the countryside, with primary or vocational education, not receiving information from healthcare professionals, gave an incorrect answer.

A cause-and-effect relationship between stressful lifestyle and HT and ischaemic heart disease has been proven [12]. As many as 91% of respondents gave the correct answer, which was the second result in this study. Incorrect knowledge concerned especially men and those not receiving information from health professionals.

Conclusions

1. Incorrect knowledge of factors affecting the development of HT concerned in particular the consumption of excessive amounts of fat, raised cholesterol, nicotine/smoking and drinking coffee.
2. There were no common features of people who answered the questions incorrectly.

3. Results of the study indicate the need to educate people with HT on the factors influencing the development of this disease.

References

1. Kearney PM, Whelton M, Reynolds K, et al. Global burden of hypertension: analysis of worldwide data. *Lancet*. 2005; 365(9455): 217–223, doi: [10.1016/S0140-6736\(05\)17741-1](https://doi.org/10.1016/S0140-6736(05)17741-1), indexed in Pubmed: [15652604](https://pubmed.ncbi.nlm.nih.gov/15652604/).
2. Zdrojewski Ł, Zdrojewski T, Rutkowski M, et al. Prevalence and control of cardiovascular risk factors in Poland. Assumptions and objectives of the NATPOL 2011 Survey. *Kardiol Pol*. 2013; 71(4): 381–392, doi: [10.5603/KP.2013.0066](https://doi.org/10.5603/KP.2013.0066), indexed in Pubmed: [23788344](https://pubmed.ncbi.nlm.nih.gov/23788344/).
3. Lalonde M. A New Perspective on the Health of Canadians, a working document. : 1974.
4. Puddey IB, Beilin LJ, Vandongen R. Regular alcohol use raises blood pressure in treated hypertensive subjects. A randomised controlled trial. *Lancet*. 1987; 1(8534): 647–651, indexed in Pubmed: [2882082](https://pubmed.ncbi.nlm.nih.gov/2882082/).
5. Dickinson HO, Mason JM, Nicolson DJ, et al. Lifestyle interventions to reduce raised blood pressure: a systematic review of randomized controlled trials. *J Hypertens*. 2006; 24(2): 215–233, doi: [10.1097/01.hjh.0000199800.72563.26](https://doi.org/10.1097/01.hjh.0000199800.72563.26), indexed in Pubmed: [16508562](https://pubmed.ncbi.nlm.nih.gov/16508562/).
6. Rupp H, Brilla CG, Maisch B. [Hypertension and alcohol: central and peripheral mechanisms]. *Herz*. 1996; 21(4): 258–264, indexed in Pubmed: [8805006](https://pubmed.ncbi.nlm.nih.gov/8805006/).
7. Reisin E. Sodium and obesity in the pathogenesis of hypertension. *Am J Hypertens*. 1990; 3(2): 164–167, indexed in Pubmed: [2155002](https://pubmed.ncbi.nlm.nih.gov/2155002/).
8. Wofford MR, Hall JE. Pathophysiology and treatment of obesity hypertension. *Curr Pharm Des*. 2004; 10(29): 3621–3637, indexed in Pubmed: [15579059](https://pubmed.ncbi.nlm.nih.gov/15579059/).
9. Neter JE, Stam BE, Kok FJ, et al. Influence of weight reduction on blood pressure: a meta-analysis of randomized controlled trials. *Hypertension*. 2003; 42(5): 878–884, doi: [10.1161/01.HYP.0000094221.86888.AE](https://doi.org/10.1161/01.HYP.0000094221.86888.AE), indexed in Pubmed: [12975389](https://pubmed.ncbi.nlm.nih.gov/12975389/).
10. Cornelissen VA, Fagard RH. Effects of endurance training on blood pressure, blood pressure-regulating mechanisms, and cardiovascular risk factors. *Hypertension*. 2005; 46(4): 667–675, doi: [10.1161/01.HYP.0000184225.05629.51](https://doi.org/10.1161/01.HYP.0000184225.05629.51), indexed in Pubmed: [16157788](https://pubmed.ncbi.nlm.nih.gov/16157788/).
11. Szczepańska-Sadowska E, Sikora M. Patofizjologia nadciśnienia tętniczego. Więcek A, Januszewicz A, Szczepańska-Sadowska E, Prejzbisz A. *Hipertensjologia, patogeneza, diagnostyka i leczenie nadciśnienia tętniczego*. Medycyna Praktyczna, Kraków. 2011; 2: 23–175.
12. Pickering TG. Mental stress as a causal factor in the development of hypertension and cardiovascular disease. *Curr Hypertens Rep*. 2001; 3(3): 249–254, indexed in Pubmed: [11353576](https://pubmed.ncbi.nlm.nih.gov/11353576/).
13. Omvik P. How smoking affects blood pressure. *Blood Pressure*. 2009; 5(2): 71–77, doi: [10.3109/08037059609062111](https://doi.org/10.3109/08037059609062111).
14. James JE. Critical review of dietary caffeine and blood pressure: a relationship that should be taken more seriously. *Psychosom Med*. 2004; 66(1): 63–71, indexed in Pubmed: [14747639](https://pubmed.ncbi.nlm.nih.gov/14747639/).
15. Doll R, Peto R, Boreham J, et al. Mortality in relation to smoking: 40 years' observations on male British doctors. *BMJ*. 1994; 309(6959): 901–911, indexed in Pubmed: [7755693](https://pubmed.ncbi.nlm.nih.gov/7755693/).
16. Primates P, Falaschetti E, Gupta S, et al. Association between smoking and blood pressure: evidence from the health survey for England. *Hypertension*. 2001; 37(2): 187–193, indexed in Pubmed: [11230269](https://pubmed.ncbi.nlm.nih.gov/11230269/).
17. Steffen M, Kuhle C, Hensrud D, et al. The effect of coffee consumption on blood pressure and the development of hypertension: a systematic review and meta-analysis. *J Hypertens*. 2012; 30(12): 2245–2254, doi: [10.1097/HJH.0b013e3283588d73](https://doi.org/10.1097/HJH.0b013e3283588d73), indexed in Pubmed: [23032138](https://pubmed.ncbi.nlm.nih.gov/23032138/).
18. Appel LJ, Moore TJ, Obarzanek E, et al. A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research

- Group. *N Engl J Med.* 1997; 336(16): 1117–1124, doi: [10.1056/NEJM199704173361601](https://doi.org/10.1056/NEJM199704173361601), indexed in Pubmed: [9099655](https://pubmed.ncbi.nlm.nih.gov/9099655/).
19. Conlin PR, Chow D, Miller ER, et al. The effect of dietary patterns on blood pressure control in hypertensive patients: results from the Dietary Approaches to Stop Hypertension (DASH) trial. *Am J Hypertens.* 2000; 13(9): 949–955, indexed in Pubmed: [10981543](https://pubmed.ncbi.nlm.nih.gov/10981543/).
 20. Allison MA, Aragaki AK, Ray RM, et al. A Randomized Trial of a Low-Fat Diet Intervention on Blood Pressure and Hypertension: Tertiary Analysis of the WHI Dietary Modification Trial. *Am J Hypertens.* 2016; 29(8): 959–968, doi: [10.1093/ajh/hpv196](https://doi.org/10.1093/ajh/hpv196), indexed in Pubmed: [26708006](https://pubmed.ncbi.nlm.nih.gov/26708006/).
 21. Freedman ND, Park Y, Abnet CC, et al. Association of coffee drinking with total and cause-specific mortality. *N Engl J Med.* 2012; 366(20): 1891–1904, doi: [10.1056/NEJMoa1112010](https://doi.org/10.1056/NEJMoa1112010), indexed in Pubmed: [22591295](https://pubmed.ncbi.nlm.nih.gov/22591295/).
 22. Gać P, Poręba R, Poręba M, et al. Znajomość przyczyn nadciśnienia tętniczego i skutków jego nieleczenia w losowej grupie studentów wybranych miast Polski. *Nadciśnienie Tętnicze.* 2013; 17(5): 377–383.
 23. Bronkowska M, Martynowicz H, Żmich K, et al. Wybrane elementy stylu życia oraz wiedza żywieniowa otyłych osób z rozpoznaniem nadciśnieniem tętniczym. *Nadciśnienie Tętnicze.* 2009; 13(4): 266–274.
 24. Georgiopoulou VV, Kalogeropoulos AB, Raggi P, et al. Prevention, diagnosis, and treatment of hypertensive heart disease. *Cardiol Clin.* 2010; 28(4): 675–691, doi: [10.1016/j.ccl.2010.07.005](https://doi.org/10.1016/j.ccl.2010.07.005), indexed in Pubmed: [20937450](https://pubmed.ncbi.nlm.nih.gov/20937450/).
 25. Erbel R, Aboyans V, Boileau C, et al. Wytoczne ESC dotyczące rozpoznawania i leczenia chorób aorty w 2014 roku. *Kardiologia Polska.* 2014; 72(12): 1169–1252, doi: [10.5603/kp.2014.0225](https://doi.org/10.5603/kp.2014.0225).
 26. Montalescot G, Sechtem U, Achenbach S, et al. Wytoczne ESC dotyczące postępowania w stabilnej chorobie wieńcowej w 2013 roku. *Kard Pol.* 2013; 71(9): 224–317.
 27. Mancia G, Fagard R, Narkiewicz K, et al. Wytoczne ESH/ESC dotyczące postępowania w nadciśnieniu tętniczym w 2013 roku. *Kard Pol.* 2013; 71(3): 60–64.
 28. Kirchhof P, Benussi S, Kotecha D, et al. Wytoczne ESC dotyczące leczenia migotania przedsionków w 2016 roku, opracowane we współpracy z EACTS. *Kardiologia Polska.* 2016; 74(12): 1359–1469, doi: [10.5603/kp.2016.0172](https://doi.org/10.5603/kp.2016.0172).
 29. Szczepańska-Sadowska E, Sikora M. Nadciśnienie tętnicze pierwotne. Więcek A., Januszewicz A., Szczepańska-Sadowska E., Prejbisz A. *Hipertensjologia, patogeneza, diagnostyka i leczenie nadciśnienia tętniczego.* Medycyna Praktyczna, Kraków. 2011; 2: 291–394.