

Endothelial dysfunction: Can it differentiate Takotsubo syndrome from spontaneous coronary artery dissection?

Alicja Skrobucha^{1*}, Ewa Borowiak^{1*}, Łukasz Szarpak^{2, 3},
 Krzysztof J. Filipiak^{4, 5}, Aleksandra Gąsecka¹

¹st Chair and Department of Cardiology, Medical University of Warsaw, Poland

²Institute of Outcomes Research, Maria Skłodowska-Curie Medical Academy, Warsaw, Poland

³Henry JN Taub Department of Emergency Medicine, Baylor College of Medicine, Houston, United States

⁴Institute of Clinical Science, Maria Skłodowska-Curie Medical Academy, Warsaw, Poland

⁵Department of Hypertensiology, Angiology and Internal Medicine,
 Poznan University of Medical Sciences, Poznan, Poland

It was with interest that the present authors read an article by Victoria L. Cammann et al. [1], who compared clinical features and outcomes in patients with Takotsubo syndrome (TTS) and spontaneous coronary artery dissection (SCAD). The study compared the clinical characteristics and outcomes of 2098 TTS patients from the International Takotsubo Registry to 750 SCAD patients from the Canadian Spontaneous Coronary Artery Dissection Cohort Study. Both TTS and SCAD are increasingly recognized as non-atherosclerotic causes of myocardial infarction (MI) and have a predilection for the female gender. The authors found that TTS patients were significantly older (66.9 ± 12.8 years vs. 51.8 ± 10.2 years; $p < 0.001$), had more major cardiovascular risk factors, and had a more impaired left ventricular ejection fraction than SCAD patients. On admission, TTS patients more often had dyspnoea (46.8% vs. 0.9%; $p < 0.001$), but less frequently presented with chest pain (74.4% vs. 91.5%; $p < 0.001$). Thirty-day mortality was significantly higher in the TTS group (4.4% vs. 0.1%; $p < 0.001$). This study gave noteworthy findings about the unique populations of patients with TTS and SCAD. SCAD relates to the spontaneous formation of intra-mural

hematoma within the tunica media of a coronary artery. Two pathophysiological theories have been described: the “inside-out” proposes that primarily a disruption in the vessel wall develops and allows blood to enter and generate a false lumen and in the “outside-in” the hematoma arises de novo from the vasa vasorum [2]. As the exact pathophysiology of both conditions remains unclear, here we review the literature in search of the role of endothelial dysfunction in the pathogenesis of TTS, including (i) catecholamine-induced myocardial injury, (ii) failure of coronary vasomotor function, (iii) role of estrogens in TTS pathophysiology, (iv) role of oxidative stress in TTS pathophysiology (Fig. 1).

Catecholamine-induced myocardial injury is the most established theory of TTS pathogenesis, but it does not explain all aspects of the clinical presentations of this condition. Catecholamines seem to play a central role in the pathophysiology of TTS, as acute episodes are often caused by sudden stress, intuitively associated with an endogenous adrenergic surge. Therefore, TTS is also known as stress cardiomyopathy or broken heart syndrome. Interestingly, in Cammann et al. study [1], physical triggers were more common in TTS patients (36.1% vs. 28.8%; $p < 0.001$), while

Address for correspondence: Aleksandra Gąsecka, MD, PhD, 1st Chair and Department of Cardiology, Medical University of Warsaw, ul. Banacha 1a, 02–097 Warszawa, Poland, tel: +48 22 599 19 51, e-mail: aleksandra.gasecka@wum.edu.pl

Received: 4.05.2023

Accepted: 6.10.2023

Early publication date: 16.10.2023

*Alicja Skrobucha and Ewa Borowiak equally contributed to the manuscript and share the first authorship.

This article is available in open access under Creative Commons Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.




Figure 1. Role of endothelial dysfunction in the pathogenesis of Takotsubo syndrome; A — adrenaline; NA — noradrenaline; NO — nitric oxide.

emotional triggers (50.3% vs. 29.6%; $p < 0.001$) and anxiety disorders (19.7% vs. 6.7%; $p < 0.001$) were more common in SCAD patients. There are three main aspects of catecholamine-induced myocardial injury: (i) the release of catecholamines, (ii) the cardiac response, and (iii) results in the left ventricular dysfunction. The levels of plasma catecholamine were even three times higher in patients with TTS than in patients with acute MI [3]. Mild to moderate elevations of noradrenaline and normetanephrine concentrations were found in 48% and 35% of TTS patients, respectively [4]. TTS is presumably caused by a catecholaminergic wave, leading to myocardial stunning through an action on β_2 -receptors, which hyperactivity negatively influences endothelial function. It can be detected based on decreased brachial flow mediated dilation.

Failure of coronary vasomotor function is proposed to be another explanation of TTS, including microvascular spasm and epicardial spasm. Endothelin-1 represents one of the most potent vasoconstrictors and mediators of microvascular spasm in TTS. A significant increase of endothelin-1 was observed in patients with TTS [5]. In addition, women with a history of TTS showed an

impaired endothelium-dependent vasodilation and excessive vasoconstriction, compared to patients with MI [6]. In the Kurisu et al. study [6], epicardial coronary spasm by acetylcholine infusion was induced in 23% of TTS patients. Several studies confirmed abnormal coronary vasomotor responses to acetylcholine in TTS patients.

Takotsubo syndrome mostly develops in postmenopausal women, who have a reduced level of estrogen. The effect of estrogen supplementation upregulates cardioprotective substrates such as atrial natriuretic peptide and heat shock protein 70. Furthermore, estrogen inhibits sympathetic nervous system activity and induces endothelial nitric oxide (NO) production [7]. Increased sympathetic vasoconstriction and impaired vasodilation affect the inadequate response to increased levels of psychological stress, and are also associated with reduced estrogen levels [8]. This may also be related to the role of estrogen as a factor in increasing the sensitivity of α -receptors to catecholamines, and thus promoting vasodilation. In postmenopausal women, the sensitivity of β -receptors to catecholamines is reduced, while sensitivity of α -receptors remains unchanged, resulting in a predominance of vasoconstriction over vasodilation [8].

Nitric oxide-related oxidative stress is another factor associated with the development of TTS. Several studies investigated the levels of homoarginine, a substrate for NO synthase, which is an important marker in cardiovascular disease. Homoarginine levels were significantly lower in patients with TTS compared to healthy controls (median 1403 nmol/L vs. 1634 nmol/L; $p = 0.031$). In contrast, 3-nitrotyrosine plasma concentrations, a biomarker of NO-dependent oxidative stress, showed no significant differences between groups (median 1915 pmol/L vs. 2170 pmol/L; $p = 0.4726$) [9]. Kayacelebi et al. [9] confirmed the results of previous studies showing that homoarginine is a marker of cardiovascular disease and that oxidative stress has an influence on the pathomechanism of TTS.

Considering L-arginine, another substrate for NO synthase, a shift toward a competitive metabolic pathway involving arginase correlated with the occurrence of thin-cap fibroatheroma, that is a feature of acute coronary syndrome, and with thickening of the intima-media in the chronic phase [10].

Understanding the underlying etiology of TTS seems critical from a clinical point of view to develop appropriate treatments for both the acute phase and the prevention of long-term recurrent events. It may also bring us closer to discovering biomarkers for diagnosis and risk prediction. Endothelial dysfunction appears to play an important role in the pathogenesis of TTS. As Cammann et al. [1] showed, differences exist between the populations of patients with TTS and SCAD. However, this study did not evaluate the potential differences in endothelial dysfunction in both diseases. Hence, it would be interesting to investigate the role of endothelial dysfunction both in patients with TTS and SCAD, with the goal of comparing the pathogenesis of these two diseases.

Conflict of interest: None declared

References

1. Cammann VL, Sarcon A, Szawan KA, et al. Clinical characteristics and outcomes of patients with takotsubo syndrome versus spontaneous coronary artery dissection. *Cardiol J*. 2023; 30(1): 125–130, doi: [10.5603/CJ.a2021.0065](https://doi.org/10.5603/CJ.a2021.0065), indexed in Pubmed: [34165180](https://pubmed.ncbi.nlm.nih.gov/34165180/).
2. Hayes SN, Tweet MS, Adlam D, et al. Spontaneous coronary artery dissection: JACC state-of-the-art review. *J Am Coll Cardiol*. 2020; 76(8): 961–984, doi: [10.1016/j.jacc.2020.05.084](https://doi.org/10.1016/j.jacc.2020.05.084), indexed in Pubmed: [32819471](https://pubmed.ncbi.nlm.nih.gov/32819471/).
3. Wittstein IS, Thiemann DR, Lima JAC, et al. Neurohumoral features of myocardial stunning due to sudden emotional stress. *N Engl J Med*. 2005; 352(6): 539–548, doi: [10.1056/NEJMoa043046](https://doi.org/10.1056/NEJMoa043046), indexed in Pubmed: [15703419](https://pubmed.ncbi.nlm.nih.gov/15703419/).
4. Y-Hassan S, Henareh L. Plasma catecholamine levels in patients with takotsubo syndrome: Implications for the pathogenesis of the disease. *Int J Cardiol*. 2015; 181: 35–38, doi: [10.1016/j.ijcard.2014.11.149](https://doi.org/10.1016/j.ijcard.2014.11.149), indexed in Pubmed: [25481310](https://pubmed.ncbi.nlm.nih.gov/25481310/).
5. Martin EA, Prasad A, Rihal CS, et al. Endothelial function and vascular response to mental stress are impaired in patients with apical ballooning syndrome. *J Am Coll Cardiol*. 2010; 56(22): 1840–1846, doi: [10.1016/j.jacc.2010.03.107](https://doi.org/10.1016/j.jacc.2010.03.107), indexed in Pubmed: [21087714](https://pubmed.ncbi.nlm.nih.gov/21087714/).
6. Kurisu S, Sato H, Kawagoe T, et al. Tako-tsubo-like left ventricular dysfunction with ST-segment elevation: a novel cardiac syndrome mimicking acute myocardial infarction. *Am Heart J*. 2002; 143(3): 448–455, doi: [10.1067/mhj.2002.120403](https://doi.org/10.1067/mhj.2002.120403), indexed in Pubmed: [11868050](https://pubmed.ncbi.nlm.nih.gov/11868050/).
7. Pelliccia F, Greco C, Vitale C, et al. Takotsubo syndrome (stress cardiomyopathy): an intriguing clinical condition in search of its identity. *Am J Med*. 2014; 127(8): 699–704, doi: [10.1016/j.amjmed.2014.04.004](https://doi.org/10.1016/j.amjmed.2014.04.004), indexed in Pubmed: [24754972](https://pubmed.ncbi.nlm.nih.gov/24754972/).
8. Sachdev E, Bairey Merz CN, Mehta PK. Takotsubo cardiomyopathy. *Eur Cardiol*. 2015; 10(1): 25–30, doi: [10.15420/ocr.2015.10.01.25](https://doi.org/10.15420/ocr.2015.10.01.25), indexed in Pubmed: [30310419](https://pubmed.ncbi.nlm.nih.gov/30310419/).
9. Kayacelebi AA, Nguyen THa, Neil C, et al. Homoarginine and 3-nitrotyrosine in patients with takotsubo cardiomyopathy. *Int J Cardiol*. 2014; 173(3): 546–547, doi: [10.1016/j.ijcard.2014.03.080](https://doi.org/10.1016/j.ijcard.2014.03.080), indexed in Pubmed: [24704403](https://pubmed.ncbi.nlm.nih.gov/24704403/).
10. Mołek P, Żmudzki P, Machnik A, et al. Thin-cap fibroatheroma and increased coronary intima-media thickness are associated with an altered balance of arginine metabolites. *Pol Arch Intern Med*. 2023; 133(2), doi: [10.20452/pamw.16356](https://doi.org/10.20452/pamw.16356), indexed in Pubmed: [36226830](https://pubmed.ncbi.nlm.nih.gov/36226830/).

