

# Does non-pharmacological treatment affect outcomes in dysautonomic syndromes?

Denise Hachul

Cardiac Arrhythmias Unit, Heart Institute, University of Sao Paulo Medical School, Sao Paulo, Brazil

## Abstract

*Dysautonomic syndromes are very common clinical conditions with different presentations and a wide spectrum of mechanisms. Many non-pharmacological and pharmacological interventions have been recommended to treat dysautonomic patients along these years, but the majority of them were not supported by multicenter controlled trials. General measures, like water drinking and salt intake, avoiding triggers and recognizing prodromal symptoms, as well as performing orthostatic training and physical conditioning are commonly able to improve symptoms and avoid syncope and falls. In this article, we will explore why and how such measures are applied to treat patients with dysautonomic syndromes, based on the physiopathology of these disorders. (Cardiol J 2014; 21, 6: 611–615)*

**Key words:** dysautonomic syndromes, syncope

## Introduction

Dysautonomic syndromes leading to orthostatic intolerance and syncope are common clinical issues [1]. In a recent international meta-analysis, involving 43,315 patients with syncope in emergency departments, dysautonomic conditions (situational, orthostatic or vasovagal causes) were present in 30% of patients [2].

Dysautonomic syndromes can be acquired or idiopathic. Some are genetically determined. In general, these disorders have a benign prognosis, but up to one third of patients present physical injury related to falls. Thus, dysautonomic syndromes have a great impact on quality of life [3]. Lack of a robust diagnosis of the mechanisms involved and presence of frequent recurrences can create psychosocial effects including anxiety, panic, and depression. For these reasons, all efforts have to be made to clarify the etiology of syncope, define the prognosis and, consequently, establish the best treatment.

Different clinical presentations and a wide spectrum of mechanisms are involved in the

pathophysiology of these syndromes [4], most of which are still not entirely understood. While different non-pharmacological interventions have been proposed, their results are controversial and multicenter controlled trials supporting them do not exist.

Individualizing understanding of autonomic dysfunction in relationship to an organic condition and distinguishing it from a functional disorder is fundamental before planning treatment. Concerning reflex syncope, achieving an ideal treatment is even more difficult, because its paroxysmal nature makes the assessment of the treatment efficacy particularly challenging.

## Pathophysiological mechanisms as a goal for therapeutic approaches

Vasovagal syncope seems to be a physiological response with different degrees of individual susceptibility: the reflex pathways involved in vasovagal reflex are the same as those related to the mechanisms of protection against exaggerated bleeding and extreme stress and may occur in

**Address for correspondence:** Dr Denise Hachul, Dr Eneas De Carvalho Aguiar 44. ZIP Code: 05403-000. Sao Paulo, SP, Brasil, tel/fax: 55 11 26615341, e-mail: denise.hachul@gmail.com

Received: 02.11.2014

Accepted: 11.02.2014

normal individuals after variable periods of space flights or exposure to ground models of microgravity [5].

Autonomic control of the cardiovascular system depends on the integrity of the brain, spinal cord and nerves; it depends also on the integrity of the heart conduction system, on vasoconstrictor reserve, blood volume, and influences of pharmacotherapy and age. The first step of the treatment is to reassure patients about the benign nature of the condition and to teach patients how to avoid triggers and recognize premonitory symptoms in order to prevent falls and physical injuries [6].

It has been observed that in dysautonomic syndromes, most patients have greater venous volumes and impaired vasoconstriction in response to orthostatic stress and exercise [7, 8]. This condition leads to an inefficient venous return and abnormal central blood distribution. The level of vasoconstriction deficiency is associated with different degrees of orthostatic intolerance, being worse in those patients who faint early during tilt table test [9, 10].

### Salt and water

Salt supplementation has an important role for patients with dysautonomic syndromes and it is associated with improvement in orthostatic tolerance [11]. It is recommended in the absence of coexisting hypertension and mainly in patients with 24-h urinary excretion of sodium inferior to 170 mmol/dL. Salt supplementation provokes improvement of the sympathetic control of the vasculature, leading to a better vasoconstriction response to orthostatic stress and a better cerebral auto-regulation, with no change in the resting blood pressure (BP). The recommendation is to add enough salt to the meal as taste allows. If this sodium is insufficient, oral tablets can be added.

The augmented ingestion of water is recommended not only to increase blood volume. Indeed, this is not the most important consequence of the water ingestion on dysautonomic patients. When water is drunk, a pressor response is observed, reaching the maximum effect around 20 min after the ingestion [12, 13]. This consequence was demonstrated in older normal subjects and in patients with autonomic failure. The mechanisms of the pressor response remain unclear, but it has been observed that water incites an increase in the peripheral vascular resistance and consequently, less fall of the stroke volume. There is indirect evidence that water provides a pressor effect even in normal subjects. It was observed that water

ingestion increased orthostatic tolerance to head-up tilt test combined with lower body negative pressure [12]. Its effect alone in vasovagal syncope remains unclear but, in the postural orthostatic tachycardia syndrome, water ingestion attenuates the postural rise in heart rate (HR) and in patients with autonomic failure, and also reduces the degree of orthostatic and postprandial hypotension [13].

### Tilt training, head-up sleeping

Orthostatic intolerance is a common clinical syndrome observed after space flights and its ground based simulation models, the 6° head down bed rest and diving [14, 15]. In these cases, orthostatic intolerance cannot be explained as a simple consequence of hypovolemia, because restoration of plasma volume does not completely improve the condition. It is already known that microgravity provokes many modifications in the hemodynamic parameters and cardiovascular autonomic responses to orthostatic stress, like impairment in the baroreflex sensitivity, attenuation of vasoconstriction and increase of venous capacitance, which is exacerbated by a decrease in muscular tonus. These data show that similar cardio-circulatory changes are present in ground based microgravity models and in dysautonomic syndromes. Trying to clarify the role of orthostatic training in the management of these conditions, 1 study evaluated healthy subjects, who were divided into two groups. They were submitted to 21 days of head-down tilt bed rest (HDT) [16]. The study group was also submitted to lower-body negative pressure (LBNP). LBNP is an experimental model of orthostatic stress and it is considered similar to head-up tilt table test in assessing predisposition to hypotension. It can also simulate orthostatic or tilt training. Before HDT, all the subjects of the study group and the control group completed the tilt tests. After 10 and 21 days of HDT, all the subjects of the control group and 1 subject of the LBNP group could not complete the tilt test due to presyncopal or syncopal symptoms. The mean upright time before symptoms in the control group was significantly shorter than the one in the LBNP group. Stroke volume and cardiac output decreased significantly in the control group on days 3 and 10 of HDT, but remained unchanged throughout HDT in the LBNP group. The results suggest that brief daily orthostatic training sessions were effective in diminishing the consequence of HDT on orthostatic tolerance. Extrapolating the findings for dysautonomic patients, postural training could be applied

to obtain a similar effect. A practical method of performing orthostatic training is head-up sleeping (HUS) [17]. The effect of 3 to 4 months of 10° HUS on orthostatic tolerance was studied in patients with reflex syncope and early presyncope during tilting test associated with graded LBNP. Time to presyncope significantly increased after HUS. Before HUS, all patients experimented presyncope by the end of first level of lower body suction. After HUS, almost 50% of the patients were able to tolerate the test. HUS did not affect HR, whereas mean BP suffered a significant increase. Overall, there was a significant increase in plasma volume after HUS.

Ector et al. [18] proposed that repeated exposure to orthostatic stress and tilt training might be helpful to treat patients with dysautonomic syndromes, based on their potential positive effects on the autonomic function of the cardiovascular system. Experimental studies demonstrated a possible effect on the vasoconstriction reserve [19]. However, randomized controlled trials are controversial and do not confirm the effectiveness of tilt testing, especially because of poor compliance in continuing the training for longer periods [20–22].

### Exercise training

Although exercise training is highly recommended for patients with dysautonomic syncope, there are few published data demonstrating its benefits in improving orthostatic tolerance [6].

It is well known that regular programs of exercises improve stroke volume and increase muscle mass and tonus. The active “muscle pump” in the legs, help venous blood return, preventing venous pooling. The stronger the muscle pump, the better [23, 24]. Physical training also induces modifications in the autonomic nervous system, increasing parasympathetic activity to the heart, lowering HR and improving baroreflex sensitivity [24, 25]. All these effects are highly desirable for patients with dysautonomic syndromes.

The most important obstacle faced by dysautonomic patients to perform a program of exercise is that they are unable to tolerate exercise and, consequently, most of them become deconditioned. Deconditioning is especially common in patients with postural orthostatic tachycardia syndrome (POTS) [26, 27]. The intolerance to exercise is usually secondary to high abnormal standing HR and BP responses to exercise. The difficulty to perform exercises increases the deconditioning and a vicious cycle of exercise intolerance is developed in these patients.

However, it was demonstrated that exercise training increases the baroreflex sensitivity in patients with POTS, decreasing the upright HR [28]. Performing low intensity to moderate exercises in recumbent bike, rowing or swimming at first, avoiding the upright position helps the patient to improve physical conditioning and to tolerate the upgrades along time.

A recent report demonstrated that patients with POTS have “smaller hearts” in terms of mass and diameter, associated with reduced blood volume, when compared to control subjects [29]. A program of exercise training applied to these patients increased left ventricular mass and blood volume and reduced HR in all of them. Quality of life was also improved in all patients after the training. The practice of supervised mild exercise in the supine position at the beginning of the program is critical to provide the patient a proper exercise and warrants their adherence to long-term therapy (Fig. 1).

### Isometric counterpressure maneuvers

The physical counterpressure maneuvers have been introduced as a manner to avoid loss of consciousness and are recommended when symptoms of impending syncope occur. During the prodromal phase, BP falls prior to HR, due to vasodilatation in the skeletal muscles. Isometric maneuvers induce BP increase, delaying the loss of consciousness or avoiding it. This effect is mediated mainly by sympathetic nerve discharge and mechanical compression of the venous vascular bed [30].

The PC trial [31] was a multicenter, prospective trial, which randomized patients with recurrent vasovagal syncope and prodromal symptoms to conventional therapy alone and to conventional therapy plus physical maneuvers. The treated group had a significantly better outcome, with a relative risk reduction of 39% (95% CI 11–53%). These results motivated the inclusion of physical maneuvers as first-line treatment in patients presenting with vasovagal syncope and recognizable prodromal symptoms.

### Conclusions

In conclusion, non-pharmacological therapies, based on lifestyle modifications, postural training and physical exercises are potentially helpful for dysautonomic syndromes. The main objective of



**Figure 1.** Physical rehabilitation: the patient starts the aerobic training in seated position, on an ergometric bicycle. Tilt training is recommended from the beginning of the rehabilitation program, as well as the lower extremities resistance training. When orthostatic tolerance improves, patient continues the aerobic training on the treadmill; **A.** Lower-extremity resistance training; **B.** Seated in an ergometric bicycle; **C.** Tilt training; **D.** Treadmill.

these recommendations is to restore quality of life and self-confidence of the patients. Although there is some evidence that these interventions can affect the outcome, prospective studies enrolling

larger number of patients are necessary to prove their efficacy.

**Conflict of interest:** None declared

## References

- Freeman R, Wieling W, Axelrod FB et al. Consensus statement on the definition of orthostatic hypotension, neurally mediated syncope and the postural tachycardia syndrome. *Clin Auton Res*, 2011; 2: 69–72.
- D'Ascenzo F, Biondi-Zoccai G, Reed MJ et al. Incidence, etiology and predictors of adverse outcomes in 43,315 patients presenting to the Emergency Department with syncope: An international meta-analysis. *Int J Cardiol*, 2013; 167: 57–62.
- Baron-Esquivias G, Gómez S, Aguilera A et al. Short-term evolution of vasovagal syncope: Influence on the quality of life. *Int J Cardiol*, 2005; 102: 315–319.
- Grubb BP. Neurocardiogenic syncope and related disorders of orthostatic intolerance. *Circulation*, 2005; 111: 2997–3006.
- Alboni P, Brignole M, Degli Uberti EC. Is vasovagal syncope a disease? *Europace*, 2007; 9: 83–87.
- Moya A, Sutton R, Ammirati F et al. Guidelines for the diagnosis and management of syncope (version 2009). The Task Force for the Diagnosis and Management of Syncope of the European Society of Cardiology (ESC). Developed in collaboration with European Heart Rhythm Association (EHRA), Heart Failure Association (HFA), and Heart Rhythm Society (HRS). *Eur Heart J*, 2009; 30: 2631–2671.
- Bécher M, Binggeli C, Corti R et al. Dysfunctional baroreflex regulation of sympathetic nerve activity in patients with vasovagal syncope. *Circulation*, 2003; 107: 1620–1625.
- Morillo CA, Ellenbogen KA, Pava LF. Pathophysiologic basis for vasodepressor syncope. *Cardiol Clin*, 1997; 15: 233–249.
- Cooper VL, Hainsworth R. Effects of head-up tilting on baroreceptor control in subjects with different tolerances to orthostatic stress. *Clin Sci*, 2002; 103: 221–226.
- Stewart JM, Weldon A. Inappropriate early hypotension in adolescents: A form of chronic orthostatic intolerance with defective dependent vasoconstriction. *Pediatr Res*, 2001; 50: 97–103.
- Claydon VE, Hainsworth R. Salt supplementation improves orthostatic cerebral and peripheral vascular control in patients with syncope. *Hypertension*, 2004; 43: 809–813.
- Shannon JR, Diedrich A, Biaggioni I et al. Water drinking as a treatment for orthostatic syndromes. *Am J Med*, 2002; 112: 355–360.
- Mathias CJ, Young TM. Water drinking in the management of orthostatic intolerance due to orthostatic hypotension, vasovagal syncope and the postural tachycardia syndrome. *Eur J Neurol*, 2004; 11: 613–619.
- Miwa C, Sugiyama Y, Iwase S et al. Effects of three days of dry immersion on heart rate and blood pressure variabilities during head-up tilting in humans. *Environ Med*, 1997; 41: 135–137.
- Kamiya A, Michikami D, Fu Q et al. Pathophysiology of orthostatic hypotension after bed rest: Paradoxical sympathetic withdrawal. *Am J Physiol Heart Circ Physiol*, 2003; 285: H1158–H1167.
- Sun XQ, Yao YJ, Yang CB et al. Effect of lower body negative pressure on orthostatic tolerance and cardiac function during 21 days head-down tilt bed rest. *J Gravit Physiol*, 2003; 10: 11–17.
- Cooper VL, Hainsworth R. Head-up sleeping improves orthostatic tolerance in patients with syncope. *Clin Auton Res*, 2008; 18: 318–324.
- Ector H, Reybrouck T, Heidbüchel H et al. Tilt training: A new treatment for recurrent neurocardiogenic syncope and severe orthostatic intolerance. *Pacing Clin Electrophysiol*, 1998; 21 (Part 2): 193–196.
- Verheyden B, Ector H, Aubert AE, Reybrouck T. Tilt training increases the vasoconstrictor reserve in patients with neurally mediated syncope evoked by head-up tilt testing. *Eur Heart J*, 2008; 29: 1523–1530.
- On YK, Park J, Huh J, Kim JS. Is home orthostatic self-training effective in preventing neurally mediated syncope? *Pacing Clin Electrophysiol*, 2007; 30: 638–643.
- Duygu M, Zoghi M, Turk U et al. The role of tilt training in preventing recurrent syncope in patients with vasovagal syncope: A prospective and randomized study. *Pacing Clin Electrophysiol*, 2008; 31: 592–596.
- Foglia-Manzillo G, Giada F, Gaggioli G et al. Efficacy of tilt training in the treatment of neurally mediated syncope. A randomized study. *Europace*, 2004; 6: 199–204.
- Mtinangi B, Hainsworth R. Increased orthostatic tolerance following moderate exercise training in patients with unexplained syncope. *Heart*, 1998; 80: 596–600.
- Mtinangi BL, Hainsworth R. Effects of moderate exercise training on plasma volume, baroreceptor sensitivity and orthostatic tolerance in healthy subjects. *Exp Physiol*, 1999; 84: 121–130.
- Gardenghi G, Rondon MU, Braga AM et al. The effects of exercise training on arterial baroreflex sensitivity in neurally mediated syncope patients. *Eur Heart J*, 2007; 28: 2749–2755.
- Masaki S, Eisenach JH, Schrage WG et al. Reduced stroke volume during exercise in postural tachycardia syndrome. *J Appl Physiol*, 2007; 103: 1128–1135.
- Van Lieshout JJ. Exercise training and orthostatic intolerance: A paradox? *J Physiol*, 2003; 551: 401.
- Galbreath MM, Shibata S, Van Gundy TB et al. Effects of exercise training on arterial-cardiac baroreflex function in POTS. *Clin Auton Res*, 2011; 21: 73–80.
- Fu Q, Van Gundy TB, Galbreath MM et al. Cardiac origins of the postural orthostatic tachycardia syndrome. *J Am Coll Cardiol*, 2010; 55: 2858–2868.
- Croci F, Brignole M, Menozzi C et al. Efficacy and feasibility of isometric arm counter-pressure manoeuvres to abort impending vasovagal syncope during real life. *Europace*, 2004; 6: 287–291.
- Van Dijk N1, Quartieri F, Blanc JJ et al. Effectiveness of physical counterpressure maneuvers in preventing vasovagal syncope: The Physical Counterpressure Manoeuvres Trial (PC-Trial). *J Am Coll Cardiol*, 2006; 48: 1652–1657.