


The effectiveness of drug-coated balloons for two dissimilar calcific lesions assessed by near-infrared spectroscopy intravascular ultrasound and optical coherence tomography

Takao Konishi¹ , Kohei Saiin¹, Youji Tamaki¹, Hiroyuki Natsui¹, Tomoya Sato¹, Sakae Takenaka¹, Atsushi Tada¹, Yoshifumi Mizuguchi¹, Yuta Kobayashi¹, Hirokazu Komoriyama¹, Yoshiya Kato¹, Takuma Sato¹, Rui Kamada¹, Kiwamu Kamiya¹, Toshiyuki Nagai¹, Shinya Tanaka², Toshihisa Anzai¹

¹Department of Cardiovascular Medicine, Faculty of Medicine and Graduate School of Medicine, Hokkaido University, Sapporo, Japan

²Department of Cancer Pathology, Faculty of Medicine, Hokkaido University, Sapporo, Japan

A 63-year-old woman, who had undergone an everolimus-eluting stent implantation in the middle right coronary artery (mRCA) 3 years ago, was referred to our hospital for chest pain. Coronary angiography revealed 90% stenosis of the proximal right coronary artery (pRCA, Fig. 1A, white arrow) and mRCA (Fig. 1A, white arrowhead). Near-infrared spectroscopy-intravascular ultrasound (NIRS-IVUS) showed a fibro-fatty plaque (thick yellow arrow) with deep calcification (thin yellow arrow) in the pRCA (Fig. 1B, a). A calcified nodule (yellow arrowhead) was found in the mRCA (Fig. 1B, b). Angioplasties with paclitaxel-coated balloons 3.5/15 mm and 3.0/20 mm intra-stent were performed for pRCA and mRCA, respectively. Final angiography showed no significant RCA stenosis (Fig. 1C). However, 6 months later, sig-

nificant in-stent restenosis was observed (Fig. 1D). Optical coherence tomography (OCT) showed no significant restenosis in the pRCA (Fig. 1E, c). Meanwhile, a calcified nodule protruding intra-stent was detected in the mRCA (Fig. 1E, d, yellow arrowhead). Recent studies have shown that the stent-less strategy using drug-coated balloon (DCB) might be an effective option for calcific lesions in patients with coronary artery disease. Paclitaxel-coated balloon can inhibit the growth of smooth muscle cells, thus inhibiting neointimal proliferation. This case highlighted that DCB treatment was more effective for deep calcification with superficial fibrous plaques than for calcified nodules. NIRS-IVUS and OCT were useful for identifying different types of coronary calcifications and for predicting the effectiveness of DCB treatment.

Conflict of interest: None declared

Address for correspondence: Takao Konishi, MD, PhD, Department of Cardiovascular Medicine, Faculty of Medicine and Graduate School of Medicine, Hokkaido University, West 7, North 15, Kita-ku, Sapporo, 060-8638, Japan, tel: 011-706-6973, fax: 011-706-7874, e-mail: takaokonishi0915@gmail.com

Received: 28.03.2021 Accepted: 7.06.2021

This article is available in open access under Creative Commons Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.




Figure 1. A. The initial coronary angiogram (CAG); B. Near-infrared spectroscopy-intravascular ultrasound; C. Final CAG; D. CAG at 6 month-follow-up; E. Optical coherence tomography.