

REVIEW ARTICLE

Cardiology Journal 2020, Vol. 27, No. 6, 848–856 DOI: 10.5603/CJ.a2018.0067 Copyright © 2020 Via Medica ISSN 1897–5593 eISSN 1898–018X

# Echocardiographic predictors of atrial fibrillation recurrence after catheter ablation: A literature review

Aleksandra Liżewska-Springer, Alicja Dąbrowska-Kugacka, Ewa Lewicka, Łukasz Drelich, Tomasz Królak, Grzegorz Raczak

Department of Cardiology and Electrotherapy, Medical University of Gdansk, Poland

### Abstract

**Background:** Catheter ablation (CA) is a well-known treatment option for patients with symptomatic drug-resistant atrial fibrillation (AF). Multiple factors have been identified to determine AF recurrence after CA, however their predictive value is rather small. Identification of novel predictors of CA outcome is therefore of primary importance to reduce health costs and improve long-term results of intervention. The recurrence of AF following CA is related to severity of left ventricular (LV) dysfunction, extent of atrial dilatation and fibrosis. The aim of this paper was to present and discuss the latest studies on the utility of echocardiographic parameters in terms of CA effectiveness in patients with paroxysmal and persistent AF.

**Methods:** PubMed, Google Scholar, EBSCO databases were searched for studies reporting echocardiographic preprocedural predictors of AF recurrence after CA. LV systolic and diastolic function, as well as atrial size, strain and dyssynchrony were taken into consideration.

**Results:** Twenty one full-text articles were analyzed, including three meta-analyses. Several echocardiographic parameters have been reported to determine a risk of AF recurrence after CA. There are conventional methods that measure left atrial size and volume, LV ejection fraction, parameters assessing LV diastolic dysfunction, and methods using more innovative technologies based on speckle tracking echocardiography to determine left atrial synchrony and strain. Each of these parameters has its own predictive value.

**Conclusions:** Regarding CA effectiveness, every patient has to be evaluated individually to estimate the risk of AF recurrence, optimally using a combination of several echocardiographic parameters. (Cardiol J 2020; 27, 6: 848–856)

Key words: atrial fibrillation, catheter ablation, pulmonary vein isolation, echocardiography, predictors, recurrence

## Introduction

Atrial fibrillation (AF) is the most common cardiac arrhythmia with a projected prevalence of 14–17 million by the year 2030 in the European Union [1]. AF remains one of the major causes of stroke, heart failure, sudden death, and cardiovascular morbidity. With large increases in the burden of AF expected in the coming decades, better diagnosis and stratification of treatment selection is of paramount importance.

In recent years catheter ablation (CA) became a common treatment for patients with symptomatic, drug-resistant AF. The success rate of CA, defined as no AF relapse, of up to 70% for patients with paroxysmal AF and around 50% in those with persistent AF [1]. Multiple factors have been identified for AF recurrence after CA, such

Address for correspondence: Dr. Alicja Dąbrowska-Kugacka, Department of Cardiology and Electrotherapy MedicalUniversity of Gdansk, ul. Dębinki 7, 80–952 Gdańsk, Poland, e-mail: alicja.dabrowska-kugacka@gumed.edu.plReceived: 26.03.2018Accepted: 10.06.2018

as age, AF duration, ventricular and atrial function and comorbidities, however their predictive value is rather small [1, 2]. Therefore, new predictors of procedural outcome are needed for better identification of the most suitable candidates for CA.

Optimal patient selection is crucial to avoid unnecessary risk associated with CA, which can be accompanied by serious complications such as cardiac tamponade, stroke, pulmonary vein stenosis and atrio-esophageal fistula. Finally more accurate prediction can influence a decision for continuation of long-term oral anticoagulant and antiarrhythmic drug therapy.

Much of the information on ventricular and atrial function can be derived from cardiac magnetic resonance and tomography, but for practical reasons echocardiography is mostly used in clinical settings. Active deformation of the heart muscle during the cardiac cycle can be assessed with strain imaging from two-dimentional speckle tracking echocardiography (STE). This technique enables the recognition of subtle cardiac dysfunction and markers myocardial damage. The ultimate goal of these markers is to define different types of AF that are characterized by specific pathophysiology which may warrant early aggressive intervention and will respond favourably to CA. Recurrent AF after CA seems to be higher in patients with signs of atrial cardiomyopathy, also ventricular function plays a major role in the efficacy of this procedure.

The aim of this paper was to present and discuss the latest studies on echocardiographic parameters in terms of CA effectiveness in the treatment of patients with paroxysmal and persistent AF.

PubMed, Google Scholar, EBSCO databases were searched using the key words "echocardiographic predictors of atrial fibrillation after catheter ablation" or "echocardiography atrial fibrillation," "echocardiography pulmonary vein isolation" or "echocardiography catheter ablation." The search returned 104 abstracts, published from 1997 to 2017, all in English. After screening the abstracts, 42 were included for full-text analysis, according to their relevance to the subject. The criteria to include studies were as follows: (A) patients with paroxysmal or persistent AF referred for CA, (B) endpoint analysis taking into account the first recurrence of AF (defined as any documented episode of AF lasting > 30 s). Finally, 21 full-text articles were included in relation to CA in patients with AF. The following echocardiographic criteria were taken into consideration: left (LA) and right atrial (RA) size (diameter, area and volume), left ventricular (LV) ejection fraction (LVEF), parameters assessing LV diastolic dysfunction, atrial strain and dyssynchrony (Table 1).

## Methods

## Atrial size

It is established that LA size contributes to structural remodeling and therefore to atrial fibrosis [3]. Dilatation of the LA is an independent predictor of new onset AF [4]. It is also a wellknown predictor of low success for CA in terms of AF recurrence [5–7].

Zhuang et al. [8] performed a meta-analysis of 22 studies which included over 3700 patients, which showed that an increased antero-posterior diameter of the LA was associated with a higher risk of AF recurrence after CA. The antero-posterior diameter of LA before CA, obtained from the parasternal long-axis view, was 35-50 mm. In patients with arrhythmia recurrences it was 1.87 mm larger than in those with successful pulmonary vein isolation [8]. According to European expert consensus from 2012 the LA diameter over 50-55 mm indicates limited success of CA [6]. In a study performed by Liao et al. [9] on 589 patients with paroxysmal AF, LA diameter over 43 mm and LV systolic diameter over 31 mm were the best cutoff values for predicting AF progression after CA. Their predictive value was highest when both of the above-mentioned diameters were exceeded [9]. LA dilation can be asymmetric and the anteroposterior dimension underestimates and does not truly reflect LA size. Studies focusing on LA area, however, did not confirm its predictive utility. Tomas at al. [10] found that LA area > 24 cm<sup>2</sup> did not predict AF recurrence at 12 month follow up. Njoku et al. [11] in a meta-analysis of 21 observational studies encompassing 3850 patients reported that patients with AF recurrence had larger mean left atrial volume (LAV) and LAV index (LAVi) compared to patients with no arrhythmia relapse. Moreover, they found that increased LAV/LAVi was independently associated with frequent AF recurrences after CA. There was a 3% increase in the odds of AF relapse per unit increase in LAV/ /LAVi. Shin et al. [12] found that LAVi of 34 mL/ /m<sup>2</sup> showed a sensitivity of 70% and a specificity of 91% to predict AF recurrence.

Atrial fibrillation is a biatrial disease. RA, with its enlargement and remodeling also involved in AF relapse [13]. Moon et al. [13] reported that increased RA volume index (RAVi) might affect early AF recurrence (within 3 months) after CA and RAVi over 78 mL/m<sup>2</sup> predicted the early recurrence with

Ľ.
atio
ilde
er
het
cat
st-
d
JCe
rrer
scul
e re
tio
illa
fibr
<u>.</u>
atr
i of
ors
fact
ic 1
aph
ogr
rdio
оса
cho
ē
ĭί<
sdic
pre
the
'nt
oqe
es ;
udi
Sti
е 1.
lde
Ĕ

				al.			t al.	t.al.				al.	1
Reference	Zhuang et e 2011	Njoku et al 2017		Nan Wen et 2017		Cha et al. 2011	Black-Maier e 2017	Marrouche et 2018		Hirai et al. 2014	Masuda et a 2017	Okamatsu et 2015	
Prognostic ECHO parameter	LAD was 1.87 mm higher in patients with arrhythmia recurrences	Higher mean LAVi (using ECHO, CT, or MRI) Higher mean LAV (using ECHO, CT, or MRI)	Higher mean LAV/LAVi (using ECHO, CT, or MRI)	RAD > 35.5 mm (in PAF with LAD > 35 mm)		LVEF ≤ 40%	LVEF was not a risk factor	LVEF ≤ 25%		Average E/e' > 13 (during sinus rhythm)	Average E/e' > 14	E/e' ≥ 15	
AF recurrences after CA (n)	SN	NN NN NN NN	NS	284 (60.6%)		SN	NS	60 (37%)		76 (38%)	38 (18%)	9 (41%)	
Median follow-up [months]	6–30 ± 13	17 (mean) 17 (mean)	18	24.3 ± 18		13	1	37.8		12	12 ± 6	21 ± 12	
LVEF [%]	N	× 50 × 50	> 50	> 50		≤ 40 (111 patients), > 50 (157 patients with isolated LVDD), 100 pa- tients with normal LV function	< 50 (97 patients), ≥ 50 (133 patients)	≤ 35		> 50	> 50	> 50	
Persistent AF (n)	1476	544 537*	960	274		198	150	245		25	102	17	
Paroxysmal AF (n)	2274	611 913*	1932	196	FUNCTION	170	80	118	SFUNCTION	173	113	D	
Study type	Meta- -analysis	Meta- -analysis Meta- analysis	Meta- analysis	Retrospective	<b>AR SYSTOLIC DYS</b>	Retrospective	Retrospective	Prospective	<b>AR DIASTOLIC DY:</b>	Retrospective	Prospective	Retrospective	
Total patients (n)	ARIAL SIZE 3750 (22 studies)	1425 (9 studies) 1559 (11 studies)	2886 (13 studies)	470	LEFT VENTRICUL	368	230	363	LEFT VENTRICUL	198	215	22	

tt-catheter ablation.
recurrence pos
atrial fibrillation
hic factors of a
chocardiograpl
ie predictive ec
udies about th
e 1 (cont.). St
Table

al patients	Study type	Paroxysmal AF (n)	Persistent AF (n)	LVEF (%)	Median follow-up [months]	AF recurrences after CA (n)	Prognostic ECHO parameter	Reference
	Retrospective	152	80	> 50	36	59 (25%)	LVDD defined as: septal e″ < 8 lateral e″ < 10 LAV ≥ 34 mL/m² (risk factor of VLR)	Onishi et al. 2017
	Prospective	70	54	> 50	12	42 (34%, early- during first week after PVI), 26 (27%, late)	E/A ratio (in early recurrence)	Kosiuk et al. 2014
T ATRIAL STR	IAIN							
tudies)	Meta- analysis	529	157	> 50	11	232 (94.9%)	LA strain < 22.8% (using QRS or P analysis, using different software packages)	Ma et al. 2016
	Retrospective	68	32	> 50	12	26 (26%)	Basal LA lateral strain < 25.3%	Yasuda et al. 2015
	Retrospective Prospective	0	102	> 50	15	57 (56%)	LA strain < 10% (during an episode of persistent AF)	Parwani et al. 2017
	Prospective	42	0	> 50	14 ± 7	12 (29%)	LA global peak area strain (3D-GASs) < 28.9% (measured in systole, determined by 3D speckle tracking)	Mochizuki et al. 2017
HER PARAMET	rers							
	Retrospective	25	0	> 42	20 ± 9	7 l (28%)	A mechanical dyssynchrony (speckle tracking strain analysis)	Loghin et al. 2013
	Retrospective	132	0	> 50	23 ± 7	38 (28.8%)	FMR ≥ 1 (regurgitation jet area to LA area during sinus rhythm)	Qiao et al. 2016
dy not stated; 3D- er; AF — atrial fibi 1 by pulsed wave I ation; LA — left ati in fraction; MRI — — very late recurre	GASs —left atrial glob. rillation; CA — cathetel Doppler; e' — early dis rium; LAD — left atrial - magnetic resonance ii ences (> 12 months)	al peak area strain in r ablation; CT — com astolic mitral annular diameter; LAV — lef maging; n — numbe	systole determined t nputed tomography; velocity assesed by ft atrial volume; LAVi rr of patients; PAF —	yy three-dimensior E — early mitral in tissue Doppler; e' — left atrial volum paroxysmal AF; P∖	ial speckle tracking flow velocity asses — early diastolic i e index; LV — left VI — pulmonary ve	i echocardiography; A seed by pulsed wave D mitral annular velocity ventricular; LVDD — I in isolation; RAD — ri	·— late mitral inflow velocity assess oppler; E/A — mitral early to late dis assessed by tissue Doppler; FMR — eft ventricular diastolic dysfunction; ght atrial diameter assessed in the 4.	ed by pulsed wave astolic peak ratio as- - functional mitral re- LVEF — left ventricular -chamber apical view;

74% sensitivity and 68% specificity [13]. Wen et al. [14] found that RA size predicted successful CA in patients with paroxysmal AF and LA horizontal diameter (determined as the measurement from the middle of mitral isthmus to the LA roof in the 4-chamber apical view) enlargement over 35 mm. RA was measured from the middle of tricuspid isthmus to the RA roof in the 4-chamber apical view, and when this diameter was below 35.5 mm it predicted AF recurrence-free survival at over 2-year follow up. Although a large cohort (over 400 patients) was examined, the results might be applicable only to an Asian population.

## Left ventricular systolic function

In previous years several trials comparing amiodarone with CA in AF patients with heart failure (HF) and reduced ejection fraction (HFrEF) showed ablation to be superior at maintaining sinus rhythm [15, 16]. However the efficacy of CA in HFrEF patients with AF is still a matter of debate. Development of AF in HFrEF patients occurs due to eccentric remodeling of the LA. In this group an increase in LA diameter and volume was observed. The underlying electrical substrate driving to AF is likely different than in patients with HF and preserved ejection fraction, where LA stiffness is dominant [17].

In a study performed by Cha et al. [18] three groups of patients with AF undergoing CA were analyzed: 111 with LV systolic dysfunction (LVEF  $\leq 40\%$ ), 157 with isolated LV diastolic dysfunction (LVEF > 50%) and 100 individuals with normal LV function. The authors reported that AF elimination rate was significantly lower in patients with systolic dysfunction (62%, p = 0.002) and non-significantly lower in those with diastolic dysfunction (75%, p = 0.15) when compared with the group with normal LV function.

Black-Maier et al. [16] compared two groups of about 100 patients with LVEF of less and over 50% and showed no significant differences in the rates of atrial arrhythmia recurrence between HF patients with preserved and reduced ejection fraction (33.9% vs. 32.6%; p = 0.8) at 1-year follow up. The contradictory result may be due to differences in the study population enrolled. Patients in the Black-Maier et al. [16] study were older and were more likely to have hypertension and diabetes mellitus.

The efficacy of CA in patients with HFrEF and AF was uncertain, until the lately published breakthrough CASTLE-AF trial [19]. It was a multicenter, randomized, controlled trial to assess whether

CA lowers morbidity and mortality as compared with medical therapy (rate or rhythm control) in patients with coexisting AF and medically managed HF. The study included almost 400 patients with symptomatic paroxysmal or persistent AF and HF in the New York Heart Association class II-IV, LVEF of 35% or less and an implanted defibrillator. Mortality and hospitalization for worsening HF was significantly reduced in patients treated with CA. After 5 year follow-up sinus rhythm was maintained in 63% of patients in the ablation group vs. 22% in the medically treated group (p < 0.001). The study also revealed that patients with LVEF of less than 25% were less likely to benefit from ablation than those with LVEF between 25% and 35%. It would be interesting to determine the success rate of CA in patients with advanced HF but preserved ejection fraction.

## Left ventricular diastolic dysfunction

Left ventricular diastolic dysfunction may indicate an increase in LV filling pressure, which can influence LA remodeling over the long term [20]. Wall stress due to increased atrial pressure plays an important role in the development of atrial electrical and structural remodeling. Impaired diastolic function has also been reported to be associated with AF recurrences [21]. According to guidelines of the American Society of Echocardiography, LV diastolic dysfunction can be evaluated and graded by mitral inflow assessed by pulsed wave Doppler, tissue Doppler of LV walls, and LA size [22].

In 2014 Hirai et al. [23] reported that elevated LA pressure, as determined by increased average E/e' index (E — early transmitral flow velocity obtained by pulsed wave Doppler; e'— early diastolic mitral velocity measured by tissue Doppler; averaged annular septal and lateral values), was the only echocardiographic parameter that predicted AF recurrence after CA. The E/e' value over 13 indicated increased risk of AF relapse during 12 month follow up.

Three years later Masuda et al. [24] reported that patients with E/e' > 14 before CA more frequently developed recurrent atrial tachyarrhythmias after single and multiple procedures. It was the first study demonstrating that patients with high E/e' ratio, along with age, female sex, low body mass index, and persistent AF more frequently presented low-voltage areas within the LA predisposing to AF, when endocardial voltage mapping was performed during the CA procedure. The relationship between higher E/e' ratio and the presence of low voltage areas in the LA may indicate that higher E/e' is associated with advanced atrial arrhythmogenic substrate outside the pulmonary veins. This might explain AF recurrences even after a properly performed (without reconnections) pulmonary vein isolation.

Finally Okamatsu et al. [25] who examined 24 patients with hypertrophic cardiomyopathy demonstrated that E/e' ratio was the only predictor of AF recurrence following pulmonary veins isolation (PVI). Patients with  $E/e' \ge 15$  had a significantly higher risk of AF recurrence than those with E/e' < 15. Thus, patients with AF and mild or moderate LV diastolic dysfunction (E/e' < 15) are better candidates for PVI than those with a restrictive inflow pattern.

In a study performed by Onishi et al. [26] LV diastolic dysfunction at baseline was the only independent risk factor of late AF recurrence, defined as first AF relapse after more than 12 months subsequent to CA. The authors defined LV diastolic dysfunction very strictly and patients had to fulfill all three of the following criteria: early diastolic septal annular velocity e' < 8 cm/s, lateral annular velocity e' < 10 cm/s and LAVi  $\ge$  34 mL/m<sup>2</sup>. After a single CA procedure, reconnections of pulmonary veins could affect recurrence of AF. To minimize the influence of pulmonary vein reconnections, risk factors of late recurrences after multiple CA procedures, not only after a single session were examined. LV diastolic dysfunction appeared to be the only risk factor of late AF relapses.

Kosiuk et al. [27] showed that E/A mitral inflow pattern (early to late mitral inflow velocity ratio, assessed by pulsed-wave Doppler at the level of the mitral valve), which is associated with LV diastolic dysfunction, was the best pre-procedural predictor of short-term AF recurrence during the first week after PVI. An E/A ratio of 1.35 was the cut-off value with the highest sensitivity and specificity for predicting early AF recurrence. In contrast to Onishi et al. [26], the authors reported that none of the parameters evaluating LV diastolic function predicted long-term PVI result (defined as any documented AF episode within a 3-12 month period after ablation). This discrepancy may be due to different definitions of LV diastolic dysfunction applied in both studies. Moreover, Kosiuk et al. [27] analyzed the results of CA after only a single procedure, and it can also explain the contradictory results because, as mentioned above, late AF recurrences are mainly due to pulmonary vein reconnections [28, 29]. Finally, a limited number of patients with severe LV diastolic dysfunction might preclude evaluation of other parameters, like E/e'

ratio as predictors of both, short- and long-term AF recurrences after CA.

### Left atrial strain

Schneider et al. [30] evaluated results of tissue Doppler imaging (TDI)-based LA strain analysis in patients with AF for the prediction of successful CA. The authors showed that patients with higher atrial strain and strain rate after CA appear to have a greater likelihood of maintaining sinus rhythm. Peak strain and strain rate were measured at each mid-LA segment one day prior to, within 24 hours, and 3 months after CA. A value of 20% for atrial septal systolic strain obtained before the procedure predicted sinus rhythm maintenance after CA, but with rather low sensitivity (57%) and specificity (56%). This method, however, is prone to angulation error and suffers from variable reproducibility. Moreover, it does not allow distinguishing active myocardial contraction from its passive motion.

Two-dimensional STE (2D-STE) is angle independent and thus more useful for LA strain analysis [31]. There are several studies indicating that LA strain has higher predictive value than LA size obtained from conventional echocardiography [31, 32]. The LA strain reflects LA reservoir, conduit and booster pump function. Furthermore, it correlates with the extent of LA fibrosis, especially in patients with persistent AF [7, 33].

A meta-analysis of 8 studies, which included 686 patients with paroxysmal AF showed that global LA strain is useful to identify individuals at high risk of AF recurrence after CA [34]. This analysis included both patients with sinus rhythm or AF at baseline, before PVI. In the group with AF, the beginning of QRS was set as the zero strain point. In patients with sinus rhythm the trigger for strain analysis was put either at the onset of QRS complex or P-wave. LA peak positive strain of less than 22.8% predicted AF recurrence with 78% sensitivity and 75% specificity. These results were independent of the applied method of LA strain analysis: from the beginning of QRS or P-wave, as well as software package used.

Left atrial strain measured on LA lateral wall by 2D-STE might be the most useful parameter for predicting successful AF ablation as it represents pure LA contractile function [35]. Yasuda et al. [35] indicated a significant prognostic value of basal LA lateral total strain, both in patients with sinus rhythm and AF during examination. The authors set the zero strain point at the beginning of QRS complex in the group with AF and at P wave in the group with sinus rhythm. The total strain was calculated as follows: positive peak strain – negative peak strain. They reported that a value below 25.3% in basal LA lateral strain showed 81% sensitivity and 72% specificity for predicting AF recurrence after CA [35].

Most of the studies included in the mentioned meta-analysis [34] assessed patients with mainly paroxysmal or different types of AF. Parwani et al. [36] reported that low LA peak positive strain (< 10%) during an episode of persistent AF was strongly linked to recurrence of AF after one or even after two CA procedures.

Finally, some results suggest that threedimensional STE (3D-STE) is potentially more accurate than 2D-STE for assessment of LA dysfunction in patients with AF [37, 38]. Mochizuki et al. [39] assessed patients with paroxysmal AF and found that global LA strain determined by 3D-STE less than 28.9% was a predictor of AF recurrence after the first-time CA Moreover, 3D-STE global LA strain was a better predictor of AF relapse after CA than LA strain obtained by 2D-STE. Threedimensional strain analysis reflects LA function in many directions: not only longitudinal, but also circumferential and area strains. This can explain its superiority over 2D-STE.

## Left atrial mechanical dyssynchrony

The LA mechanical dyssynchrony, which clearly indicates the presence of atrial structural and electrical remodeling, can predict recurrence of AF after CA in patients with paroxysmal AF [40, 41]. It can be determined by different echocardiographic methods.

Den Uijl et al. [41] reported that total atrial conduction time (PA-TDI), reflecting atrial electrical remodeling, was an independent predictor of AF recurrence after PVI. Total atrial conduction time was obtained by measuring the time delay between the onset of the P-wave on the surface electrocardiogram and the peak A'-wave of spectral tissue Doppler tracing on LA lateral wall. The prolonged value of 146  $\pm$  20 ms was associated with AF recurrence after CA (for comparison PA-TDI in healthy subjects was  $78 \pm 7$  ms) [42]. Similarly, Fukushima et al. [43] found that PA-TDI was an independent predictor of AF recurrence in patients with paroxysmal AF. They reported 2.5--fold higher rate of AF relapses in patients with PA-TDI duration > 151.3 ms. Evranos et al. [44] demonstrated a relationship between PA-TDI on LA lateral wall and recurrence of AF in patients treated with cryobaloon ablation. PA-TDI over 125 ms predicted AF recurrence with 80% sensitivity and 90% specificity. Unlike den Uijl et al. [41] they defined PA-TDI as a time interval from the onset of P wave on the surface electrocardiogram to the beginning of the A' wave.

Loghin at al. [40], used an algorithm based on 2D-STE — the vector velocity imaging (VVI) and looked at the timing of peak longitudinal strain obtained on opposing LA walls during atrial contractile phase. They found that maximum opposing walls delay of over 51 ms predicted AF recurrence after CA with 89% sensitivity and 72% specificity. Unfortunately, the study was retrospective and based on a small cohort of 25 patients.

Sarvari et al. [45] measured the dispersion of LA contraction duration, defined as the time difference from the peak of P wave on the surface electrocardiogram during sinus rhythm to maximum LA shortening assessed by 2D-STE strain (peak negative longitudinal strain). The standard deviation of contraction durations measured in 18 LA segments was defined as LA mechanical dispersion. The authors reported that patients with AF relapse and normal (LAV  $25 \pm 10 \text{ mL/m}^2$ ) presented with significantly greater LA mechanical dispersion compared with patients after successful CA (38  $\pm$  14 ms vs. 30  $\pm$  12 ms; p < 0.001). Therefore LA mechanical dispersion can be a useful tool to predict AF recurrence after CA in patients with structurally normal heart.

## **Functional mitral regurgitation**

Qiao et al. [46] reported that functional mitral regurgitation, defined as regurgitation jet area to LA area ratio  $\geq 0.1$  in subjects without any primary valvular disease, independently predicted long-term outcomes post ablation in patients with paroxysmal AF. Functional mitral regurgitation was strongly correlated with the presence and extent of low voltage zones within the LA, assessed invasively prior to ablation.

## Conclusions

Several echocardiographic parameters have been reported to determine the risk of AF recurrence after CA. These parameters reflect morphology, function and myocardial remodeling in patients with AF. There are conventional methods that measure LA size and volume, LVEF, parameters assessing LV diastolic dysfunction, and methods using more innovative technologies based on STE to determine LA synchrony and strain. Each of these parameters has their own predictive value. Unfortunately, there is no single parameter that actually enables the prediction of AF relapse after CA. To summarize, the predictors of AF recurrence after CA which were confirmed by several groups were LA diameter > 50–55 mm or LAVi > 34 mL/m<sup>2</sup>, E/e' > 13–15, LA strain assessed by STE < 20–25% and total atrial conduction time measured by TDI > 150 ms. The presence of LV systolic dysfunction also lowered CA success rate with a bottom LVEF cut-off value of < 25%. It needs underlining that risk of AF recurrence after CA should be estimated individually, optimally on the basis of several echocardiographic parameters.

#### Conflict of interest: None declared

#### References

- Kirchhof P, Benussi S, Kotecha D, et al. ESC Scientific Document Group . 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. Eur Heart J. 2016; 37(38): 2893–2962, doi: 10.1093/eurheartj/ehw210, indexed in Pubmed: 27567408.
- Vizzardi E, Curnis A, Latini MG, et al. Risk factors for atrial fibrillation recurrence: a literature review. J Cardiovasc Med (Hagerstown). 2014; 15(3): 235–253, doi: 10.2459/ JCM.0b013e328358554b, indexed in Pubmed: 23114271.
- Burstein B, Nattel S. Atrial fibrosis: mechanisms and clinical relevance in atrial fibrillation. J Am Coll Cardiol. 2008; 51(8): 802–809, doi: 10.1016/j.jacc.2007.09.064, indexed in Pubmed: 18294563.
- Vaziri SM, Larson MG, Benjamin EJ, et al. Echocardiographic predictors of nonrheumatic atrial fibrillation. The Framingham Heart Study. Circulation. 1994; 89(2): 724–730, indexed in Pubmed: 8313561.
- Berruezo A, Tamborero D, Mont L, et al. Pre-procedural predictors of atrial fibrillation recurrence after circumferential pulmonary vein ablation. Eur Heart J. 2007; 28(7): 836–841, doi: 10.1093/eurheartj/ehm027, indexed in Pubmed: 17395676.
- Calkins H, Kuck K, Cappato R, et al. 2012 HRS/EHRA/ECAS Expert Consensus Statement on Catheter and Surgical Ablation of Atrial Fibrillation: Recommendations for Patient Selection, Procedural Techniques, Patient Management and Follow-up, Definitions, Endpoints, and Research Trial Design. Heart Rhythm. 2012; 9(4): 632–696.e21, doi: 10.1016/j.hrthm.2011.12.016.
- Cameli M, Lisi M, Righini FM, et al. Usefulness of atrial deformation analysis to predict left atrial fibrosis and endocardial thickness in patients undergoing mitral valve operations for severe mitral regurgitation secondary to mitral valve prolapse. Am J Cardiol. 2013; 111(4): 595–601, doi: 10.1016/j.amjcard.2012.10.049, indexed in Pubmed: 23211360.
- Zhuang J, Wang Y, Tang K, et al. Association between left atrial size and atrial fibrillation recurrence after single circumferential pulmonary vein isolation: a systematic review and meta-analysis of observational studies. Europace. 2011; 14(5): 638–645, doi: 10.1093/europace/eur364.
- 9. Liao YC, Liao JN, Lo LW, et al. Left Atrial Size and Left Ventricular End-Systolic Dimension Predict the Progression of Paroxysmal Atrial Fibrillation After Catheter Ablation. J Cardiovasc Elec-

trophysiol. 2017; 28(1): 23–30, doi: 10.1111/jce.13115, indexed in Pubmed: 27779351.

- Tomas L, Orosco A, Vergara JM, et al. Predictors of recurrence and outcomes in catheter ablation of paroxysmal atrial fibrillation. Rev Argent Cardiol. 2017; 85(3): 240–246.
- 11. Njoku A, Kannabhiran M, Arora R, et al. Left atrial volume predicts atrial fibrillation recurrence after radiofrequency ablation: a meta-analysis. Europace. 2018; 20(1): 33–42, doi: 10.1093/europace/eux013, indexed in Pubmed: 28444307.
- Shin SH, Park MY, Oh WJ, et al. Left atrial volume is a predictor of atrial fibrillation recurrence after catheter ablation. J Am Soc Echocardiogr. 2008; 21(6): 697–702, doi: 10.1016/j. echo.2007.10.022, indexed in Pubmed: 18187293.
- Moon J, Hong YJ, Shim J, et al. Right atrial anatomical remodeling affects early outcomes of nonvalvular atrial fibrillation after radiofrequency ablation. Circulation . 2012; 76(4): 860–867, doi: 10.1253/circj.cj-11-1232.
- Wen SN, Liu N, Bai R, et al. Right atrial diameter and outcome of catheter ablation of atrial fibrillation. J Interv Card Electrophysiol. 2017; 49(2): 157–164, doi: 10.1007/s10840-017-0258-2.
- 15. Di Biase L, Mohanty P, Mohanty S, et al. Ablation versus amiodarone for treatment of persistent atrial fibrillation in patients with congestive heart failure and an implanted device: results from the AATAC multicenter randomized trial. Circulation. 2016; 133(17): 1637–1644, doi: 10.1161/CIRCULATIONA-HA.115.019406, indexed in Pubmed: 27029350.
- Black-Maier E, Ren X, Steinberg BA, et al. Catheter ablation of atrial fibrillation in patients with heart failure and preserved ejection fraction. Heart Rhythm. 2018; 15(5): 651–657, doi: 10.1016/j. hrthm.2017.12.001, indexed in Pubmed: 29222043.
- Melenovsky V, Hwang SJ, Redfield MM, et al. Left atrial remodeling and function in advanced heart failure with preserved or reduced ejection fraction. Circ Heart Fail. 2015; 8(2): 295–303, doi: 10.1161/CIRCHEARTFAILURE.114.001667, indexed in Pubmed: 25593126.
- Cha YM, Wokhlu A, Asirvatham SJ, et al. Success of ablation for atrial fibrillation in isolated left ventricular diastolic dysfunction: a comparison to systolic dysfunction and normal ventricular function. Circ Arrhythm Electrophysiol. 2011; 4(5): 724– -732, doi: 10.1161/CIRCEP.110.960690, indexed in Pubmed: 21747059.
- Marrouche NF, Brachmann J, Andresen D, et al. Catheter ablation for atrial fibrillation with heart failure. N Engl J Med. 2018; 378(5): 417–427, doi: 10.1056/NEJMoa1707855, indexed in Pubmed: 29385358.
- Hu YF, Hsu TL, Yu WC, et al. The impact of diastolic dysfunction on the atrial substrate properties and outcome of catheter ablation in patients with paroxysmal atrial fibrillation. Circulation. 2010; 74(10): 2074–2078, indexed in Pubmed: 20668352.
- Park J, Joung B, Uhm JS, et al. High left atrial pressures are associated with advanced electroanatomical remodeling of left atrium and independent predictors for clinical recurrence of atrial fibrillation after catheter ablation. Heart Rhythm. 2014; 11(6): 953–960, doi: 10.1016/j.hrthm.2014.03.009, indexed in Pubmed: 24607916.
- Nagueh SF, Smiseth OA, Appleton CP, et al. Recommendations for the Evaluation of Left Ventricular Diastolic Function by Echocardiography: An Update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. J Am Soc Echocardiogr. 2016; 29(4): 277–314, doi: 10.1016/j. echo.2016.01.011, indexed in Pubmed: 27037982.

- Hirai T, Cotseones G, Makki N, et al. Usefulness of left ventricular diastolic function to predict recurrence of atrial fibrillation in patients with preserved left ventricular systolic function. Am J Cardiol. 2014; 114(1): 65–69, doi: 10.1016/j.amjcard.2014.03.061, indexed in Pubmed: 24819904.
- Masuda M, Fujita M, Iida O, et al. An E/e' ratio on echocardiography predicts the existence of left atrial low-voltage areas and poor outcomes after catheter ablation for atrial fibrillation. Europace. 2018; 20(5): e60–e68, doi: 10.1093/europace/eux119, indexed in Pubmed: 28651348.
- Okamatsu H, Ohara T, Kanzaki H, et al. Impact of left ventricular diastolic dysfunction on outcome of catheter ablation for atrial fibrillation in patients with hypertrophic cardiomyopathy. Circ J. 2015; 79(2): 419–424, doi: 10.1253/circj.CJ-14-0823, indexed in Pubmed: 25452101.
- Onishi N, Kaitani K, Amano M, et al. Relationship between left ventricular diastolic dysfunction and very late recurrences after multiple procedures for atrial fibrillation ablation. Heart Vessels. 2018; 33(1): 41–48, doi: 10.1007/s00380-017-1027-y, indexed in Pubmed: 28766046.
- Kosiuk J, Breithardt OA, Bode K, et al. The predictive value of echocardiographic parameters associated with left ventricular diastolic dysfunction on short- and long-term outcomes of catheter ablation of atrial fibrillation. Europace. 2014; 16(8): 1168–1174, doi: 10.1093/europace/eut415, indexed in Pubmed: 24569573.
- Ouyang F, Antz M, Ernst S, et al. Recovered pulmonary vein conduction as a dominant factor for recurrent atrial tachyarrhythmias after complete circular isolation of the pulmonary veins: lessons from double Lasso technique. Circulation. 2005; 111(2): 127–135, doi: 10.1161/01.CIR.0000151289.73085.36, indexed in Pubmed: 15623542.
- Sauer WH, McKernan ML, Lin D, et al. Clinical predictors and outcomes associated with acute return of pulmonary vein conduction during pulmonary vein isolation for treatment of atrial fibrillation. Heart Rhythm. 2006; 3(9): 1024–1028, doi: 10.1016/j. hrthm.2006.05.007, indexed in Pubmed: 16945795.
- Schneider C, Malisius R, Krause K, et al. Strain rate imaging for functional quantification of the left atrium: atrial deformation predicts the maintenance of sinus rhythm after catheter ablation of atrial fibrillation. Eur Heart J. 2008; 29(11): 1397–1409, doi: 10.1093/eurheartj/ehn168, indexed in Pubmed: 18436560.
- Hammerstingl C, Schwekendiek M, Momcilovic D, et al. Left atrial deformation imaging with ultrasound based two-dimensional speckle-tracking predicts the rate of recurrence of paroxysmal and persistent atrial fibrillation after successful ablation procedures. J Cardiovasc Electrophysiol. 2012; 23(3): 247–255, doi: 10.1111/j.1540-8167.2011.02177.x, indexed in Pubmed: 21955059.
- Mirza M, Caracciolo G, Khan U, et al. Left atrial reservoir function predicts atrial fibrillation recurrence after catheter ablation: a two-dimensional speckle strain study. J Interv Card Electrophysiol. 2011; 31(3): 197–206, doi: 10.1007/s10840-011-9560-6, indexed in Pubmed: 21424845.
- 33. Kuppahally SS, Akoum N, Burgon NS, et al. Left atrial strain and strain rate in patients with paroxysmal and persistent atrial fibrillation: relationship to left atrial structural remodeling detected by delayed-enhancement MRI. Circ Cardiovasc Imaging. 2010; 3(3): 231–239, doi: 10.1161/CIRCIMAGING.109.865683, indexed in Pubmed: 20133512.
- Ma XX, Boldt LH, Zhang YL, et al. Clinical relevance of left atrial strain to predict recurrence of atrial fibrillation after catheter ablation: a meta-analysis. Echocardiography. 2016; 33(5): 724–733, doi: 10.1111/echo.13184, indexed in Pubmed: 26857344.

- 35. Yasuda R, Murata M, Roberts R, et al. Left atrial strain is a powerful predictor of atrial fibrillation recurrence after catheter ablation: study of a heterogeneous population with sinus rhythm or atrial fibrillation. Eur Heart J Cardiovasc Imaging. 2015; 16(9): 1008–1014, doi: 10.1093/ehjci/jev028, indexed in Pubmed: 25750193.
- Parwani AS, Morris DA, Blaschke F, et al. Left atrial strain predicts recurrence of atrial arrhythmias after catheter ablation of persistent atrial fibrillation. Open Heart. 2017; 4(1): e000572, doi: 10.1136/openhrt-2016-000572, indexed in Pubmed: 28674624.
- Mochizuki A, Yuda S, Oi Y, et al. Assessment of left atrial deformation and synchrony by three-dimensional speckle-tracking echocardiography: comparative studies in healthy subjects and patients with atrial fibrillation. J Am Soc Echocardiogr. 2013; 26(2): 165–174, doi: 10.1016/j.echo.2012.10.003, indexed in Pubmed: 23140846.
- 38. Kobayashi Y, Okura H, Kobayashi Y, et al. Assessment of atrial synchrony in paroxysmal atrial fibrillation and impact of pulmonary vein isolation for atrial dyssynchrony and global strain by three-dimensional strain echocardiography. J Am Soc Echocardiogr. 2014; 27(11): 1193–1199, doi: 10.1016/j.echo.2014.08.004, indexed in Pubmed: 25240493.
- Mochizuki A, Yuda S, Fujito T, et al. Left atrial strain assessed by three-dimensional speckle tracking echocardiography predicts atrial fibrillation recurrence after catheter ablation in patients with paroxysmal atrial fibrillation. J Echocardiogr. 2017; 15(2): 79–87, doi: 10.1007/s12574-017-0329-5, indexed in Pubmed: 28155065.
- Loghin C, Karimzadehnajar K, Ekeruo IA, et al. Outcome of pulmonary vein isolation ablation for paroxysmal atrial fibrillation: predictive role of left atrial mechanical dyssynchrony by speckle tracking echocardiography. J Interv Card Electrophysiol. 2014; 39(1): 7–15, doi: 10.1007/s10840-013-9841-3, indexed in Pubmed: 24310816.
- den Uijl DW, Gawrysiak M, Tops LF, et al. Prognostic value of total atrial conduction time estimated with tissue Doppler imaging to predict the recurrence of atrial fibrillation after radiofrequency catheter ablation. Europace. 2011; 13(11): 1533–1540, doi: 10.1093/europace/eur186, indexed in Pubmed: 21712264.
- Erdem FH, Erdem A, Özlü F, et al. Electrophysiological validation of total atrial conduction time measurement by tissue doppler echocardiography according to age and sex in healthy adults. J Arrhythm. 2016; 32(2): 127–132, doi: 10.1016/j.joa.2015.11.006, indexed in Pubmed: 27092194.
- 43. Fukushima K, Fukushima N, Ejima K, et al. Left atrial appendage flow velocity and time from P-wave onset to tissue Doppler-derived A' predict atrial fibrillation recurrence after radiofrequency catheter ablation. Echocardiography. 2015; 32(7): 1101–1108, doi: 10.1111/echo.12823, indexed in Pubmed: 25362992.
- Evranos B, Aytemir K, Oto A, et al. Predictors of atrial fibrillation recurrence after atrial fibrillation ablation with cryoballoon. Cardiol J. 2013; 20(3): 294–303, doi: 10.5603/cj.2013.0075.
- Sarvari SI, Haugaa KH, Stokke TM, et al. Strain echocardiographic assessment of left atrial function predicts recurrence of atrial fibrillation. Eur Heart J Cardiovasc Imaging. 2016; 17(6): 660–667, doi: 10.1093/ehjci/jev185, indexed in Pubmed: 26219297.
- Qiao Yu, Wu L, Hou B, et al. Functional mitral regurgitation: predictor for atrial substrate remodeling and poor ablation outcome in paroxysmal atrial fibrillation. Medicine (Baltimore). 2016; 95(30): e4333, doi: 10.1097/MD.00000000004333, indexed in Pubmed: 27472715.