



# Plurihormonalność guzów przysadki w świetle badań immunohistochemicznych

Wielohormonalność guzów przysadki w świetle badań immunohistochemicznych

Marek Pawlikowski<sup>1</sup>, Jolanta Kunert-Radek<sup>1</sup>, Maciej Radek<sup>2</sup>

<sup>1</sup>Department of Neuroendocrinology, Chair of Endocrinology, Medical University, Łódź, Poland

<sup>2</sup>Chair of Neurosurgery and Surgery of Peripheral Nerves, Medical University, Łódź, Poland

## Abstract

**Introduction:** Plurihormonalność guzów przysadki może być zdefiniowana jako zdolność guza przysadki do wyrażania więcej niż jednego hormonu przysadkowego. Zastosowanie immunohistochemii do diagnostyki usuniętych chirurgicznie guzów przysadki pozwoliło na stwierdzenie, że w istocie znaczną część guzów przysadki jest wielohormonalna. Jednak dane o częstości występowania i znaczeniu klinicznym wielohormonalności są skąpe i częściowo sprzeczne.

**Material and methods:** Stosowano 55 guzów przysadki, usuniętych chirurgicznie, które badano immunohistochemicznie z użyciem przeciwciał przeciw hormonom przysadki lub ich podjednostkom. Dodatkowo, 40 guzów przysadki badano z użyciem przeciwciała Ki-67, aby ocenić potencjał proliferacyjny.

**Results:** Według zaleceń Światowej Organizacji Zdrowia (WHO), nie uważamy guzów przysadki wyrażających zarówno FSH i LH (gonadotropinomas) jak i somatotropinomas wyrażających GH i prolaktynę za wielohormonalne. Mimo tej ograniczonej definicji, plurihormonalność została stwierdzona w guzach przysadki zarówno w grupie guzów hormonalnie czynnych, jak i w grupie guzów klinicznie nieczynnych. Wykazano, że ponad jedna trzecia (36.1%) badanych guzów przysadki wyraża więcej niż jeden hormon. Plurihormonalność, szczególnie ta, która jest związana z współwystępowaniem ACTH, jest częściej obserwowana w guzach nawrotowych. Guzy wielohormonalne wykazują także wyższy indeks Ki-67 w porównaniu z guzami monohormonalnymi.

**Conclusions:** Plurihormonalność jest częstym zjawiskiem zarówno w grupie guzów hormonalnie czynnych, jak i w grupie guzów klinicznie nieczynnych. Ponadto wielohormonalność wydaje się wiązać z wyższym ryzykiem nawrotu guza. (Pol J Endocrinol 2010; 61 (1): 63-66)

**Key words:** guzy przysadki, immunohistochemia, plurihormonalność, nawrót guza

## Streszczenie

**Wstęp:** Wielohormonalność guzów przysadki jest definiowana jako ich zdolność do ekspresji więcej niż jednego hormonu przysadkowego. Zastosowanie immunohistochemii do diagnostyki usuniętych chirurgicznie guzów przysadki pozwoliło na stwierdzenie, że w istocie znaczną część guzów przysadki jest wielohormonalna. Jednak dane o częstości występowania i znaczeniu klinicznym wielohormonalności są skąpe i częściowo sprzeczne.

**Materiał i metody:** Stosowano 55 guzów przysadki, usuniętych chirurgicznie, które badano immunohistochemicznie z użyciem przeciwciał przeciw hormonom przysadkowym i/lub ich podjednostkom. Dodatkowo 40 guzów przysadki badano z użyciem przeciwciała Ki-67, w celu ustalenia ich potencjału proliferacyjnego.

**Wyniki:** Zgodnie z zaleceniami Światowej Organizacji Zdrowia (WHO, *World Health Organization*) nie zaliczamy do guzów przysadki wielohormonalnych guzów z łączną ekspresją FSH i LH (gonadotropinoma) ani somatotropinoma wykazujących łączną ekspresję GH i prolaktyny. Nawet przy tym ograniczeniu, wielohormonalność jest częstym zjawiskiem zarówno w przypadku guzów hormonalnie czynnych, jak i w grupie guzów klinicznie nieczynnych. Wykazano, że w ponad jednej trzeciej (36.1%) badanych guzów przysadki wykrywa się obecność więcej niż jednego hormonu. Wielohormonalność, zwłaszcza związana z współwystępowaniem ACTH, jest częściej obserwowana w guzach nawrotowych. Guzy wielohormonalne wykazują także wyższy indeks Ki-67 w porównaniu z guzami monohormonalnymi.

**Wnioski:** Wielohormonalność jest częstym zjawiskiem zarówno w grupie guzów klinicznie czynnych, jak i w grupie guzów klinicznie nieczynnych hormonalnie. Ponadto wielohormonalność wydaje się wiązać z wyższym ryzykiem nawrotu guza.

(Endokrynol Pol 2010; 61 (1): 63-66)

**Słowa kluczowe:** guzy przysadki, immunohistochemia, wielohormonalność, nawrót guza



## Introduction

Plurihormonality of pituitary adenomas can be defined as the ability of an adenoma to express more than one pituitary hormone. According to the recommendations of the WHO, the gonadotropinomas expressing both gonadotropins, *i.e.* FSH and LH, are not considered plurihormonal. The same concerns adenomas in patients with acromegaly, secreting prolactin in addition to GH [1]. Plurihormonal adenomas leading to the elevated concentrations of more than one pituitary hormone in blood are considered relatively rare. The application of immunohistochemistry to diagnose surgically removed pituitary tumours revealed that a great number of pituitary adenomas, including the so-called “clinically non-functioning” pituitary adenomas (CNFPA), are in fact plurihormonal [2–8]. However, the data on the incidence and the clinical relevance of pituitary adenoma plurihormonality remain scarce.

The aim of this study was to evaluate the incidence of plurihormonality in a non-selected group of immunohistochemically examined pituitary adenomas and to answer the question whether the plurihormonality of the tumours is relevant for their prognosis.

## Material and methods

The archival material of 155 surgically removed pituitary adenomas was studied. Paraffin sections of each tumour were immunostained using the primary antibodies against the following pituitary hormones or their subunits: prolactin (PRL, polyclonal, Dako, Denmark), growth hormone (GH, polyclonal, Dako or Immunon, USA), LH (Dako, monoclonal), FSH (Dako, monoclonal), TSH (monoclonal, Immunotech, France), ACTH (polyclonal, Sigma), and alpha-subunit (alpha-SU, monoclonal, Immunotech, France). Additionally, 40 adenomas were immunostained with anti-Ki-67 antigen (MIB-1) antibody (Dako-Cytomation, Denmark). Visualization of the immune reactions was done by means of streptavidin-biotin-peroxidase technique with use of 3,3'-diaminobenzidine as chromogen. The presence of more than 1% of hormone immunopositive cells was considered important. Statistical analysis of the numeric data was done by means of McNemara's and  $\chi^2$  tests (recurrence rates) or ANOVA followed by LSD test (Ki-67 indices); statistical significance was at the level  $p < 0.05$ .

## Results

The results are presented in Tables I–III. As can be seen, over one third of pituitary adenomas examined by immunohistochemistry are plurihormonal (Table I). If, following the WHO recommendations, we do not consi-

Table I. Incidence of plurihormonal pituitary adenomas

Tabela I. Zachorowalność na gruczolaki wielohormonalne przysadki

Total	56/155 (36.1%)
Acromegaly	
including GH/PRL	18/29 (62.1%)
excluding GH/PRL	7/29 (24.1%)
Cushing's disease	4/14 (28.6%)
CNFPA	38/85 (44.7%)

Table II. Recurrence rate of pituitary adenomas

Tabela II. Wskaźniki występowania nawrotów gruczolaków przysadki

Total	32/155 (20.6%)
Plurihormonal monohormonal	10/56 (17.8%)
Including gonadotropinomas	22/99 (22.2%)
Excluding gonadotropinomas	6/64 (9.4%)
Acromegaly	2/29 (6.9%)
GH/PRL	1/18 (5.5%)
Plurihormonal (GH/alfaSU)	1/7 (14.3%)
Cushing's disease	3/14 (21.4%)
Plurihormonal	3/4 (75%)

Table III. Clinically non-functioning pituitary adenomas (CNFPA)

Tabela III. Klinicznie nieczynne gruczolaki przysadki (CNFPA)

Total	85/155 (55.2%)	Recurrent 29/85 (34.1%)
"Pure" gonadotropinomas	35/85 (41.1%)	Recurrent 16/35 (45.7%)
Monohormonal (excluding gonadotropinomas)	7/85 (8.2%)	Recurrent 4/7 (57.1%)
Plurihormonal	38/85 (44.1%)	Recurrent 9/38 (23.7%)
Inactive	=5/85 (5.9%)	Recurrent 0/5 (0%)

der GH/PRL adenomas in acromegaly or FSH/LH adenomas as plurihormonal, then plurihormonality is most frequent in clinically non-functioning adenomas (CNFPA) (see Table I). Approximately 1/5 of the investigated adenomas were recurrent. The recurrence rate is highest in case of gonadotropinomas (45.7%). However, if we exclude this type of adenoma from the group of monohormonal tumours, the recurrence rate of plu-

rihormonal adenomas is twice as high (17.8% *v.* 9.4%), although the difference is not significant statistically. A statistically significant difference in the recurrence rate is found in corticotropinomas manifesting themselves as Cushing's disease. All the recurrent tumours observed in patients with Cushing's disease were plurihormonal (Table II). In the case of CNFPA, the highest recurrence rate is observed in monohormonal adenomas. However, the number of such tumours is relatively low, and, interestingly, almost all recurrent adenomas in this group (3/4) were "silent" (*i.e.* not manifested as Cushing's disease) corticotropinomas. The next position among the recurrent CNFPAs belongs to "pure" gonadotropinomas. In contrast, no recurrent tumours were found in the group of hormonally immunonegative tumours (Table III). We also analyzed the relationship between tumour recurrence and expression of ACTH in plurihormonal CNFPA. Some tendency towards a higher recurrence rate was observed in the group of plurihormonal CNFPAs co-expressing ACTH, as compared to ACTH-negative tumours (6/17, 35.3% *v.* 2/14, 14.2%, respectively). Moreover, the mean Ki-67 index (% of Ki-67-immunopositive cell nuclei), calculated for plurihormonal tumours, was significantly higher than the monohormonal adenoma group ( $6.0 \pm 1.9\%$  vs.  $1.7 \pm 0.5\%$ , mean  $\pm$  SEM,  $p < 0.05$ ).

## Discussion

Our data show that plurihormonality is a frequent phenomenon in both hormonally active and clinically non-functioning pituitary adenomas. Their frequency in our material (36.1%) is very similar to that (31%) reported earlier by Ho et al. [5]. This finding supports the opinion of quoted authors that the immunostaining of all pituitary hormones is mandatory for correct classification of pituitary adenomas. Earlier studies with the application of double immunostaining at the light microscopy and the electron microscopy levels showed that the multiple hormones are expressed within the same tumoural cell, and even within the same secretory granule [9]. Although double immunostaining was not applied in the present study, the sum of the percentages of cells immunopositive for different hormones often exceeded 100. This means that the cells must present double or even triple hormone immunopositivity. These findings indicate that plurihormonal adenomas are constituted from plurihormonal cells rather than from multiple cellular lines expressing different pituitary hormones. The question arises whether the expression of multiple hormones in tumoural cells is relevant for their natural history, and specifically whether it possesses a prognostic value. The recurrence of pituitary

adenoma after surgery may depend on early or late diagnosis and the skilfulness of the neurosurgeon, but also depends on tumour aggressiveness. The latter is linked with tumour invasiveness, *i.e.* the ability of the tumour cells to invade the adjacent structures consequently enabling total excision of the tumour, and with its high proliferation potency. The latter is usually measured by calculation of the percentage of cell nuclei positively immunostained with antibody to Ki-67 antigen (Ki-67 index). Generally, Ki-67 index is positively correlated with tumour invasiveness and recurrence, although some exceptions have been observed, and the low Ki-67 index can be noticed in aggressive tumours, and, in contrast, the high proliferation index may occur in indolent adenomas [10]. It was important to see whether plurihormonality increases the risk of the tumour recurrence. Although the high recurrence rate in our material concerns gonadotropinomas, which according to WHO recommendations are considered "monohormonal" adenomas, and monohormonal ACTH-immunopositive adenomas without Cushing's disease manifestation ("silent" corticotropinomas), in the remaining types of adenomas the plurihormonality seems to be linked with the increased risk of tumour recurrence. This suggestion is supported by our finding that the mean value of Ki-67 index in plurihormonal pituitary adenomas is significantly higher than in monohormonal tumours. In our earlier paper we showed that plurihormonal adenomas present higher values of another proliferation index, namely proliferating cells nuclear antigen, PCNA [11]. Our observation of the higher recurrence rate in cases of plurihormonal versus monohormonal corticotropinomas with Cushing's disease is compatible with the findings of Desai et al. [12]. The quoted authors have found that corticotropinomas co-secreting the alpha-subunit are more aggressive and have a higher recurrence rate than corticotropinomas without the expression of this peptide. The increased risk of tumour recurrence seems also to be linked with ACTH co-expression in plurihormonal CNFPA, which corroborates with the data of Bradley et al. [13]. We suggest that plurihormonality of pituitary adenomas, especially that linked with ACTH co-expression, seems to predict a higher risk of tumour recurrence. However, this presumption needs further studies based on higher number of examined tumours.

## Acknowledgements

This paper is financially supported by the Medical University of Lodz. The technical assistance of Mrs Maria Jaranowska, Mrs Małgorzata Jędrzejewska, and Mrs Anna Opładowska is greatly appreciated.

## References

1. Solcia E, Kloppel G, Sobin LH. Histological typing of endocrine tumors World Health Organization, International Histological Classification of Tumors, Springer 2004, 16–22.
2. Ikeda H, Yoshimoto, Yandell DW. Hormone production and c-myc protein labeling in plurihormonal pituitary adenomas. *Endocr Pathol* 1994; 5: 162–168.
3. Matsuno A, Teramoto A, Takekoshi S et al. HGH, PRL and ACTH gene expression in clinically nonfunctioning adenomas detected with nonisotopic in situ hybridization method. *Endocr Pathol* 1995; 6: 13–20.
4. Scheithauer BW, Kovacs K, Horvath E. The adenohypophysis In: Lechago J, Gould VE (eds). *Bloodworth Endocrine Pathology*. Williams and Wilkins, Baltimore 1997: 85–152.
5. Ho DM, Hsu CY, Ting LT et al. Plurihormonal pituitary adenomas: immunostaining of all hormones is mandatory for correct classification *Histopathology* 2001; 39: 310–319.
6. Kunert-Radek J, Radek A, Gruszka A et al. Immunohistochemical investigation of clinically nonfunctioning pituitary tumors as a prognostic factor of tumor recurrence Proceedings, 12th Int Congress of Endocrinology, Lisbon 31.VIII-4.IX.2004, Medimond.
7. Salehi F, Cohen S, Syro LV et al. Plurihormonality in pituitary adenomas associated with acromegaly *Endocr Pathol* 2006; 17: 291–296.
8. Coculescu M, Badiu C, Galoiu S et al. Evolution under complex therapy of acromegaly due to a pituitary plurihormonal adenoma with colocalization of GH and FSH *Acta Endocrinol* 2006; 2: 337–348.
9. Zielińska M, Zielińska A, Kunert-Radek J et al. Immunohistochemical electron microscope study on coexistence of hormones in pituitary adenomas *Endokrynologia Polska* 2008; 59 (Suppl. A): P2–46, 62–63.
10. Turner HE, Wass JA Are markers of proliferation valuable in the histological assessment of pituitary tumours? *Pituitary* 1999; 1: 147–151.
11. Pawlikowski M, Gruszka A, Kurnatowska I et al. Proliferating cell nuclear antigen (PCNA) expression in pituitary adenomas: relationship to the endocrine phenotype of adenoma *Folia Histochem Cytobiol* 2006; 44: 37–41.
12. Desai B, Burrin M, Nott CA, Geddes JF, Lamb EJ, Wood DF et al. Glycoprotein hormone alpha-subunit production and plurihormonality in human corticotroph tumours — an in vitro and immunohistochemical study *Eur J Endocrinol* 1995; 133: 25–32.
13. Bradley KJ, Wass JA, Turner HE. Non-functioning pituitary adenomas with positive immunoreactivity for ACTH behave more aggressively than ACTH immunonegative tumours but not recur more frequently *Clin Endocrinol* 2003; 58: 59–64.