

Supplementary File

Table S1. Median values from Q1 and Q3 and minimum and maximum values for age differences between study groups.

A) For major osteoporotic fractures

| | Age (N = 93) | |
|---|---------------------|-----------------|
| | Median (Q1–Q3) | Minimum–Maximum |
| Very high risk (FRAX for MOF) (N = 38) | 70.00 (64.00–74.00) | 58.00–85.00 |
| High risk (FRAX for MOF) (N = 55) | 71.00 (67.00–77.00) | 58.00–89.00 |

B) For hip fracture

| | Age (N = 92) | |
|--|---------------------|-----------------|
| | Median (Q1–Q3) | Minimum–Maximum |
| Very high risk (FRAX for HF) (N = 76) | 72.00 (66.00–75.50) | 58.00–89.00 |
| High risk (FRAX for HF) (N = 16) | 69.50 (67.50–75.50) | 61.00–81.00 |

C) For extended assessment criteria

| | Age (N=115) | |
|---------------------------------|---------------------|-----------------|
| | Median (Q1–Q3) | Minimum–Maximum |
| Very high risk (N = 115) | 70.00 (66.00–75.00) | 53.00–89.00 |
| High risk (N = 40) | 67.00 (63.50–69.50) | 55.00–82.00 |