

# Diosmetin ameliorates osteoarthritic inflammation *in vivo* and ECM macromolecules degradation in interleukin-1 $\beta$ -stimulated murine chondrocytes through the Nrf2/NF- $\kappa$ B pathway

Liang Qian<sup>1</sup>, Chuang Li<sup>1</sup>, Hong Liu<sup>1</sup>, Hui Zhou<sup>1</sup>, Tao Tan<sup>2</sup>

<sup>1</sup>Department of Orthopedics, The Eighth Hospital of Wuhan, Wuhan, China

<sup>2</sup>Department of Trauma Surgery, The Eighth Hospital of Wuhan, Wuhan, China

## Abstract

**Introduction.** Osteoarthritis (OA) is a prevailing degenerative disease in elderly population and can lead to severe joint dysfunction. Studies have revealed various pharmacological activities of diosmetin, including the anti-OA efficacy. The present study further investigated its effect on interleukin (IL)-1 $\beta$ -induced OA in chondrocytes.

**Material and methods.** Primary chondrocytes were isolated from young mice, stimulated with IL-1 $\beta$  (10 ng/mL), and pretreated with diosmetin (10 and 20  $\mu$ M) to conduct the *in vitro* assays. CCK-8 assay assessed the cytotoxicity of diosmetin whereas the levels of inflammatory factors (PGE2, nitrite, TNF- $\alpha$ , and IL-6) in homogenized cells were evaluated by ELISA. The levels of inflammatory cytokines, content of extracellular matrix (ECM), and signaling-related proteins (Nrf2, HO-1, and NF- $\kappa$ B p65) were assessed by western blotting. Expression of collagen II, p65, and Nrf2 in the chondrocytes was confirmed by immunofluorescence staining. The chondrocytes treated with IL-1 $\beta$  and diosmetin were transfected with Nrf2 knockdown plasmid (si-Nrf2) to investigate the role of Nrf2. *In vivo* OA mouse model was induced by surgically destabilizing the medial meniscus (DMM). Safranin O staining was conducted to assess the OA severity in the knee-joint tissue.

**Results.** Diosmetin suppressed the expression of iNOS, COX-2, PGE2, nitrite, TNF- $\alpha$ , IL-6, MMP-13, and ADAMTS-5 induced by IL-1 $\beta$  in chondrocytes. The expression of p-p65, p-I $\kappa$ B $\alpha$ , and nuclear p65 was decreased whereas that of Nrf2 and HO-1 increased by diosmetin treatment in IL-1 $\beta$ -treated chondrocytes. Nrf2 knockdown by siRNA reversed the inhibitory effect of diosmetin on IL-1 $\beta$ -induced degradation of ECM proteins and inflammatory factors in cultured chondrocytes. In the DMM-induced model of OA, diosmetin alleviated cartilage degeneration and decreased the Osteoarthritis Research Society International score.

**Conclusions.** Diosmetin ameliorates expression of inflammation biomarkers and ECM macromolecules degradation in cultured murine chondrocytes *via* inactivation of NF- $\kappa$ B signaling by activating Nrf2/HO-1 signaling pathway. (*Folia Histochem Cytobiol* 2024; 62, 2:99–109)

**Keywords:** mouse; osteoarthritis; chondrocytes; diosmetin; ECM proteins; Nrf2/HO-1; NF- $\kappa$ B

## INTRODUCTION

Osteoarthritis (OA) is a degenerative joint disorder that leads to unbearable pain, seriously affecting the

life quality of more than 250 million people worldwide [1]. Degeneration and inflammation of articular cartilage are the two main features of it [2]. Current treatments for OA mainly focus on the pain management and symptom alleviation [3]. However, those pain-relief drugs, such as corticosteroids, opioids, and nonsteroidal anti-inflammatory drugs (NSAIDs) have certain side effects, and surgical treatment also carries significant risks, which remain an obstacle to preventing OA [4]. Thus, novel therapeutic targets with fewer side effects are needed.

### Correspondence address:

Tao Tan  
The Eighth Hospital of Wuhan,  
No.1 Huasheng Road, Tazihu Street, Jiang'an District,  
Wuhan 430000, Hubei Province, China  
e-mail: tantaodoctor@hotmail.com

Increasing studies have revealed inflammatory mechanisms and cartilage extracellular matrix (ECM) damage as the crucial risk factors for the OA progression [5]. It is well known that the key pro-inflammatory cytokine interleukin-1 $\beta$  (IL-1 $\beta$ ) is highly expressed in the cartilage tissue of OA patients [6]. It can trigger the secretion of other inflammatory mediators synthesized by inducible nitric oxide synthase 2 (iNOS) and cyclooxygenase-2 (COX-2) and proteolytic enzymes including matrix metalloproteinases (MMPs) in OA chondrocytes, resulting in the disorders of the homeostasis between ECM synthesis and degradation [7]. Therefore, inhibition of IL-1 $\beta$ -induced inflammation and ECM degradation has become a potential approach to alleviate the progression of OA.

Diosmetin (3',5,7-trihydroxy-4'-methoxy flavone) is a type of flavonoid, which is abundant in lemon peel and citrus fruits [8]. Hesperidin, an another flavanone derived from *Citrus aurantium L.*, has been reported to repress the OA development [9]. Multiple pharmacological activities of diosmetin have been found, such as anti-cancer [10], anti-oxidant [11], anti-microbial [12], anti-apoptotic [13], anti-estrogenic [14], anti-hyperglycemic [15], and anti-diabetic [16]. Particularly, it has also been reported to be effective in inhibiting inflammatory reaction in various diseases, including myocardial ischemia-reperfusion injury [17], colitis [18], and rheumatoid arthritis [19]. In recent years, the beneficial effects of diosmetin on bone-related diseases have been revealed. For instance, diosmetin regulates bone formation and has an anti-osteoporosis function [20, 21]. Shao *et al.* revealed its inhibitory effect on osteolysis induced by lipopolysaccharide [22]. It can also protect against osteopenia induced by chronic kidney disease [23]. In a recent study, diosmetin has been shown to prevent the loss of subchondral bone in OA mouse model [24]. However, its specific role in IL-1 $\beta$ -treated chondrocytes has not been detected yet.

In this study, we investigated the effect of diosmetin on IL-1 $\beta$ -induced degradation of ECM proteins and inflammatory response in cultured murine chondrocytes. We also revealed the possible signaling pathways that participate in the regulatory role of diosmetin in OA. The results of this study may provide novel research option for improving OA treatments.

## MATERIALS AND METHODS

### Animal model

C57BL/6 mice (male; 5–8 weeks old) were provided by the Animal Center of Chinese Academy of Sciences (Shanghai, China) and randomly divided into three groups (n = 12 in each group): Sham; DMM; and DMM + diosmetin. All the mice were anesthetized

using 1% sodium pentobarbital (60 mg/kg). To establish an OA model, the right knee joint capsule of the mouse was exposed, and then medial meniscotibial ligaments was sectioned to induce the destabilization of the medial meniscus (DMM) as described by Glasson *et al.* [25]. Mice in the Sham group received arthrotomy without transecting the meniscotibial ligament. Finally, the right knee joint capsule and skin were sutured using surgical absorbable suture. Two weeks after the DMM surgery, diosmetin (10 mg/kg; purity: 99.80%; 520-34-3; MedChemExpress, China) dissolved in sterile saline was subcutaneously injected into the mice in the DMM + diosmetin group every other day for 7 consecutive days. The treatment dose was used according to the previous study [22]. Meanwhile, the mice in the Sham and DMM groups only received 1 ml of sterile 0.9% NaCl. At the end of the drug treatment, all the mice were euthanized using an overdose injection of pentobarbital sodium (100 mg/kg), and the knee joints of all mice were resected for the subsequent experiments. All animal experiments were performed in accordance with the National Institutes of Health Guidelines for the Care and Use of Laboratory Animals and were approved by the Ethical Review System for Laboratory Animal Welfare of the Wuhan Myhalic Biotechnology Co., Ltd (approval number: HLK-20207112; approval date: 2023.7.26).

### Safranin O staining

The collected knee joints were fixed in paraformaldehyde (4%; Sigma-Aldrich, MO, USA) for 24 h and then decalcified in EDTA (10%; Sigma-Aldrich). Followed by paraffin embedding, the specimens were sectioned into slices (5  $\mu$ m-thick) and stained with safranin O/fast green (Solarbio, Beijing, China) [26]. Then, the Osteoarthritis Research Society International (OARSI) scoring system was used to grade the severity of OA (between 0 and 15) in each group according to Glasson *et al.* recommendations [27].

### Cell culture and treatment

Primary chondrocytes were isolated from C57BL/6 mice (2 weeks old; Animal Center of Chinese Academy of Sciences). Briefly, the articular cartilage was separated from the knee joint of the mice after euthanasia (cervical dislocation under inhalation anesthesia) and cut into 1–3 mm pieces, which were then treated with trypsin (0.25%; Gibco, CA, USA) for 10 min and collagenase type II (0.25%; Gibco) for 1.5 h at 37°C and centrifuged for 5 min at 1000 g to obtain the cell pellet. Then, the collected chondrocytes were cultured in Dulbecco's modified Eagle's medium (Gibco) containing penicillin/streptomycin (1%; Gibco) and

fetal bovine serum (10%; Gibco) at 37°C. Cells were passaged when the confluency reached 80–90%, and the second-generation chondrocytes were selected. Chondrocytes were stimulated with IL-1 $\beta$  (10 ng/mL; PeproTech, Drive Cranbury, NJ, USA) for 24 h and then the cells were treated with diosmetin (10 and 20  $\mu$ M) for 48 h. In rescue assay, cells were transfected with si-Nrf2 (sc-37030; Santa Cruz, USA) and its control using Lipofectamine 2000 (Thermo Fisher, Waltham, MA, USA). Transfection efficiency was verified using western blotting analysis.

### CCK-8 assay

Cell counting kit-8 (Beyotime, Haimen, Jiangsu, China) was used to assess the cytotoxicity of diosmetin to mouse chondrocytes. Chondrocytes were seeded on 96-well plates and treated with diosmetin (5, 10, 20, or 30  $\mu$ M) for 24 h and 48 h. Then, the CCK-8 solution (10  $\mu$ L) was added. After 4 h of incubation, the absorbance at 450 nm was determined using a microplate reader (Thermo Fisher).

### Western blotting

Total protein extraction from the treated mouse chondrocytes was conducted using RIPA lysis buffer (Beyotime), and the BCA protein assay kit (Beyotime) was used to evaluate the protein concentration. Proteins (50  $\mu$ g) were separated by SDS-PAGE (10% gels) and electro-transferred onto PVDF membrane (Millipore, USA), which was subsequently blocked with 5% skim milk. Primary antibodies against iNOS (ab178945; 1:1000; Abcam, UK), COX-2 (ab179800; 1:1000), collagen II (ab34712; 1:1000), aggrecan (FNab00213; 1:500; FineTest, China), MMP-13 (ab39012; 1:3000), ADAMTS-5 (ab41037; 1:250), p-p65 (ab76302; 1:1000), p65 (ab32536; 1:1000), p-I $\kappa$ B $\alpha$  (ab133462; 1:10000), I $\kappa$ B $\alpha$  (ab32518; 1:1000), Nrf2 (abs130481; 1:500; Absin Bioscience, China), HO-1 (ab52947; 1:2000), GAPDH (ab181602; 1:10000), and lamin B (ab151735; 1:2000) were incubated at 4°C with the membranes overnight. Then, goat anti-rabbit secondary antibody (ab205718; 1:2000) was added. Finally, electrochemiluminescence plus reagent (Invitrogen, China) was prepared for visualizing the protein bands, and the intensity of protein bands was quantified with Image Lab 3.0 software (Bio-Rad, USA).

### ELISA

Mouse chondrocytes were seeded on 6-well plates and treated with IL-1 $\beta$  (10 ng/mL) for 24 h and diosmetin (10 and 20  $\mu$ M) for 48 h. Then the levels of PGE<sub>2</sub>, Nitrite, TNF- $\alpha$ , and IL-6 in the cell culture supernatants were measured by ELISA kits (Invitrogen) according to the manufacturer's protocol.

### Immunofluorescence

The chondrocytes were fixed with 4% paraformaldehyde for 10 min and permeabilized with Triton X-100 (0.1%; Sigma-Aldrich) for 15 min. After rinsing with phosphate-buffered saline (PBS), cells were incubated with primary antibodies against Collagen II (ab34712), p65 (ab32536), and Nrf2 (abs130481; 1:100; Absin Bioscience) at 4°C overnight. Then, the secondary antibodies were added (ab150077; 1:200) for 1 h at room temperature. Then, the nuclei of cells were stained with 4',6-diamidino-2'-phenylindole (Sigma-Aldrich) for 30 min. At the end, an Olympus IX73 fluorescence microscope (Olympus, Tokyo, Japan) was used for analyzing the results.

### Statistical analysis


SPSS18.0 software (IBM, USA) was used to conduct the statistical analysis, and the data are presented as mean  $\pm$  SD. Significant differences ( $P < 0.05$ ) were analyzed using Student's *t*-test or one-way analysis of variance followed by the Tukey *post hoc* tests. Nonparametric data (OARSI scores) were evaluated using the Kruskal-Wallis H test. All the experiments were repeated at least three times.

## RESULTS

### Diosmetin inhibits IL-1 $\beta$ -induced markers of inflammation in chondrocytes

To confirm the optimal concentrations of diosmetin, CCK-8 assay was conducted to evaluate the cell viability of chondrocytes under the treatment with diosmetin (5, 10, 20, and 30  $\mu$ M). Chondrocytes viability had no significant alteration when the cells were treated with 5, 10, and 20  $\mu$ M of diosmetin for 24 and 48 h; however, 30  $\mu$ M of diosmetin slightly reduced the viability of cells after 48 h (Fig. 1A). Therefore, diosmetin at 10  $\mu$ M and 20  $\mu$ M concentrations was used for the subsequent experiments. After the stimulation of chondrocytes by IL-1 $\beta$ , the protein levels of enzymes producing inflammatory mediators (iNOS and COX-2) were increased in the chondrocytes. Subsequent treatment of cells with diosmetin remarkably reduced the protein levels of iNOS and COX-2 (Fig. 1B, C).

To further check the effect of diosmetin on inflammatory biomarkers induced by IL-1 $\beta$  in chondrocytes, ELISA was performed to further assess the changes in PGE<sub>2</sub>, nitrite, TNF- $\alpha$ , and IL-6 levels in the supernatants. Consistently, the treatment with diosmetin suppressed the release of PGE<sub>2</sub>, nitrite, TNF- $\alpha$ , and IL-6, indicating its anti-inflammatory role in the *in vitro* OA model (Fig. 1D–G).



**Figure 1.** Diosmetin inhibits IL-1 $\beta$ -induced inflammation in cultured murine chondrocytes. **A.** CCK-8 assessed the viability of chondrocytes under the treatment of diosmetin (5, 10, 20, and 30  $\mu$ M). **B.** **C.** Western blotting of iNOS and COX-2 expression in chondrocytes treated with IL-1 $\beta$  and diosmetin. **D.** **G.** The levels of inflammatory markers in the supernatants of chondrocytes treated with IL-1 $\beta$  and diosmetin were measured by ELISA as described in Methods. \*\* $P < 0.01$ , \*\*\*  $P < 0.001$

### Diosmetin suppresses IL-1 $\beta$ -induced ECM molecules' degradation in chondrocytes

In this part of our study, we further explored the effect of diosmetin on ECM molecules degradation. We found (Fig. 2) that the levels of collagen II and aggrecan proteins were decreased while the levels of MMP-13 and ADAMTS-5 proteins were increased in the chondrocytes stimulated by IL-1 $\beta$ , suggesting severe degradation of ECM molecular components in chondrocytes. By contrast, diosmetin promoted collagen II and aggrecan expression and restrained MMP-13 and ADAMTS-5 expression in IL-1 $\beta$ -treated chondrocytes (Fig. 2A–C). Immunofluorescence also showed decreased collagen II expression after IL-1 $\beta$  stimulation, while collagen II expression was restored in the cells incubated with IL-1 $\beta$  and diosmetin (Fig. 2D), indicating that diosmetin had inhibitory effect on ECM molecules' degradation in IL-1 $\beta$ -treated chondrocytes.


### Diosmetin inactivates NF- $\kappa$ B signaling pathway in IL-1 $\beta$ -treated chondrocytes

To investigate the potential mechanisms of diosmetin action, expression of signaling-related

proteins was evaluated in cultured chondrocytes. Relative to the control group, IL-1 $\beta$  upregulated p-p65 and p-I $\kappa$ B $\alpha$  expression and downregulated I $\kappa$ B $\alpha$  expression, and diosmetin reversed these alterations (Fig. 3A, B). Additionally, the protein level of nuclear p65 in the chondrocytes was increased by IL-1 $\beta$  and was reduced by diosmetin (Fig. 3C, D). Consistent with that, immunofluorescence showed that diosmetin decreased the fluorescence of nuclear p65 induced by IL-1 $\beta$  in the chondrocytes (Fig. 3E), suggesting that diosmetin inhibited NF- $\kappa$ B signaling by blocking NF- $\kappa$ B p65 nuclear translocation.

### Diosmetin activates Nrf2/HO-1 signaling pathway in IL-1 $\beta$ -treated chondrocytes

Western blotting revealed that the levels of nuclear Nrf2 and HO-1 proteins were not affected by IL-1 $\beta$  in chondrocytes. However, diosmetin treatment markedly enhanced the protein levels of nuclear Nrf2 and HO-1 in IL-1 $\beta$ -treated chondrocytes (Fig. 4A, B). Consistently, enhanced fluorescence intensity of nuclear Nrf2 was observed in the IL-1 $\beta$  + diosmetin group relative to the IL-1 $\beta$  group (Fig. 4C), suggesting




**Figure 2.** Diosmetin suppresses IL-1 $\beta$ -induced ECM degradation in chondrocytes. **A–C.** Western blotting of ECM-related proteins in chondrocytes treated with IL-1 $\beta$  and diosmetin. **D.** Immunofluorescence of collagen II expression in chondrocytes treated with IL-1 $\beta$  and diosmetin. \*\* $P < 0.01$ , \*\*\* $P < 0.001$ .

that diosmetin activated Nrf2/HO-1 pathway in IL-1 $\beta$ -treated chondrocytes.


### Diosmetin inhibits IL-1 $\beta$ -induced changes in chondrocytes *via* activation of Nrf2/HO-1 signaling pathway

In this part of our study, chondrocytes were treated with IL-1 $\beta$  (10 ng/mL) and diosmetin (20  $\mu$ M) and then transfected with Nrf2 knockdown plasmid (si-Nrf2) or its control (si-NC) to investigate how Nrf2/HO-1 signaling possibly influences the diosmetin action in OA. Transfection efficiency was then confirmed by western blotting, which showed that the protein levels of nuclear Nrf2 and HO-1 were markedly reduced in the si-Nrf2-treated chondrocytes si-Nrf2 compared with the cells transfected


with control plasmid (Fig. 5A, B). Interestingly, the downregulated protein levels of nuclear p65 and p-p65 induced by diosmetin were restored by Nrf2 knockdown (Fig. 5C, D). In addition, the elevated protein levels of inflammatory markers (iNOS and COX-2) and ECM degradation markers (MMP-13 and ADAMTS-5) induced by IL-1 $\beta$  were decreased by diosmetin, whereas Nrf2 knockdown counteracted these diosmetin effects (Fig. 5E, F). Moreover, the effect of diosmetin on the levels of inflammatory biomarkers PGE2, nitrite, TNF- $\alpha$ , and IL-6 in the supernatants of IL-1 $\beta$ -treated cells was also abolished by Nrf2 knockdown (Fig. 5G–J). These results suggested that diosmetin may inhibit NF- $\kappa$ B signaling-mediated inflammatory response by activating Nrf2.



**Figure 3.** Diosmetin inactivates NF-κB signaling pathway in IL-1β-treated chondrocytes. **A–D.** Western blotting of NF-κB signaling-related proteins in chondrocytes treated with IL-1β and diosmetin. **E.** Immunofluorescence of p65 expression in chondrocytes treated with IL-1β and diosmetin. \*\*P < 0.01, \*\*\* P < 0.001.



**Figure 4.** Diosmetin activates Nrf2/HO-1 signaling pathway in IL-1β-treated chondrocytes. **A, B.** Western blotting of Nrf2/HO-1 signaling-related proteins in chondrocytes treated with IL-1β and diosmetin. **C.** Immunofluorescence of Nrf2 expression in chondrocytes treated with IL-1β and diosmetin. \*\*P < 0.01, \*\*\*P < 0.001.




**Figure 5.** Diosmetin inhibits IL-1β-induced markers of OA in chondrocytes *via* activation of Nrf2/HO-1 signaling pathway. **A**, **B**. Western blotting of NF-κB and **C**, **D**. Nrf2/HO-1 signaling-related proteins, as well as **(E, F)** inflammation and ECM-related proteins. **G–J**. The levels of inflammatory markers in IL-1β-treated chondrocytes pretreated with diosmetin and transfected with siRNA targeting Nrf2 were assessed by ELISA. \*\*P < 0.01, \*\*\*P < 0.001.


**Diosmetin improves osteoarthritis progression in destabilized medial meniscus (DMM) mouse model**

The OA mouse model was established by the DMM surgery to investigate the role of diosmetin in protecting articular cartilage. According to the results of Safranin O staining, the mice in the DMM group presented early OA-like manifestations, including erosion of the cartilage and massive loss of proteo-

glycan staining with higher OARSI score compared to the control (Fig. 6). In contrast, diosmetin treatment significantly slowed the OA progression in mouse model, which was evidenced by less cartilage erosion, smoother cartilage surface, and more proteoglycan staining than the DMM group. In addition, the increase of OARSI score in the DMM group was suppressed by diosmetin as well, indicating its protective effect against pathological osteoarthritic changes (Fig. 6A, B).



**Figure 6.** Diosmetin improves osteoarthritis progression in destabilization of the medial meniscus mouse model. **A.** Safranin O staining of the articular cartilage from different experimental groups. **B.** Osteoarthritis Research Society International (OARSI) scores of cartilages from different experimental groups. \*\*P < 0.01, \*\*\*P < 0.001.



**Figure 7.** Graphical summary of diosmetin effects on IL-1 $\beta$ -induced murine chondrocytes.



## DISCUSSION

In recent years, increasing plant-based natural agents have caught much attention of researchers and have been confirmed as potential drugs to treat OA, such as Astilbin extracted from *Dimorphandra mollis* [28], murine OA model),  $\beta$ -HIVS derived from *Lithospermum erythrorhizon* [29], murine OA model), and Xanthohumol isolated from hops [30], rat OA model). Diosmetin from *Citrus aurantium L.* has been shown to protect cartilage in a surgically-induced OA mouse model [24]. We for the first time confirmed the anti-OA effect of diosmetin occurs *via* Nrf2/NF- $\kappa$ B signaling pathways. Inflammation has been proved to be a critical factor in the pathogenesis of OA [28]. IL-1 $\beta$  is often used as an inducer of OA in articular chondrocytes, and it was shown to stimulate the secretion of inflammatory cytokines IL-6 and TNF- $\alpha$ , leading to an inflammatory response [29]. Diosmetin is recognized as an anti-inflammatory agent in various diseases. For example, in atopic dermatitis model, it has been reported to inhibit the production of nitric oxide (NO) and reduce the expression of iNOS [30]. Yu *et al.* also demonstrated its inhibitory effect on iNOS expression in cerulein-induced acute pancreatitis [31]. Similarly, we found that the levels of iNOS and COX-2 proteins and their products were elevated in mouse chondrocytes under the stimulation by IL-1 $\beta$ , and their levels were significantly reduced by diosmetin treatment, suggesting the anti-inflammatory action of diosmetin in the IL-1 $\beta$ -induced OA *in vitro* model. It has been reported that PGE<sub>2</sub>, nitrite, TNF- $\alpha$ , and IL-6 are upregulated in OA animal models [32, 33]. PGE<sub>2</sub>, nitrite, TNF- $\alpha$ , and IL-6 are regarded as important catabolic factors that can trigger the degradation of ECM macromolecules in chondrocytes [7, 34]. In our study, diosmetin remarkably reduced levels of those catabolic factors in IL-1 $\beta$ -treated chondrocytes, suggesting that during OA *in vivo* diosmetin may suppress the ECM degradation and protect chondrocytes and articular cartilage against inflammation-induced dysfunction.

Many studies revealed that impaired ECM metabolism also contributes to OA progression [35]. Under IL-1 $\beta$  stimulation in chondrocytes, key ECM degradative enzymes, including MMPs and aggrecanases, are upregulated, and collagen II and aggrecan are downregulated, which contribute to ECM molecules' degradation [29]. Consistently, here, IL-1 $\beta$ -induced murine chondrocytes showed decreased expression of collagen II and aggrecan and increased levels of MMP-13 and ADAMTS-5, while diosmetin reversed the changes, suggesting its suppressive effect on IL-1 $\beta$ -induced ECM degradation.

As an upstream regulatory factor of inflammation-related factors (iNOS, IL-1 $\beta$ , and TNF- $\alpha$  and other) and cartilage-degrading enzymes (ADAMTs and MMPs), NF- $\kappa$ B signaling has been widely reported to participate in inflammatory processes and degradation of ECM in articular cartilage and chondrocytes [36, 37]. After stimulation by IL-1 $\beta$ , the I $\kappa$ B subunit can be activated, resulting in the transport of NF- $\kappa$ B p65 to the nucleus [38, 39]. Diosmetin has been previously reported to exert anti-inflammatory effect by decreasing the p-JNK and p-NF- $\kappa$ B protein expression in the aorta of hypertensive rats [40]. Chen *et al.* reported that diosmetin inhibited inflammatory response in synoviocyte-like MH7A cells by inhibiting Akt and NF- $\kappa$ B pathways, suppressing features of rheumatoid arthritis *in vitro* model [19]. Here, we found that diosmetin reversed the upregulated p-p65, p-I $\kappa$ B $\alpha$ , nuclear p65 expression and downregulated I $\kappa$ B $\alpha$  expression induced by IL-1 $\beta$  in chondrocytes, indicating that probably *via* inactivation of NF- $\kappa$ B signaling diosmetin inhibits markers of inflammation and ECM degradation in IL-1 $\beta$ -exposed chondrocytes.

Nrf2 is demonstrated to be responsible for the inactivation of NF- $\kappa$ B pathway [41], and emerging evidence has implied the key role of Nrf2 pathway in alleviating IL-1 $\beta$ -induced ECM molecule's degradation in chondrocytes *via* the inhibition of NF- $\kappa$ B pathway [42, 43]. Consistent with the previous study showing the effect of diosmetin on activation of Nrf2 in thoracic aorta of hypertensive rats [40], also in our *in vitro* OA model, diosmetin upregulated the expression of Nrf2 and HO-1 protein in IL-1 $\beta$ -treated chondrocytes, which confirmed the beneficial involvement of Nrf2/HO-1 pathway. Furthermore, we were the first to show that Nrf2 knockdown in IL-1 $\beta$ -treated chondrocytes significantly abolished the inhibitory effect of diosmetin on NF- $\kappa$ B signaling activation, inflammation, and ECM molecules' degradation, indicating that diosmetin mitigates OA development *via* the inactivation of NF- $\kappa$ B signaling by promoting Nrf2/HO-1 pathway.

To verify the anti-inflammatory effect of diosmetin elucidated in the *in vitro* OA model, we conducted a histological analysis of the articular cartilage of mice subjected to the classical DMM OA model. The protective effect of diosmetin against IL-1 $\beta$ -induced inflammatory in the *in vitro* model of OA has been confirmed in the *in vivo* model. Diosmetin dampened pathological osteoarthritic changes including cartilage erosion and massive loss of proteoglycans. In addition, the reduced OARSI score was shown in diosmetin-treated OA mice, which confirmed the therapeutic effect of diosmetin in murine OA model.

In conclusion, the current study revealed that diosmetin effectively alleviated OA progression under *in vivo* and *in vitro* conditions acting *via* the inactivation of NF- $\kappa$ B signaling pathway by activating Nrf2/HO-1 signaling (Fig. 7). The results of our study proved that diosmetin could be considered as a potential drug for OA treatment.

## Article information and declarations

### Data availability statement

The datasets used or analyzed during the current study are available from the corresponding author on reasonable request.

### Ethics statement

All animal experiments were performed in accordance with the National Institutes of Health Guidelines for the Care and Use of Laboratory Animals and were approved by the Ethical Review System for Laboratory Animal Welfare of the Wuhan Myhalic Biotechnology Co., Ltd (approval number: HLK-20207112; approval date: 2023.7.26).

### Author contributions

Liang Qian was the main designer of this study. Liang Qian, Chuang Li, Hong Liu, Hui Zhou and Tao Tan collected and analyzed the data. Liang Qian, Chuang Li, Hong Liu, Hui Zhou and Tao Tan drafted the manuscript. All authors read and approved the final manuscript.

### Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

### Acknowledgments

We appreciate all participants who contributed to the study.

### Conflict of interest

The authors declared no competing interests in this study.

## REFERENCES

- Hunter DJ, Bierma-Zeinstra S. Osteoarthritis. *Lancet*. 2019; 393(10182): 1745–1759, doi: [10.1016/S0140-6736\(19\)30417-9](https://doi.org/10.1016/S0140-6736(19)30417-9), indexed in Pubmed: [31034380](https://pubmed.ncbi.nlm.nih.gov/31034380/).
- Glyn-Jones S, Palmer AJR, Agricola R, et al. Osteoarthritis. *Lancet*. 2015; 386(9991): 376–387, doi: [10.1016/S0140-6736\(14\)60802-3](https://doi.org/10.1016/S0140-6736(14)60802-3), indexed in Pubmed: [25748615](https://pubmed.ncbi.nlm.nih.gov/25748615/).
- Zhang W, Ouyang H, Dass C, et al. Current research on pharmacologic and regenerative therapies for osteoarthritis. *Bone Res*. 2016; 4(1), doi: [10.1038/boneres.2015.40](https://doi.org/10.1038/boneres.2015.40).
- Bindu S, Mazumder S, Bandyopadhyay U. Non-steroidal anti-inflammatory drugs (NSAIDs) and organ damage: A current perspective. *Biochem Pharmacol*. 2020; 180: 114147, doi: [10.1016/j.bcp.2020.114147](https://doi.org/10.1016/j.bcp.2020.114147), indexed in Pubmed: [32653589](https://pubmed.ncbi.nlm.nih.gov/32653589/).
- Philp AM, Davis ET, Jones SW. Developing anti-inflammatory therapeutics for patients with osteoarthritis. *Rheumatology (Oxford)*. 2017; 56(6): 869–881, doi: [10.1093/rheumatology/kew278](https://doi.org/10.1093/rheumatology/kew278), indexed in Pubmed: [27498352](https://pubmed.ncbi.nlm.nih.gov/27498352/).
- Kobayashi M, Squires GR, Mousa A, et al. Role of interleukin-1 and tumor necrosis factor alpha in matrix degradation of human osteoarthritic cartilage. *Arthritis Rheum*. 2005; 52(1): 128–135, doi: [10.1002/art.20776](https://doi.org/10.1002/art.20776), indexed in Pubmed: [15641080](https://pubmed.ncbi.nlm.nih.gov/15641080/).
- Kapoor M, Martel-Pelletier J, Lajeunesse D, et al. Role of proinflammatory cytokines in the pathophysiology of osteoarthritis. *Nat Rev Rheumatol*. 2011; 7(1): 33–42, doi: [10.1038/nrrheum.2010.196](https://doi.org/10.1038/nrrheum.2010.196), indexed in Pubmed: [21119608](https://pubmed.ncbi.nlm.nih.gov/21119608/).
- Zheng Y, Zhang R, Shi W, et al. Metabolism and pharmacological activities of the natural health-benefiting compound diosmin. *Food Funct*. 2020; 11(10): 8472–8492, doi: [10.1039/d0fo01598a](https://doi.org/10.1039/d0fo01598a), indexed in Pubmed: [32966476](https://pubmed.ncbi.nlm.nih.gov/32966476/).
- Fu J, Zhang Y, Yan T, et al. Comprehensive characterization of monoclonal antibody by Fourier transform ion cyclotron resonance mass spectrometry. *MAbs*. 2019; 11(1): 106–115, doi: [10.1080/19420862.2018.1525253](https://doi.org/10.1080/19420862.2018.1525253), indexed in Pubmed: [30230956](https://pubmed.ncbi.nlm.nih.gov/30230956/).
- Pakraddoni R, Shukla N, Gupta K, et al. Diosmetin Induces Modulation of Igf-1 and Il-6 Levels to Alter Rictor-Akt-PKC $\alpha$  Cascade in Inhibition of Prostate Cancer. *J Clin Med*. 2021; 10(20), doi: [10.3390/jcm10204741](https://doi.org/10.3390/jcm10204741), indexed in Pubmed: [34682865](https://pubmed.ncbi.nlm.nih.gov/34682865/).
- Guo G, Dong J. Diosmetin attenuates oxidative stress-induced damage to lens epithelial cells via the mitogen-activated protein kinase (MAPK) pathway. *Bioengineered*. 2022; 13(4): 11072–11081, doi: [10.1080/21655979.2022.2068755](https://doi.org/10.1080/21655979.2022.2068755), indexed in Pubmed: [35481411](https://pubmed.ncbi.nlm.nih.gov/35481411/).
- Wang SY, Sun ZL, Liu T, et al. Flavonoids from *Sophora moorcroftiana* and their synergistic antibacterial effects on MRSA. *Phyther Res*. 2014; 28(7): 1071–1076, doi: [10.1002/ptr.5098](https://doi.org/10.1002/ptr.5098), indexed in Pubmed: [24338874](https://pubmed.ncbi.nlm.nih.gov/24338874/).
- Si Q, Shi Y, Huang D, et al. Diosmetin alleviates hypoxia-induced myocardial apoptosis by inducing autophagy through AMPK activation. *Mol Med Rep*. 2020; 22(2): 1335–1341, doi: [10.3892/mmr.2020.11241](https://doi.org/10.3892/mmr.2020.11241), indexed in Pubmed: [32627001](https://pubmed.ncbi.nlm.nih.gov/32627001/).
- Xie B, Pan D, Liu H, et al. Diosmetin protects against obesity and metabolic dysfunctions through activation of adipose estrogen receptors in mice. *Mol Nutr Food Res*. 2021; 65(17): e2100070, doi: [10.1002/mnfr.202100070](https://doi.org/10.1002/mnfr.202100070), indexed in Pubmed: [34223710](https://pubmed.ncbi.nlm.nih.gov/34223710/).
- Juárez-Reyes K, Brindis F, Medina-Campos ON, et al. Hypoglycemic, antihyperglycemic, and antioxidant effects of the edible plant *Anoda cristata*. *J Ethnopharmacol*. 2015; 161: 36–45, doi: [10.1016/j.jep.2014.11.052](https://doi.org/10.1016/j.jep.2014.11.052), indexed in Pubmed: [25490313](https://pubmed.ncbi.nlm.nih.gov/25490313/).
- Gong X, Xiong Li, Bi C, et al. Diosmetin ameliorate type 2 diabetic mellitus by up-regulating *Corynebacterium glutamicum* to regulate IRS/PI3K/AKT-mediated glucose metabolism disorder in KK-Ay mice. *Phytomedicine*. 2021; 87: 153582, doi: [10.1016/j.phymed.2021.153582](https://doi.org/10.1016/j.phymed.2021.153582), indexed in Pubmed: [34091150](https://pubmed.ncbi.nlm.nih.gov/34091150/).
- Shi M, Wang J, Bi F, et al. Diosmetin alleviates cerebral ischemia-reperfusion injury through Keap1-mediated Nrf2/ARE signaling pathway activation and NLRP3 inflammasome inhibition. *Environ Toxicol*. 2022; 37(6): 1529–1542, doi: [10.1002/tox.23504](https://doi.org/10.1002/tox.23504), indexed in Pubmed: [35191607](https://pubmed.ncbi.nlm.nih.gov/35191607/).
- Li HL, Wei YY, Li XH, et al. Diosmetin has therapeutic efficacy in colitis regulating gut microbiota, inflammation, and oxidative stress via the circ-Sirt1/Sirt1 axis. *Acta Pharmacol Sin*. 2022; 43(4): 919–932, doi: [10.1038/s41401-021-00726-0](https://doi.org/10.1038/s41401-021-00726-0), indexed in Pubmed: [34262136](https://pubmed.ncbi.nlm.nih.gov/34262136/).

19. Chen Y, Wang Y, Liu M, et al. Diosmetin exhibits anti-proliferative and anti-inflammatory effects on TNF- $\alpha$ -stimulated human rheumatoid arthritis fibroblast-like synoviocytes through regulating the Akt and NF- $\kappa$ B signaling pathways. *Phytother Res.* 2020; 34(6): 1310–1319, doi: [10.1002/ptr.6596](https://doi.org/10.1002/ptr.6596), indexed in Pubmed: 31833613.
20. Hu S, Huang Y, Chen Y, et al. Diosmetin reduces bone loss and osteoclastogenesis by regulating the expression of TRPV1 in osteoporosis rats. *Ann Transl Med.* 2020; 8(20): 1312, doi: [10.21037/atm-20-6309](https://doi.org/10.21037/atm-20-6309), indexed in Pubmed: 33209892.
21. Hsu YL, Kuo PL. Diosmetin induces human osteoblastic differentiation through the protein kinase C/p38 and extracellular signal-regulated kinase 1/2 pathway. *J Bone Miner Res.* 2008; 23(6): 949–960, doi: [10.1359/jbmr.080219](https://doi.org/10.1359/jbmr.080219), indexed in Pubmed: 18269307.
22. Shao S, Fu F, Wang Z, et al. Diosmetin inhibits osteoclast formation and differentiation and prevents LPS-induced osteolysis in mice. *J Cell Physiol.* 2019; 234(8): 12701–12713, doi: [10.1002/jcp.27887](https://doi.org/10.1002/jcp.27887), indexed in Pubmed: 30515812.
23. Sharma S, Porwal K, Kulkarni C, et al. Diosmin, a citrus fruit-derived phlebotonic bioflavonoid protects rats from chronic kidney disease-induced loss of bone mass and strength without deteriorating the renal function. *Food Funct.* 2022; 13(4): 2184–2199, doi: [10.1039/d1fo03867b](https://doi.org/10.1039/d1fo03867b), indexed in Pubmed: 35119062.
24. Ding H, Ding H, Mu P, et al. Diosmetin inhibits subchondral bone loss and indirectly protects cartilage in a surgically-induced osteoarthritis mouse model. *Chem Biol Interact.* 2023; 370: 110311, doi: [10.1016/j.cbi.2022.110311](https://doi.org/10.1016/j.cbi.2022.110311), indexed in Pubmed: 36563736.
25. Glasson SS, Blanchet TJ, Morris EA. The surgical destabilization of the medial meniscus (DMM) model of osteoarthritis in the 129/SvEv mouse. *Osteoarthritis Cartilage.* 2007; 15(9): 1061–1069, doi: [10.1016/j.joca.2007.03.006](https://doi.org/10.1016/j.joca.2007.03.006), indexed in Pubmed: 17470400.
26. Obeidat AM, Kim SY, Burt KG, et al. A standardized approach to evaluation and reporting of synovial histopathology in two surgically induced murine models of osteoarthritis. *Osteoarthritis Cartilage.* 2024 [Epub ahead of print], doi: [10.1016/j.joca.2024.05.006](https://doi.org/10.1016/j.joca.2024.05.006), indexed in Pubmed: 38823432.
27. Glasson SS, Chambers MG, Van Den Berg WB, et al. The OARSI histopathology initiative — recommendations for histological assessments of osteoarthritis in the mouse. *Osteoarthritis Cartilage.* 2010; 18 Suppl 3: S17–S23, doi: [10.1016/j.joca.2010.05.025](https://doi.org/10.1016/j.joca.2010.05.025), indexed in Pubmed: 20864019.
28. Bonnet CS, Walsh DA. Osteoarthritis, angiogenesis and inflammation. *Rheumatology (Oxford).* 2005; 44(1): 7–16, doi: [10.1093/rheumatology/keh344](https://doi.org/10.1093/rheumatology/keh344), indexed in Pubmed: 15292527.
29. Daheshia M, Yao JQ. The interleukin 1beta pathway in the pathogenesis of osteoarthritis. *J Rheumatol.* 2008; 35(12): 2306–2312, doi: [10.3899/jrheum.080346](https://doi.org/10.3899/jrheum.080346), indexed in Pubmed: 18925684.
30. Lee DH, Park JK, Choi J, et al. Anti-inflammatory effects of natural flavonoid diosmetin in IL-4 and LPS-induced macrophage activation and atopic dermatitis model. *Int Immunopharmacol.* 2020; 89(Pt A): 107046, doi: [10.1016/j.intimp.2020.107046](https://doi.org/10.1016/j.intimp.2020.107046), indexed in Pubmed: 33045572.
31. Yu Ge, Wan R, Yin G, et al. Diosmetin ameliorates the severity of cerulein-induced acute pancreatitis in mice by inhibiting the activation of the nuclear factor- $\kappa$ B. *Int J Clin Exp Pathol.* 2014; 7(5): 2133–2142, indexed in Pubmed: 24966921.
32. Sun S, Yan Z, Shui X, et al. Astilbin prevents osteoarthritis development through the TLR4/MD-2 pathway. *J Cell Mol Med.* 2020; 24(22): 13104–13114, doi: [10.1111/jcmm.15915](https://doi.org/10.1111/jcmm.15915), indexed in Pubmed: 33063931.
33. Zhang M, Zhang R, Zheng T, et al. Xanthohumol attenuated inflammation and ECM degradation by mediating HO-1/C/EBP $\beta$  pathway in osteoarthritis chondrocytes. *Front Pharmacol.* 2021; 12: 680585, doi: [10.3389/fphar.2021.680585](https://doi.org/10.3389/fphar.2021.680585), indexed in Pubmed: 34017261.
34. Cook AE, Cook JL, Stoker AM. Metabolic responses of meniscus to IL-1 $\beta$ . *J Knee Surg.* 2018; 31(9): 834–840, doi: [10.1055/s-0037-1615821](https://doi.org/10.1055/s-0037-1615821), indexed in Pubmed: 29294496.
35. Rahmati M, Nalesso G, Mobasheri A, et al. Aging and osteoarthritis: central role of the extracellular matrix. *Ageing Res Rev.* 2017; 40: 20–30, doi: [10.1016/j.arr.2017.07.004](https://doi.org/10.1016/j.arr.2017.07.004), indexed in Pubmed: 28774716.
36. Marcu KB, Otero M, Olivetto E, et al. NF-kappaB signaling: multiple angles to target OA. *Curr Drug Targets.* 2010; 11(5): 599–613, doi: [10.2174/138945010791011938](https://doi.org/10.2174/138945010791011938), indexed in Pubmed: 20199390.
37. Choi MC, Jo J, Park J, et al. NF- $\kappa$ B signaling pathways in osteoarthritic cartilage destruction. *Cells.* 2019; 8(7), doi: [10.3390/cells8070734](https://doi.org/10.3390/cells8070734), indexed in Pubmed: 31319599.
38. Wang J, Ma J, Gu JH, et al. Regulation of type II collagen, matrix metalloproteinase-13 and cell proliferation by interleukin-1 $\beta$  is mediated by curcumin via inhibition of NF- $\kappa$ B signaling in rat chondrocytes. *Mol Med Rep.* 2017; 16(2): 1837–1845, doi: [10.3892/mmr.2017.6771](https://doi.org/10.3892/mmr.2017.6771), indexed in Pubmed: 28627596.
39. Shu C, Chen J, Lv M, et al. Plumbagin relieves rheumatoid arthritis through nuclear factor kappa-B (NF- $\kappa$ B) pathway. *Bioengineered.* 2022; 13(5): 13632–13642, doi: [10.1080/21655979.2022.2081756](https://doi.org/10.1080/21655979.2022.2081756), indexed in Pubmed: 35653787.
40. Meeapat S, Prasatthong P, Potue P, et al. Diosmetin ameliorates vascular dysfunction and remodeling by modulation of nrf2/ho-1 and p-JNK/p-NF- $\kappa$ B expression in hypertensive rats. *Antioxidants (Basel).* 2021; 10(9), doi: [10.3390/antiox10091487](https://doi.org/10.3390/antiox10091487), indexed in Pubmed: 34573119.
41. Wu S, Liao X, Zhu Z, et al. Antioxidant and anti-inflammation effects of dietary phytochemicals: the Nrf2/NF- $\kappa$ B signalling pathway and upstream factors of Nrf2. *Phytochemistry.* 2022; 204: 113429, doi: [10.1016/j.phytochem.2022.113429](https://doi.org/10.1016/j.phytochem.2022.113429), indexed in Pubmed: 36096269.
42. Miao Z, Dong M, Wang Ze, et al. Linalool inhibits the progression of osteoarthritis via the Nrf2/HO-1 signal pathway both in vitro and in vivo. *Int Immunopharmacol.* 2022; 113(Pt A): 109338, doi: [10.1016/j.intimp.2022.109338](https://doi.org/10.1016/j.intimp.2022.109338), indexed in Pubmed: 36330908.
43. Li W, Tao C, Mao M, et al. The Nrf2/HMGB1/NF- $\kappa$ B axis modulates chondrocyte apoptosis and extracellular matrix degradation in osteoarthritis. *Acta Biochim Biophys Sin (Shanghai).* 2023; 55(5): 818–830, doi: [10.3724/abbs.2023078](https://doi.org/10.3724/abbs.2023078), indexed in Pubmed: 37232576.

*Submitted: 2 April, 2024*

*Accepted after reviews: 7 June, 2024*

*Available as Online first: 21 June, 2024*