

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.



ISSN: 0015-5659

e-ISSN: 1644-3284

A previously unknown variant of the calcaneofibular ligament

Authors: Kacper Ruzik, Anna Czech, Marek Drobniowski, Andrzej Borowski, Łukasz Olewnik

DOI: 10.5603/fm.100002

Article type: Case report

Submitted: 2024-03-26

Accepted: 2024-04-25

Published online: 2024-06-06

This article has been peer reviewed and published immediately upon acceptance. It is an open access article, which means that it can be downloaded, printed, and distributed freely, provided the work is properly cited.

Articles in "Folia Morphologica" are listed in PubMed.

CASE REPORT

A previously unknown variant of the calcaneofibular ligament

Kacper Ruzik^{1,2}, Anna Czech¹, Marek Drobniewski², Andrzej Borowski², Łukasz Olewnik¹

¹*Department of Anatomical Dissection and Donation, Medical University of Lodz, Łódź, Poland*

²*Clinic of Orthopaedic and Paediatric Orthopaedics, Medical University of Lodz, Łódź, Poland*

Address for correspondence: Łukasz Olewnik, MD, PhD, Department of Anatomical Dissection and Donation, Chair of Anatomy and Histology, Medical University of Lodz, ul. Żeligowskiego 7/9, 90–410 Łódź, Poland, e-mail: lukaszolewnik@gmail.com

ABSTRACT

The lateral ankle joint is composed of three ligaments: the anterior talofibular ligament (ATFL), posterior talofibular ligament (PTFL) and calcaneofibular ligament (CFL). The ATFL and CFL demonstrate morphological variation, especially regarding their shape and number of bands. During standard anatomical dissection, an unusual type of triple CFL was observed: the CFL was composed of two bands originating on the lateral malleolus, and the presence of a lateral talocalcaneal ligament (LTC) originating on the talus bone. The insertion point of each band was located on the calcaneal bone. An understanding of these anatomical patterns provides a clearer view of ankle joint biomechanics, and improved the planning and performance of surgical treatment.

Keywords: ankle joint, anatomical variations, calcaneofibular ligament, lateral talocalcaneal ligament

INTRODUCTION

The ankle joint ligament may be divided into three subgroups based on its anatomical position: lateral ligaments, medial ligaments and the ligaments of the tibiofibular syndesmosis

[15]. All demonstrate considerable anatomical variation [6, 10, 11, 15, 17, 20, 21, 26, 39]. The medial side of the ankle joint is created by the deltoid ligament and lateral side by the lateral collateral ligament complex (LCL). The LCL consists of the anterior talofibular ligament (ATFL), posterior talofibular ligament (PTFL) and calcaneofibular ligament (CFL) [14].

CFL fibers course posteroinferiorly from the tip of the lateral malleolus onto the lateral surface of the calcaneal bone. Being bi-articular, the CFL bridges both the talocrural joint and subtalar joint. The primary role of the CFL is to prevent inversion in the neutral or dorsiflexed position and restrain subtalar inversion; it also limits talar tilt within the ankle mortise [41].

All anatomical structures, including muscles or ligaments, exhibit anatomical variation [24, 33–35]. The variations of ligaments may concern the location of the attachments, their morphology and the occurrence of additional bands [27–29]. There are several classifications based on CFL morphology [6, 14, 20, 30]. Most recent classifications were proposed by Ruzik et al. [31, 32].

The most common injury in the locomotor system is ankle joint sprain, which mostly occurs as result of supination with extreme inversion and plantar flexion at the ankle joint. Although the most commonly-injured ligament is the AFL, up to 20 percent of ankle sprains also include CFL damage [2, 10].

This study presents a case report of a double CFL accompanied by the presence of lateral talocalcaneal ligament (LTC). An awareness of the morphological variability of CFL may be beneficial for anatomists, radiologists or orthopedic surgeons.

CASE REPORT

A male cadaver, 70 years old at death, was subjected to routine anatomical dissection for teaching and research purposes. The cadaver had been donated to the Department of Anatomical Dissection and Donation, Medical University of Lodz. The lateral side of the foot was dissected using standard techniques as described previously.

Dissection of the ankle joint began with the removal of the skin of the lateral compartment of the leg. Following this, the subcutaneous tissue and fascia were carefully removed to reveal the fibularis brevis and fibularis longus tendons. The tendons were then removed to display the CFL. After the dissection, a double-banded CFL with presence of lateral talocalcaneal ligament (LTC) was identified. (Fig. 1, 2).

Measurements were taken using an electronic caliper (Mitutoyo Corporation, Kawasaki-shi, Kanagawa, Japan). The measurements were obtained twice by two researchers. The results are given in Table 1.




Figure 1. Case report of double calcaneofibular ligament with lateral talocalcaneal ligament




Figure 2. Schematic drawing of double calcaneofibular ligament with lateral talocalcaneal ligament

Table 1.

Morphometric measurements of the presented case report of double CFL with presence of LTC

	Main band	Second band	LTC
Origin	Lateral malleolus	Lateral malleolus	Talus bone

Insertion	Lateral surface of calcaneal bone	Lateral surface of calcaneal bone	Lateral surface of calcaneal bone
Length	24.77	23.92	26.89
Width at proximal attachment point	2.82	2.40	2.16
Thicknes at proximal attachment point	1.56	1.50	1.51
Width in the center	2.80	2.44	2.10
Thicknes in the center	1.56	1.44	1.51
Width at distal attachment point	2.86	2.48	1.78
Thicknes at distal attachment point	1.52	1.51	1.44

CFL — calcaneofibular ligament; LTC — lateral talocalcaneal ligament

DISCUSSION

To understand the complexity of CFL variations, it is necessary to consider the embryology of the ankle joint. Foot and ankle development begins in the fourth week of fetal life. At that time, the lower limb buds are visible in external rotation. Starting from the eighth week, the rotation decreases, and the feet are set in a near neutral position in the eleventh week. In the tenth week of fetal life, joint cavities with synovial linings begin to appear, followed by the development of ATFL, PTFL and CFL [22, 23]. The former is created by the thickening of the joint capsule. The PTFL and CFL arise from a thickening of loose tissue distant from the ankle joint; following which, the CFL fibers elongate, forming the shape of a ligament [22, 23]. A study by Ruzik et al. [32] on human fetuses found the shape of the ligament to be determined from the onset of development; as such, any potential division into a double-banded CFL or the creation of an additional ligament, such as the LTC, must take place at this early stage.

The CFL runs along the lateral side of the ankle joint and is almost completely covered by the fibularis longus and fibularis brevis tendons. The ligament originates on the anterior part of the lateral malleolus and inserts onto the small tubercle at the posterior aspect of the lateral calcaneus [14].

Like other ligaments of the ankle joint, the CFL is characterized by high morphological variability. In a study based on 39 lower limbs, Burks et al. [6] describe the CFL as a band-shaped ligament; however, they did not note any anatomical variations and only mentioned the LTC without considering its frequency. Trouilloud et al. [37] present a threefold classification of the CFL based on the presence of the LTC; however, they do not describe the morphology of main band of the CFL. Wiersma et al. [40] were among the first to describe the morphological variations of the CFL, reporting a cord-like structure (66%) and a fan shape (34%). Kitsoulis et al. [20] propose a CFL classification based on band number, i.e. one, two or three bands, but do not consider any attachments; the study does not include any photographic documentation of the specified morphological types.

Pereira et al. [30] describe four types of CFL based on 47 lower limbs: Type 1 (21 out of 47 cases) is a band-shaped ligament; Type 2 is a Y-type CFL; Type 3 is a V-shaped CFL; Type 4 includes an associated lateral talocalcaneal ligament. More recent studies by Ruzik et al. [31, 32] based on human fetuses and adult human cadavers divided Type 2 and Type 4 into subtypes with detailed types of origin: for Type 2 (Y-shaped), subtype 2a included both origins located on the lateral malleolus while subtype 2b included origins located on the lateral malleolus and the talus bone. In addition, Type 4, i.e. with additional bands, was divided into three subtypes: Type 4a, with the additional band originating on the lateral malleolus, Type 4b originating on the talus bone, and Type 4c, with two additional bands, as in the present study. Each subtype included a main band originating on the lateral malleolus, with the proximal attachments located on the lateral malleolus. Of all the subtypes, only Type 4b accommodates the lateral talocalcaneal ligament. Prevalence of LTC is reported from 42% to 66% [19]. However, the presented case represents as a new subtype, because it is the first example in which an LTC coexists with a double CFL [31, 32].

The most common injury to the system is ankle sprain occurring during inversion and supination of a plantar-flexed foot. The main stabilizer of the ankle joint, which prevents this type of injury, are the lateral ankle ligaments [12, 26]. Most sprains affect the ATFL; however, the CFL is also ruptured in 20% of cases. While rest, ice, compression and elevation therapy can achieve excellent results, the patient may develop chronic ankle instability (CAI) [3, 4, 38]. CAI is initially treated conservatively with rehabilitation; however, if symptoms last longer than six months, surgical treatment should be considered [9, 18, 36].

While surgical treatment of ankle instability was originally based on non-anatomical techniques, Brostrom [5] developed the first anatomical technique, which was modified by Gould et al. [16]. Many surgical techniques have been developed for the anatomical

reconstruction of ankle ligaments based on the principles described by Brostrom and Gouland [1, 7, 8, 13, 25]. Surgical CAI treatment should be accompanied by detailed diagnostic imaging to assess whether the ATFL is isolated or whether there is a concomitant CFL injury. If both ligaments are ruptured, they must be reconstructed to avoid recurrence. As both open and arthroscopic techniques are used to reconstruct the ligaments, an understanding of the anatomical variants presented herein is needed to ensure optimal treatment results [1, 7, 8, 13, 25].

CONCLUSIONS

An unusual CFL ligament characterized by the co-occurrence of a double CFL and the presence of a LTC was identified. The described case is significant for anatomists, orthopaedic surgeons and radiologists, as its findings can be used to support the correct diagnosis and treatment.

Article information and declarations

Ethics statement

The study protocol was accepted by the Bioethics Committee of the Medical University of Lodz (resolution RNN/242/22/KE). The cadavers were the property of the Department of Anatomical Dissection and Donation, Medical University of Lodz. Informed consent was obtained from all participants before they died.

Author contributions

Kacper Ruzik — data collection and analysis and manuscript writing. **Anna Czech** — data collection and analysis. **Marek Drobniewski** — data analysis and manuscript editing. **Andrzej Borowski** — numerous consultations, observations, and suggestions related to the paper. **Łukasz Olewnik** — data analysis and manuscript editing, data analysis and manuscript writing. All authors have read and approved the manuscript

Funding

The authors have no financial or personal relationship with any third party whose interests could be influenced positively or negatively by the article's content. This research received no specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of interest

The authors declare that they have no competing interests.

References

1. Acevedo JI, Mangone P. Arthroscopic brostrom technique. *Foot Ankle Int.* 2015; 36(4): 465–473, doi: [10.1177/1071100715576107](https://doi.org/10.1177/1071100715576107), indexed in Pubmed: [25743426](https://pubmed.ncbi.nlm.nih.gov/25743426/).
2. van den Bekerom MPJ, Kerkhoffs GM, McCollum GA, et al. Management of acute lateral ankle ligament injury in the athlete. *Knee Surg Sports Traumatol Arthrosc.* 2013; 21(6): 1390–1395, doi: [10.1007/s00167-012-2252-7](https://doi.org/10.1007/s00167-012-2252-7), indexed in Pubmed: [23108678](https://pubmed.ncbi.nlm.nih.gov/23108678/).
3. van den Bekerom MPJ, Sjer A, Somford MP, et al. What is the evidence for rest, ice, compression, and elevation therapy in the treatment of ankle sprains in adults? *J Athl Train.* 2012; 47(4): 435–443, doi: [10.4085/1062-6050-47.4.14](https://doi.org/10.4085/1062-6050-47.4.14), indexed in Pubmed: [22889660](https://pubmed.ncbi.nlm.nih.gov/22889660/).
4. Bleakley CM, O'Connor SR, Tully MA, et al. Effect of accelerated rehabilitation on function after ankle sprain: randomised controlled trial. *BMJ.* 2010; 340: c1964, doi: [10.1136/bmj.c1964](https://doi.org/10.1136/bmj.c1964), indexed in Pubmed: [20457737](https://pubmed.ncbi.nlm.nih.gov/20457737/).
5. Broström L. Sprained ankles. V. Treatment and prognosis in recent ligament ruptures. *Acta Chir Scand.* 1966; 132(5): 537–550, indexed in Pubmed: [5972556](https://pubmed.ncbi.nlm.nih.gov/5972556/).
6. Burks RT, Morgan J. Anatomy of the lateral ankle ligaments. *Am J Sports Med.* 1994; 22(1): 72–77, doi: [10.1177/036354659402200113](https://doi.org/10.1177/036354659402200113), indexed in Pubmed: [8129114](https://pubmed.ncbi.nlm.nih.gov/8129114/).
7. Corte-Real NM, Moreira RM. Arthroscopic repair of chronic lateral ankle instability. *Foot Ankle Int.* 2009; 30(3): 213–217, doi: [10.3113/FAI.2009.0213](https://doi.org/10.3113/FAI.2009.0213), indexed in Pubmed: [19321097](https://pubmed.ncbi.nlm.nih.gov/19321097/).
8. Cottom JM, Rigby RB. The "all inside" arthroscopic Broström procedure: a prospective study of 40 consecutive patients. *J Foot Ankle Surg.* 2013; 52(5): 568–574, doi: [10.1053/j.jfas.2013.02.022](https://doi.org/10.1053/j.jfas.2013.02.022), indexed in Pubmed: [23669003](https://pubmed.ncbi.nlm.nih.gov/23669003/).
9. Dias S, Lewis TL, Alkhalafan Y, et al. Current concepts in the surgical management of chronic ankle lateral ligament instability. *J Orthop.* 2022; 33: 87–94, doi: [10.1016/j.jor.2022.07.006](https://doi.org/10.1016/j.jor.2022.07.006), indexed in Pubmed: [35874042](https://pubmed.ncbi.nlm.nih.gov/35874042/).
10. Doherty C, Delahunt E, Caulfield B, et al. The incidence and prevalence of ankle sprain injury: a systematic review and meta-analysis of prospective epidemiological studies. *Sports Med.* 2014; 44(1): 123–140, doi: [10.1007/s40279-013-0102-5](https://doi.org/10.1007/s40279-013-0102-5), indexed in Pubmed: [24105612](https://pubmed.ncbi.nlm.nih.gov/24105612/).
11. Edama M, Kageyama I, Kikumoto T, et al. Morphological features of the anterior talofibular ligament by the number of fiber bundles. *Ann Anat.* 2018; 216: 69–74, doi: [10.1016/j.aanat.2017.11.001](https://doi.org/10.1016/j.aanat.2017.11.001), indexed in Pubmed: [29196235](https://pubmed.ncbi.nlm.nih.gov/29196235/).
12. Edama M, Takabayashi T, Yokota H, et al. Number of fiber bundles in the fetal anterior talofibular ligament. *Surg Radiol Anat.* 2021; 43(12): 2077–2081, doi: [10.1007/s00276-021-02816-4](https://doi.org/10.1007/s00276-021-02816-4), indexed in Pubmed: [34379153](https://pubmed.ncbi.nlm.nih.gov/34379153/).

13. Espinosa N, Smerek J, Kadakia AR, et al. Operative management of ankle instability: reconstruction with open and percutaneous methods. *Foot Ankle Clin.* 2006; 11(3): 547–565, doi: [10.1016/j.fcl.2006.07.003](https://doi.org/10.1016/j.fcl.2006.07.003), indexed in Pubmed: [16971248](https://pubmed.ncbi.nlm.nih.gov/16971248/).
14. Golanó P, Vega J, de Leeuw PAJ, et al. Anatomy of the ankle ligaments: a pictorial essay. *Knee Surg Sports Traumatol Arthrosc.* 2010; 18(5): 557–569, doi: [10.1007/s00167-010-1100-x](https://doi.org/10.1007/s00167-010-1100-x), indexed in Pubmed: [20309522](https://pubmed.ncbi.nlm.nih.gov/20309522/).
15. Gomes T, Oliva X, Sanchez EV, et al. Anatomy of the ankle and subtalar joint ligaments. *Foot Ankle Clin.* 2023; 28(2): 201–216, doi: [10.1016/j.fcl.2022.12.003](https://doi.org/10.1016/j.fcl.2022.12.003).
16. Gould N, Seligson D, Gassman J. Early and late repair of lateral ligament of the ankle. *Foot Ankle.* 1980; 1(2): 84–89, doi: [10.1177/107110078000100206](https://doi.org/10.1177/107110078000100206), indexed in Pubmed: [7274903](https://pubmed.ncbi.nlm.nih.gov/7274903/).
17. Hayashi K, Tajima G, Oikawa R, et al. Morphology of the proximal tibiofibular joint and ligaments using three-dimensional computed tomography: A cadaveric study. *Knee.* 2023; 45: 27–34, doi: [10.1016/j.knee.2023.08.017](https://doi.org/10.1016/j.knee.2023.08.017), indexed in Pubmed: [37769379](https://pubmed.ncbi.nlm.nih.gov/37769379/).
18. Hertel J, Corbett RO. An updated model of chronic ankle instability. *J Athl Train.* 2019; 54(6): 572–588, doi: [10.4085/1062-6050-344-18](https://doi.org/10.4085/1062-6050-344-18), indexed in Pubmed: [31162943](https://pubmed.ncbi.nlm.nih.gov/31162943/).
19. Higashiyama R, Sekiguchi H, Takata K, et al. Arthroscopic reconstruction of the anterior talofibular ligament, lateral talocalcaneal ligament, and calcaneofibular ligament using a triangle-shaped tendon graft (alc-triangle). *Arthrosc Tech.* 2020; 9(2): e217–e223, doi: [10.1016/j.eats.2019.09.020](https://doi.org/10.1016/j.eats.2019.09.020), indexed in Pubmed: [32099775](https://pubmed.ncbi.nlm.nih.gov/32099775/).
20. Kitsoulis P, Marini A, Pseftinakou A, et al. Morphological study of the calcaneofibular ligament in cadavers. *Folia Morphol.* 2011; 70(3): 180–184, indexed in Pubmed: [21866529](https://pubmed.ncbi.nlm.nih.gov/21866529/).
21. De Leeuw PAJ, Vega J, Karlsson J, et al. The posterior fibulotalocalcaneal ligament complex: a forgotten ligament. *Knee Surg Sports Traumatol Arthrosc.* 2021; 29(5): 1627–1634, doi: [10.1007/s00167-020-06431-5](https://doi.org/10.1007/s00167-020-06431-5), indexed in Pubmed: [33486559](https://pubmed.ncbi.nlm.nih.gov/33486559/).
22. Marchi D. Using the morphology of the hominoid distal fibula to interpret arboreality in *Australopithecus afarensis*. *J Hum Evol.* 2015; 85: 136–148, doi: [10.1016/j.jhevol.2015.06.002](https://doi.org/10.1016/j.jhevol.2015.06.002), indexed in Pubmed: [26142774](https://pubmed.ncbi.nlm.nih.gov/26142774/).
23. Marchi D, Rimoldi A, García-Martínez D, et al. Morphological correlates of distal fibular morphology with locomotion in great apes, humans, and *Australopithecus afarensis*. *Am J Biol Anthropol.* 2022; 178(2): 286–300, doi: [10.1002/ajpa.24507](https://doi.org/10.1002/ajpa.24507), indexed in Pubmed: [36790753](https://pubmed.ncbi.nlm.nih.gov/36790753/).
24. Maślanka K, Zielinska N, Tubbs RS, et al. Five-headed superior omohyoid. *Folia Morphol.* 2023; 82(4): 975–979, doi: [10.5603/FM.a2022.0091](https://doi.org/10.5603/FM.a2022.0091), indexed in Pubmed: [36385428](https://pubmed.ncbi.nlm.nih.gov/36385428/).
25. Matsui K, Takao M, Miyamoto W, et al. Arthroscopic Broström repair with Gould augmentation via an accessory anterolateral port for lateral instability of the ankle. *Arch Orthop Trauma Surg.* 2014; 134(10): 1461–1467, doi: [10.1007/s00402-014-2049-x](https://doi.org/10.1007/s00402-014-2049-x), indexed in Pubmed: [25015792](https://pubmed.ncbi.nlm.nih.gov/25015792/).

26. Matsui K, Takao M, Tochigi Y, et al. Anatomy of anterior talofibular ligament and calcaneofibular ligament for minimally invasive surgery: a systematic review. *Knee Surg Sports Traumatol Arthrosc.* 2017; 25(6): 1892–1902, doi: [10.1007/s00167-016-4194-y](https://doi.org/10.1007/s00167-016-4194-y), indexed in Pubmed: [27295109](https://pubmed.ncbi.nlm.nih.gov/27295109/).
27. Olewnik Ł, Gonera B, Kurtys K, et al. The anterolateral ligament of the knee: a proposed classification system. *Clin Anat.* 2018; 31(7): 966–973, doi: [10.1002/ca.23267](https://doi.org/10.1002/ca.23267), indexed in Pubmed: [30144325](https://pubmed.ncbi.nlm.nih.gov/30144325/).
28. Olewnik Ł, Gonera B, Kurtys K, et al. A proposal for a new classification of the fibular (lateral) collateral ligament based on morphological variations. *Ann Anat.* 2019; 222: 1–11, doi: [10.1016/j.aanat.2018.10.009](https://doi.org/10.1016/j.aanat.2018.10.009), indexed in Pubmed: [30408521](https://pubmed.ncbi.nlm.nih.gov/30408521/).
29. Olewnik Ł, Gonera B, Kurtys K, et al. Classification of the popliteofibular ligament. *Clin Anat.* 2022; 35(3): 375–382, doi: [10.1002/ca.23842](https://doi.org/10.1002/ca.23842), indexed in Pubmed: [35119143](https://pubmed.ncbi.nlm.nih.gov/35119143/).
30. Pereira BS, van Dijk CN, Andrade R, et al. The calcaneofibular ligament has distinct anatomic morphological variants: an anatomical cadaveric study. *Knee Surg Sports Traumatol Arthrosc.* 2020; 28(1): 40–47, doi: [10.1007/s00167-019-05797-5](https://doi.org/10.1007/s00167-019-05797-5), indexed in Pubmed: [31776625](https://pubmed.ncbi.nlm.nih.gov/31776625/).
31. Ruzik K, Gonera B, Borowski A, et al. Anatomic variations of the calcaneofibular ligament. *Foot Ankle Int.* 2024 [Epub ahead of print]: 10711007241241073, doi: [10.1177/10711007241241073](https://doi.org/10.1177/10711007241241073), indexed in Pubmed: [38590202](https://pubmed.ncbi.nlm.nih.gov/38590202/).
32. Ruzik K, Gonera B, Podgórski M, et al. Anatomical variations of the calcaneofibular ligament in human fetuses. *Sci Rep.* 2023; 13(1): 11016, doi: [10.1038/s41598-023-37799-2](https://doi.org/10.1038/s41598-023-37799-2), indexed in Pubmed: [37419905](https://pubmed.ncbi.nlm.nih.gov/37419905/).
33. Ruzik K, Olewnik L, Westrych K, et al. Anatomical variation of co-existing bilaminar tensor of the vastus intermedius muscle and new type of sixth head of the quadriceps femoris. *Folia Morphol.* 2022; 81(4): 1082–1086, doi: [10.5603/FM.a2021.0095](https://doi.org/10.5603/FM.a2021.0095), indexed in Pubmed: [34590299](https://pubmed.ncbi.nlm.nih.gov/34590299/).
34. Ruzik K, Westrych K, Tubbs RS, et al. Case report of the double-headed extensor hallucis longus. *Folia Morphol.* 2023; 82(2): 429–433, doi: [10.5603/FM.a2022.0018](https://doi.org/10.5603/FM.a2022.0018), indexed in Pubmed: [35187631](https://pubmed.ncbi.nlm.nih.gov/35187631/).
35. Ruzik K, Westrych K, Zielinska N, et al. The morphological variability of fibularis tertius origin in human fetuses. *Ann Anat.* 2022; 243: 151920, doi: [10.1016/j.aanat.2022.151920](https://doi.org/10.1016/j.aanat.2022.151920), indexed in Pubmed: [35278660](https://pubmed.ncbi.nlm.nih.gov/35278660/).
36. Shorter AL, Richardson JK, Finucane SB, et al. Characterization and clinical implications of ankle impedance during walking in chronic stroke. *Sci Rep.* 2021; 11(1): 16726, doi: [10.1038/s41598-021-95737-6](https://doi.org/10.1038/s41598-021-95737-6), indexed in Pubmed: [34408174](https://pubmed.ncbi.nlm.nih.gov/34408174/).
37. Trouilloud P, Dia A, Grammont P, et al. Variations in the calcaneo-fibular ligament (lig. calcaneofibulare). Application to the kinematics of the ankle]. *Bull Assoc Anat (Nancy)*. 1988; 72(216): 31–5.
38. Vuurberg G, Hoorntje A, Wink LM, et al. Diagnosis, treatment and prevention of ankle sprains: update of an evidence-based clinical guideline. *Br J Sports Med.* 2018; 52(15): 956, doi: [10.1136/bjsports-2017-098106](https://doi.org/10.1136/bjsports-2017-098106), indexed in Pubmed: [29514819](https://pubmed.ncbi.nlm.nih.gov/29514819/).

39. Wiersma PH, Griffioen F. Variations of three lateral ligaments of the ankle. A descriptive anatomical study. *The Foot*. 1992; 2(4): 218–224, doi: [10.1016/0958-2592\(92\)90051-p](https://doi.org/10.1016/0958-2592(92)90051-p).
40. Wiersma PH, Griffioen F. Variations of three lateral ligaments of the ankle. A descriptive anatomical study. *The Foot*. 1992; 2(4): 218–224, doi: [10.1016/0958-2592\(92\)90051-p](https://doi.org/10.1016/0958-2592(92)90051-p).
41. Yoshizuka H, Kuraoka A. Calcaneofibular ligament may act as a tensioner of peroneal tendons as revealed by a contactless three-dimensional scan system on cadavers. *Sci Rep*. 2022; 12(1): 16650, doi: [10.1038/s41598-022-21115-5](https://doi.org/10.1038/s41598-022-21115-5), indexed in Pubmed: [36198736](https://pubmed.ncbi.nlm.nih.gov/36198736/).