

The influence of open field exposure on neurons containing nitric oxide synthase in the basolateral complex and paracapsular intercalated nerve cells of the rat amygdala

Beata Ludkiewicz, Ilona Klejbor, Beata Domaradzka-Pytel, Sławomir Wójcik, Anna Łuczyńska, Ewa Badowska-Szałewska, Jerzy Dziwiątkowski, Janusz Moryś

Department of Anatomy and Neurobiology, Medical University, Gdańsk, Poland

[Received 14 July 2004; Accepted 14 October 2004]

Our intention in the present study was to ascertain whether NO-producing cells in the basolateral complex (BLC) and paracapsular intercalated nerve cell groups (Ip) of the amygdala are activated in the open field (OF) test. The material consisted of 8 adult rat brains. The OF test was applied throughout 10 min and 90 min before the death of the animals. The brain sections were double stained using the antibodies against c-Fos (marker of neuronal activation) and against nitric oxide synthase (NOS — marker of NO-producing cells).

The neurons containing NOS and those revealing c-Fos activity constituted distinct populations within both the BLC and Ip but NOS-immunoreactive fibres often surrounded the c-Fos-immunoreactive neurons. Our results suggest that (1) neurons of the basolateral complex of the amygdala and paracapsular intercalated islands are involved but probably not crucial for the open field stress response and (2) NOS-immunoreactive cells in the BLC and Ip are not activated after OF exposure.

Key words: stress, c-Fos protein, NO-producing neurons

INTRODUCTION

The stress system co-ordinates the adaptive responses of the organism to stressors of any kind [31]. Stressors can be divided into two categories: (1) systemic, which directly threaten survival and activate the paraventricular hypothalamic nucleus (PVH) through the ascending pathway from the brainstem and (2) processive (emotional, psychological), which activate the cortical and limbic areas before the PVH is activated [28]. The amygdala, throughout its connections, activates the hypothalamic-pituitary-adrenal axis by processive stressors such as an unfriendly and unfamiliar environment [28].

The basolateral complex (BLC), composed of lateral (L) and basolateral nuclei (BL) [5], is a critical component of the amygdaloid body. The lateral nu-

cleus receives most of the cortical and subcortical input from the visual, auditory and somatosensory modalities and may integrate this sensory information [9, 18, 19, 25]. Direct and indirect projections from the lateral nucleus connect it with the central nucleus (Ce) of the amygdala [25] and by this pathway can generate the fear response [19]. The intercalated masses of the amygdala (Ip) [2] surround the BLC. The cell group lying between the basolateral complex and the central nucleus receives inputs from the BLC and sends them to the central nucleus [11], representing an inhibitory interface between the input and output nuclei of the amygdala.

Nitric oxide (NO) is well-known as a gas that acts in the nervous system as a neurotransmitter or a neuromodulator [5]. Neuronal NOS (nNOS) is the

brain-specific isozyme that makes NO in the brain [13, 22]. The role of NO is not confined to a cellular level but is also involved in regulation of the activity of specific physiological systems [17].

It has been demonstrated that NO can mediate the stress responses in the amygdala [16, 29]. In view of this, our intention was to ascertain whether NOS-containing cells in the basolateral complex of the amygdala and paracapsular intercalated nerve cells are activated by the open field (OF) test, a model of an unfamiliar and unfriendly environment. The engagement of these cells in the stress response can be evaluated by the detection of c-Fos protein, commonly recognised as a marker of neuronal activation [15, 26, 30].

MATERIAL AND METHODS

The material consisted of 8 adult rats of the Wistar strain. The experimental group consisted of 5 animals, while the control consisted of 3. Care and treatment of the animals were in accordance with the guidelines for laboratory animals established by the National Institutes of Health as well as by the Local Ethical Committee. The rats of the experimental group were exposed to the open field test throughout 10 minutes. The open field was constructed of a white wooden floor and walls (100 × 100 × 40 cm) and was illuminated with a 500-watt halogen light. The control animals were kept in a home cage. After 90 min all the animals were deeply anaesthetised with lethal doses of Thiopental (80 mg/kg of body weight) and then transcardially perfused with 0.9% solution of NaCl with heparin, followed by 4% paraformaldehyde solution in 0.1 M phosphate buffer (pH 7.4). The brains were postfixed in 4% paraformaldehyde fixative for 3–4 hours, and then kept in 0.1 M phosphate buffer containing 10% sucrose (overnight at 4°C) and 30% sucrose (until sunk). Coronal 40- μ m-thick serial sections of the brain were cut on a JUNG 1800 cryostat (Leica, Germany). The sections were then double stained with the immunohistochemical method. The free floating sections were blocked with 3% normal goat serum (NGS) containing 0.3% Triton X-100 for 1 hour and then incubated with a mixture of rabbit anti-c-Fos antibody (Santa Cruz; diluted 1:1000) together with mouse anti-NOS (Sigma, diluted 1:1500) in 3% NGS for 48 hours in 4°C. After multiple rinses in PBS, sections were incubated (2–3 hours, at room temperature) with a mixture of secondary fluorophore conjugated antibodies: Alexa Fluor 488 goat anti-mouse (Symbios; dilution 1:150) and Cy3-conjugated goat anti-rabbit (Jackson ImmunoResearch; dilution 1: 600).

The immunohistochemically stained slides were examined with fluorescent microscope BX-51 (Olympus, Japan) and the confocal system MicroRadiance (Bio-Rad, UK), equipped with an Argon ion laser and mounted on a light microscope Eclipse 600 (Nikon, Japan), using the software LaserSharp 2000 (Bio-Rad, UK). The confocal laser scanning microscopy (CLSM) images were obtained using 40 × and 60 × oil immersion objective lenses of NA = 1.3 and 1.4, respectively. The optimal iris was used for each magnification. For the reconstruction of the images the program LaserSharp 2000 v. 2.0 (Bio-Rad; UK) was used.

RESULTS

NOS-immunoreactivity

The distribution of NOS-immunoreactive (-ir) neurons in the basolateral complex was similar in all the brains studied. The lateral nucleus showed the highest number of labelled cells. There were considerably less NOS-ir neurons in the basolateral nucleus and many of these cells were located along the ventral and medial borders of the nucleus. There were also numerous stained NOS-ir cells surrounding the basolateral complex — located in the paracapsular intercalated nerve cell groups (Fig. 1).

Intensely stained NOS-ir neurons both in BLC and Ip typically had ovoid or fusiform cell bodies and 3 to 4 primary dendrites, although some were bipolar (Fig. 1A, B).

NOS-ir structures in neuropil comprised fibres and axon terminals. NOS-ir fibres in the lateral nucleus were of two kinds, (1) thin and (2) thick with varicosities (Fig. 1A). The latter were less numerous. In the lateral part of the basolateral nucleus the thin fibres predominated (Fig. 1C).

NOS-ir neurons interposed between the basolateral complex and the central nucleus and located in the adjacent external capsule were intensely stained, with elongated ovoid or fusiform cell bodies and with dendrites oriented parallel to the basolateral complex borders. NOS-ir fibres were intensely stained, thick and with varicosities (Fig. 1B).

c-Fos-immunoreactivity

Single scattered c-Fos positive cells were observed in the basolateral complex of the control rats. Both the number and the distribution of c-Fos-ir cells in the response to the open field exposure were differentiated among individual animals. c-Fos-ir cells were found in the lateral and basolateral nuclei and in the




Figure 1. NOS-immunoreactive cells and fibres (green) and c-Fos-immunoreactive nuclei (red) in the rat amygdala after open field exposure. **A** — lateral nucleus, **B** — paracapsular intercalated nerve cells, **C** — basolateral nucleus. L — lateral nucleus, Ce — central nucleus, BL — basolateral nucleus.

paracapsular intercalated cells (Fig. 1). Their intensity characterised the diverse level of immunolabelling, from high to low. No clearly visible pattern of distribution of c-Fos-ir cells was observed within BLC in the response to the open field exposure, but it seems that they were more often localised in the medial part of the basolateral nucleus (Fig. 1).

Double NOS/c-Fos immunolabelling

The double immunolabelling study revealed that cells containing NOS and those containing c-Fos constituted distinct populations within the BLC (Fig. 1A, C). Very few cells showed co-localisation — they were observed mostly in the medial part of the basolateral nucleus. Among cells of paracapsular intercalated groups no co-localisation with c-Fos was observed (Fig. 1B).

Relatively often we observed thin NOS-ir fibres surrounding cells containing c-Fos-ir nuclei (sometimes they formed basket-like structures; Fig. 1).

DISCUSSION

The results of the present study demonstrate that the nitric oxide synthase-containing neurons in the basolateral complex of the amygdala do not reveal c-Fos activity in response to open field exposure.

The exposure to the open field did not significantly influence either the morphology or the distribution patterns of NOS-ir cells in BLC. They were similar to those reported previously [20, 33].

The morphology and localisation of NOS-ir neurons belonging to the paracapsular intercalated cell groups surrounding BLC suggest that they can be GABAergic neurons [11, 23] receiving glutamergic input from the basolateral complex and sending GABAergic fibres to the central nucleus [11].

The relatively low number of c-Fos-ir cells both in the BLC and paracapsular intercalated islands were confirmed earlier, revealing that after exposure to a processive stressor c-fos expression was more prominent in the medial (Me) than in other amygdaloid nuclei [7, 8, 10, 24, 28]. In contrast to the consistency of Me activation by processive stressors, activation of other amygdaloid nuclei is, according to various authors, differentiated, even for the same stressor. For example, restraint generally elicited very little c-Fos expression in BLC [8], although, in contrast, Bhatnagar and Dallman reported substantial c-fos expression throughout the whole amygdala in the same conditions [3]. In the present study we observed relatively low c-Fos expression in BLC and Ip. However, in the case of an

unconditioned emotional stressor, namely open field exposure, it appears that these structures are in fact not critical. It seems that BLC, Ce and paracapsular intercalated nerve cell groups are more important for the generation of responses of conditioned emotional stressors [8, 32].

The c-Fos-ir cells observed by us in the BLC support the hypothesis that the basolateral complex of the amygdala could be a locus of neuroplasticity underlying fear-based memory formed and stored within this structure [21]. These c-Fos-ir neurons in the BLC probably "mark" their readiness for memory processing. According to Kaczmarek and Chaudhuri [14] and Savonenko et al. [27], c-Fos accumulation is not a mere indicator of functional neural activity but reflects a predisposition to plastic change. This hypothesis is additionally supported by the presence of long-term potentiation (LTP) of excitatory synaptic transmission, which is a form of activity-dependent plasticity which may underlie learning and memory [1, 4]. It has been demonstrated that synapses in the amygdala (especially in the lateral and basolateral nuclei) display LTP [1, 6, 12, 34], which confirms the hypothesis concerning memory-related plasticity in this structure [35]. It has also been revealed that NO plays a role in facilitating the induction of LTP in the medial amygdala [1]. It is known, that neurons cannot sequester NO; thus the key to regulating NO activity is to control NO synthesis [13]. In this case, the question arises as to why NOS-ir neurons in BLC are hardly activated after open field stress. Some reasons have been put forward by Abe et al. [1]. According to them, long-term potentiation in the lateral amygdala is NO-independent. Additionally, the presence of NOS-ir endings located around the cells containing c-Fos-ir nuclei suggests that the nitric oxide may influence (modulate) the action of stress-activated neurons of BLC.

CONCLUSION

Our results suggest that (1) neurons of the basolateral amygdaloid complex and paracapsular intercalated islands are involved but probably not crucial for open field stress response and (2) NOS-ir cells in these structures are not activated after open field exposure.

ACKNOWLEDGEMENT

The authors wish to thank Ms Sylwia Scisłowska for preparation of the illustration. This study was supported by funds from the Committee of Scientific Research, CSR Grant No. 4 P05A 068 22.

REFERENCES

1. Abe K, Watanabe Y, Saito H (1996) Differential role of nitric oxide in long-term potentiation in the medial and lateral amygdala. *Eur J Pharmacol*, 297: 43–46.
2. Alheid GF, de Olmos JS, Beltramino CA (1995) Amygdala and extended amygdala. In: Paxinos G (ed.). *The rat nervous system*. 2nd ed. Academic Press, San Diego: 495–578.
3. Bhatnagar S, Dallman M (1998) Neuroanatomical basis for facilitation of hypothalamic-pituitary-adrenal responses to a novel stressor after chronic stress. *Neuroscience*, 84: 1025–1039.
4. Bliss TV, Collingridge GL (1993) A synaptic model of memory: long-term potentiation in the hippocampus. *Nature*, 361: 31–39.
5. Bredt DS, Snyder SH (1992) Nitric oxide, a novel neuronal messenger. *Neuron*, 8: 3–11.
6. Chapman PF, Kairiss EW, Keenan CL, Brown TH (1990) Long-term synaptic potentiation in the amygdala. *Synapse*, 6: 271–278.
7. Cullinan WE, Herman JP, Battaglia DF, Akil H, Watson SJ (1995) Pattern and time course of immediate early gene expression in rat brain following acute stress. *Neuroscience*, 64: 477–505.
8. Dayas CV, Buller KM, Day TA (1999) Neuroendocrine responses to an emotional stressor: evidence for involvement of the medial but not the central amygdala. *Eur J Neurosci*, 11: 2312–2322.
9. Doron NN, LeDoux JE (1999) Organization of projections to the lateral amygdala from auditory and visual areas of the thalamus in the rat. *J Comp Neurol*, 412: 383–409.
10. Duncan GE, Knapp DJ, Breese GR (1996) Neuroanatomical characterization of Fos induction in rat behavioral models of anxiety. *Brain Res*, 713: 79–91.
11. Fuxe K, Jacobsen KX, Hoistad M, Tinner B, Jansson A, Staines WA, Agnati LF (2003) The dopamine D1 receptor-rich main and paracapsular intercalated nerve cell groups of the rat amygdala: relationship to the dopamine innervation. *Neuroscience*, 119: 733–746.
12. Gean PW, Chang FC, Huang CC, Lin JH, Way LJ (1993) Long-term enhancement of EPSP and NMDA receptor-mediated synaptic transmission in the amygdala. *Brain Res Bull*, 31: 7–11.
13. Joo KM, Chung YH, Shin CM, Lee YJ, Cha CI (2004) Region-specific alterations of neuronal nitric oxide synthase (nNOS) expression in the amygdala of the aged rats. *Brain Res*, 999: 231–236.
14. Kaczmarek L, Chaudhuri A (1997) Sensory regulation of immediate-early gene expression in mammalian visual cortex: implications for functional mapping and neural plasticity. *Brain Res Rev*, 23: 237–256.
15. Kovacs KJ (1998) c-Fos as a transcription factor: a stressful (re)view from a functional map. *Neurochem Int*, 33: 287–297.
16. Krukoff TL, Khalili P (1997) Stress-induced activation of nitric oxide-producing neurons in the rat brain. *J Comp Neurol*, 377: 509–519.
17. Krukoff TL (1999) Central actions of nitric oxide in regulation of autonomic functions. *Brain Res Rev*, 30: 52–65.
18. Linke R, De Lima AD, Schwegler H, Pape HC (1999) Direct synaptic connections of axons from superior colliculus with identified thalamo-amygdaloid projection neurons in the rat: Possible substrates of a subcortical visual pathway to the amygdala. *J Comp Neurol*, 403: 158–170.
19. Malkani S, Rosen JB (2000) Specific induction of early growth response gene 1 in the lateral nucleus of the amygdala following contextual fear conditioning in rats. *Neuroscience*, 97: 693–702.
20. McDonald AJ, Payne DR, Mascagni F (1993) Identification of putative nitric oxide producing neurons in the rat amygdala using NADPH-diaphorase histochemistry. *Neuroscience*, 52: 97–106.
21. McGaugh JL (2002) Memory consolidation and the amygdala: a systems perspective. *Trends Neurosci*, 25: 456.
22. Moncada S, Higgs A, Furchgott R (1997) International Union of Pharmacology Nomenclature in Nitric Oxide Research. *Pharmacol Rev*, 49: 137–142.
23. Nitecka L, Ben-Ari Y (1987) Distribution of GABA-like immunoreactivity in the rat amygdaloid complex. *J Comp Neurol*, 266: 45–55.
24. Pezzone MA, Lee WS, Hoffman GE, Rabin BS (1992) Induction of c-Fos immunoreactivity in the rat forebrain by conditioned and unconditioned aversive stimuli. *Brain Res*, 597: 41–50.
25. Pitkänen A, Savander V, LeDoux JE (1997) Organization of intra-amygdaloid circuitries in the rat: an emerging framework for understanding functions of the amygdala. *Trends Neurosci*, 20: 517–523.
26. Sagar SM, Sharp FR, Curran T (1988) Expression of c-fos protein in brain: metabolic mapping at the cellular level. *Science*, 240: 1328–1331.
27. Savonenko A, Filipkowski RK, Werka T, Zielinski K, Kaczmarek L (1999) Defensive conditioning-related functional heterogeneity among nuclei of the rat amygdala revealed by c-Fos mapping. *Neuroscience*, 94: 723–733.
28. Senba E, Ueyama T (1997) Stress-induced expression of immediate early genes in the brain and peripheral organs of the rat. *Neurosci Res*, 29: 183–207.
29. Seo DO, Rivier C (2001) Microinfusion of a nitric oxide donor in discrete brain regions activates the hypothalamic-pituitary-adrenal axis. *J Neuroendocrinol*, 13: 925–933.
30. Sheng M, Greenberg ME (1990) The regulation and function of c-fos and other immediate early genes in the nervous system. *Neuron*, 4: 477–485.
31. Tsigos C, Chrousos GP (2002) Hypothalamic-pituitary-adrenal axis, neuroendocrine factors and stress. *J Psychosom Res*, 53: 865–871.
32. Van de Kar LD, Piechowski RA, Rittenhouse PA, Gray TS (1991) Amygdaloid lesions: differential effect on conditioned stress and immobilization-induced increases in corticosterone and renin secretion. *Neuroendocrinology*, 54: 89–95.
33. Vincent SR, Kimura H (1992) Histochemical mapping of nitric oxide synthase in the rat brain. *Neuroscience*, 46: 755–784.
34. Watanabe Y, Saito H, Abe K (1995) Nitric oxide is involved in long-term potentiation in the medial but not lateral amygdala neuron synapses in vitro. *Brain Res*, 688: 233–236.
35. Yaniv D, Desmedt A, Jaffard R, Richter-Levin G (2004) The amygdala and appraisal processes: stimulus and response complexity as an organizing factor. *Brain Res Rev*, 44: 179–186.