

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.



**ISSN:** 0015-5659

**e-ISSN:** 1644-3284

## **The complete anatomy of the transverse facial artery: a computed tomography angiography analysis**

**Authors:** Mateusz Trzeciak, Alicia del Carmen Yika, Kinga Gładys, Patryk Ostrowski, Michał Bonczar, Michał Goncerz, Wadim Wojciechowski, Mateusz Koziej, Jerzy Walocha, Artur Pasternak

**DOI:** 10.5603/fm.97002

**Article type:** Original article

**Submitted:** 2023-08-17

**Accepted:** 2023-10-16

**Published online:** 2023-10-30

This article has been peer reviewed and published immediately upon acceptance. It is an open access article, which means that it can be downloaded, printed, and distributed freely, provided the work is properly cited.

Articles in "Folia Morphologica" are listed in PubMed.

## **The complete anatomy of the transverse facial artery: a computed tomography angiography analysis**

Mateusz Trzeciak<sup>1</sup>, Alicia del Carmen Yika<sup>1</sup>, Kinga Gładys<sup>1</sup>, Patryk Ostrowski<sup>1,2</sup>, Michał Bonczar<sup>1,2</sup>, Michał Goncerz<sup>1</sup>, Wadim Wojciechowski<sup>3</sup>, Mateusz Koziej<sup>1,2</sup>, Jerzy Walocha<sup>1,2</sup>, Artur Pasternak<sup>1</sup>

<sup>1</sup>Department of Anatomy, Jagiellonian University Medical College, Krakow, Poland

<sup>2</sup>Youthoria, Youth Research Organization, Krakow, Poland

<sup>3</sup>Department of Radiology, Jagiellonian University Medical College, Krakow, Poland

Address for correspondence: Mateusz Trzeciak, Department of Anatomy, Jagiellonian University Medical College, Mikołaja Kopernika 12, 33-332 Kraków, Poland, e-mail: trzeciakmateusz6@gmail.com

### **Abstract**

**Background:** The transverse facial artery (TFA) provides blood supply to various structures of the lateral face, including the parotid gland, parotid duct, masseter muscle, and facial skin. Knowledge about its anatomy is of utmost importance in various plastic and reconstructive procedures.

**Materials and methods:** The results of 55 (110 hemifaces) consecutive patients who underwent head and neck computed tomography angiography (CTA) in the Department of Radiology of the Jagiellonian University Medical College, Cracow, Poland, were evaluated in July 2022.

**Results:** TFA was found to originate from STA in 84 of the cases (95.5%), whereas from ECA only in 4 of the cases (4.5%). Median length of the TFA was found to be 43.39 mm (LQ = 38.53 ; UQ = 46.37). The median TFA diameter, at its origin, was established at 2.26 mm

(LQ = 1.93 ; UQ = 2.54). Median TFA cross-sectional area, at its origin, was set to be 2.54 mm (LQ = 1.67 ; UQ = 3.10).

**Conclusions:** The TFA has an important role in the arterial blood supply to the face, and the present study has demonstrated its stable prevalence in the lateral face region. The most frequent origin of the artery was from the STA; however, it also originated from the ECA in some cases. Moreover, the topographic relationships between the TFA and nearby arteries and anatomical landmarks were measured and analyzed.

**Keywords:** transverse facial artery, face, plastic surgery, perforator flap, surgery

## INTRODUCTION

The transverse facial artery (TFA) originates most commonly as a singular vessel from the superficial temporal artery (STA), within the parotid gland. It traverses the face in a superficial manner, passing over the masseter muscle. Its course is approximately a finger's breadth below the zygomatic arch (Figure 1, Figure 2). Subsequently, the artery divides into multiple branches, which provide blood supply to various structures, including the parotid gland, parotid duct, masseter muscle, and facial skin. Additionally, it forms anastomoses with branches of the facial artery (FA) [9].

Numerous studies have demonstrated the variability in the anatomy of the TFA [7, 16]. As mentioned earlier, the TFA is said to originate from the STA; however, reports have shown that the artery may also arise from the external carotid artery (ECA) or its terminal bifurcation in up to 35% of the cases [4]. Moreover, a study by Yang et al. [16] demonstrated that the number of TFAs is also variable, ranging from one to three arteries. The spatial association between the TFA and nearby structures has also been analyzed, especially between the said artery and the branches of the facial nerve [16].

Normality in anatomy can be considered relative (linked to a specific set of assumptions) and is inferred from numerous repeated observations [19]. Having appropriate knowledge of the anatomy of the TFA is of immense importance in plastic and reconstructive surgery. In recent decades, the utilization of fillers and botulinum toxin to enhance facial features and improve

the jawline has gained significant traction. However, the elevated frequency of injections may give rise to infrequent yet severe complications, notably an emerging trend of blindness reported in the literature [2, 3, 12]. The underlying mechanism behind this debilitating complication involves the injection of the toxin into an artery, such as the TFA in the lateral face, subsequently leading to the obstruction of the central retinal artery. Consequently, this occlusion disrupts the blood supply to the optic nerve, resulting in impaired vision and potential blindness [13]. Furthermore, TFA perforator flaps have been utilized in reconstructions of the zygomaticofacial region after tumor resections [18]. Hence, knowing the topographical anatomy of the TFA is of incredible importance. Therefore, the main goal of the present study was to analyze the complete anatomy of this artery and provide surgeons with useful morphometric data between the TFA and nearby structures. It is hoped that the present study may help physicians with localizing the TFA efficiently during various aesthetic and reconstructive procedures and, with that, decrease the potential of iatrogenic injury to this vessel.

## **MATERIALS AND METHODS**

### Approval of the Bioethical Committee

The research protocol was submitted for evaluation and approved by the Jagiellonian University Bioethical Committee, Cracow, Poland (1072.6120.51.2022). The study was carried out according to the allowed criteria during the next stages.

### Study Group

The results of 55 consecutive patients who underwent head and neck computed tomography angiography (CTA) in the Department of Radiology of the Jagiellonian University Medical College, Cracow, Poland, were evaluated in July 2022. Each CTA was evaluated bilaterally; therefore, a total of 110 hemifaces were initially evaluated. Exclusion criteria were established as follows: (1) head, neck, or thoracic trauma affecting the course or the morphometry of the TFA or its close anatomical area, (2) significant artifacts that prevented accurate and precise imaging and/or measurement of the TFA or its close anatomical area, (3) low quality and illegible images, and (4) significant lack of filling the whole arterial system with contrast. In cases where only a segment of the CTA exhibited any of the mentioned

abnormalities, without impacting the opposite side, the assessment of the other TFA was conducted. A substantial majority (20 cases) of the omitted hemifaces were ineligible for analysis due to prominent artifacts. The other two instances were disregarded to ensure data impartiality owing to their subpar image quality. In the end, a sum of 88 hemifaces fulfilled the stipulated inclusion criteria.

## Results Acquisition

All head and neck CTA were performed on a 128-slice scanner CT (Philips Ingenuity CT, Philips Healthcare). The main CT imaging parameters were the following: collimation/increase: 0.625 / 0.3 mm; tube current: 120 mAs; field of view: 210 mm; matrix size: 512 × 512.

All patients received intravenous administration of contrast material at a dose of 1 ml/kg (standard dose). A non-ionic contrast medium (CM) containing 350 mg of iodine per ml was used (Jowersol 741 mg/ml, Optiray®, Guerbet, France). The acquisition of CT data was initiated using a real-time bolus tracking technique (Philips Healthcare) with the region of interest (ROI) placed in the ascending aorta. CM was injected intravenously using a power injector at a flow rate of 5 ml / s. This was immediately followed by an injection of 40 ml of saline solution at the same flow rate. Following injection of CM and saline, image acquisition was automatically started with a 2 s delay when the attenuation trigger value reached a threshold of 120 Hounsfield units (HU). Scanning was performed in the caudocranial direction. The CTA examination was started at the level of the aortic arch up to the circle of Willis.

The CTAs were analyzed on a dedicated workstation in the Anatomical Department of Jagiellonian University Medical College, Cracow, Poland. To ensure the highest possible quality of the visualizations and measurements and minimize potential bias, Materialise Mimics Medical version 21.0 software (Materialise NV, Leuven, Belgium) software was used. 3-dimensional (3D) reconstructions of each scan were developed, employing a set of settings, severally adjusted to each scan. A volume rendering opacity range oscillated from 25 to 75 HU for the lower limit and up to 135 HU for a higher limit. The range was individually adjusted to each TT after a visual investigation.

## Evaluation and Measurements

At the commencement of each assessment, every TFA has been thoroughly visualized. Following that, a series of measurements was recorded within each TFA by two distinct researchers working individually. Subsequently, an average was computed, considering both sets of findings. All measurements were rounded off to two decimal places. Morphometric features of the TFA and/or its associated anatomical area were assessed in 26 categories, listed in Table 2.

## Statistical Analysis

Statistical analysis was performed with STATISTICA v13.1 (StatSoft Inc., Tulsa, OK, USA). The frequency and percentages presented qualitative features. The Shapiro-Wilk test was used to assess the normal distribution. Quantitative characteristics were presented by medians and upper and lower quartiles (UQ, LQ), as well as means and standard deviation (SD), depending on the verified normality of the data. Statistical significance was defined as  $p < 0.05$ . Mann-Whitney and Wilcoxon signed-rank tests were used to establish potential differences between groups. The Spearman rank correlation coefficient was used to determine possible correlations between the parameters.

## RESULTS

The general characteristics of the patient group are presented in Table 1. The TFA was found to originate from the STA in 84 of the cases (95.5%), whereas from ECA only in 4 of the cases (4.5%). The median length of the TFA was found to be 43.39 mm (LQ = 38.53 ; UQ = 46.37). The median TFA diameter, at its origin, was established at 2.26 mm (LQ = 1.93 ; UQ = 2.54). The median TFA cross-sectional area, at its origin, was set to be 2.54 mm (LQ = 1.67 ; UQ = 3.10). Median TFA branching angle was found to be 74.55 degrees (LQ = 59.09 ; UQ = 89.70). The median distance between the origin of the MA and the origin of the TFA was found to be 6.36 mm (LQ = 5.11 ; UQ = 8.04). The median distance between the origin of the TFA and the origin of the STA was set to be 51.81 mm (LQ = 35.90 ; UQ = 65.06). The median distance between the origin of the OA and the origin of the TFA was established at 46.48 (LQ = 39.33 ; UQ = 52.54). The median distance between the origin of the FA and the origin of the TFA was found to be 47.74 mm (LQ = 43.59 ; UQ = 53.36). The median

distance between the origin of the PAA to the origin of the TFA was established at 23.14 mm (LQ = 17.22 ; UQ = 27.14). The median distance between the origin of the TFA to the lower border of the Zygomatic Arch was set to be 24.16 mm (LQ = 21.94 ; UQ = 27.38). All of the results mentioned above, and more detailed ones in all categories, can be found in Table 2. Statistically significant differences ( $p < 0.05$ ) between females and males were found in 16 categories. There was a statistically significant difference in the length of the TFA between females and males. However, this difference was not found regarding the origin of the TFA at its origin. More detailed results with respect to the patient's sex and the performed measurements are demonstrated in Table 3.

Lastly, differences between patients' sides and correlations between parameters and patients' age were studied. However, there were no statistical differences ( $p > 0.05$ ) between patients' sides in any of the parameters. Furthermore, no statistically significant correlation between any of the parameters and patients' age was found.

## DISCUSSION

The TFA is said to originate from the STA; however, various reports have demonstrated that this is not always the case. In a study conducted by Koziej et al. [7], the TFA originated from the STA in the majority of the cases (91.7%). Nevertheless, it was stated that the artery also originated from the ECA (3.1%), as well as from a common trifurcation point along with the maxillary and superficial temporal arteries (4.7%). Interestingly, they also demonstrated a case where the TFA originated from the maxillary artery. On the other hand, in the study presented by Yang et al. [16], the TFA originated from the STA in all cases. Interesting cases of the TFA were also described by Basar et al. [1], namely, the TFA originating bilaterally from a common trunk with an artery that supplied the parotid gland. In the present study, the said vessel was present in all of the patients and originated most commonly from the STA (95.5%); however, an origin from the ECA was reported in four cases (4.5%). Moreover, the number of TFAs in the current study was constant, where only one artery was present in every patient. This differs from the results of other studies; Yang et al. [16] reported double and triple TFAs to be present in 25% and 4.5% of the cadavers, respectively. On the other hand, Koziej et al. [7] reported a significantly lower frequency of double TFAs (4.7%); and no triple TFAs were observed. Interestingly, Yang et al. [16] demonstrated anastomoses

between the TFA and the FA in one male cadaver; however, this was observed in our study. A possible explanation for this may be that small anastomoses between arteries may not contain as much contrast as larger vessels; hence, their visibility would be diminished on CTAs. On the other hand, the aforementioned study presented this occurrence in only one side out of 44 sides. Therefore, the overall prevalence of anastomoses of considerable size between the TFA and the FA may just be low in the general population, as this was neither demonstrated in the study by Koziej et al. [7] where 200 cases were analyzed.

The relationship to nearby structures, such as the zygomatic arch and nearby arteries, has been investigated in various studies. Basar et al. [1] demonstrated that the distance between the inferior region of the zygomatic arch and the TFA ranged from 3.4 to 28.6 mm (mean:  $15.49 \pm 7.56$  mm). Interestingly, Koziej et al. [7] presented higher results, with a distance of  $20.9 \pm 4.1$  mm. The results of the present study show an even larger distance of  $24.63 \pm 4.1$  mm. This data further proves that the TFA is located approximately a finger-width under the zygomatic arch. This may be a reliable rule that surgeons may follow during surgical procedures of the lateral face.

The dominance of the TFA over the FA has been reported in the available literature. A case of bilateral hypoplastic FAs and large TFAs was presented by Olson et al. [10]. In the case report, it was stated that the TFA supplied the whole midface region, giving off the superior labial artery, the lateral nasal artery, and terminating as the angular artery. The hypoplastic FAs gave off the inferior labial artery bilaterally. Interestingly, anastomoses between the branches of the TFA, FA, and the infraorbital artery were also observed. Koziej et al. [7] demonstrated only one case of a dominant TFA out of 200 cases. In another case study, Yotova et al. [17] reported a similar observation where a TFA provided compensation for a bilaterally underdeveloped FA. However, they noted that upon reaching the superficial face, the FA split into two unusual branches, ultimately terminating in the buccinator muscle. The compensatory blood flow from the TFA supported the superior labial, lateral nasal, and angular arteries. Moreover, the study did not mention the presence of an inferior labial artery. Additionally, Tubbs et al. conducted a study reporting a remarkable specimen of a large TFA compensating for the complete absence of the corresponding FA on the left side, though it was a unilateral abnormality [14]. In the present study, the FA dominated over the TFA in all of the cases. These variations of the arterial vasculature of the face develop during embryological development. The formation of aortic arches occurs during the fourth and fifth



weeks of gestation, originating from the aortic sac. Subsequently, the third aortic arch contributes to the development of the ECA, which, in turn, gives rise to the FA, and lateral to the TFA as the STA [5, 6, 8].

Having adequate knowledge about the anatomy of the TFA is of immense importance in various plastic and reconstructive procedures. The TFA has various implications in different reconstructive procedures, such as the TFA perforator flap that was extensively described by Zhang et al. [18]. In the study, 18 elderly patients with facial tumors underwent Mohr surgery, and the resection area was reconstructed using V-Y and propeller TFA perforator flaps. All of the 18 flaps survived, with proper cheek and facial symmetry, as well as proper mouth opening. Hence, the authors of the said study concluded that this reconstructive flap has a high clinical value in the treatment of zygomaticofacial tumors. Moreover, flaps that have been pedicled on the TFA have been also used for reconstructions of intraoral defects and the lower eyelid [11, 15]. Over the past few decades, there has been a growing trend in using fillers and botulinum toxin to enhance facial features and improve the jawline [12]. However, it has also raised concerns due to rare, yet severe complications that have been reported, specifically a concerning trend of blindness [2, 12]. These complications have been observed when the cosmetic filler or botulinum toxin is accidentally injected into an artery, such as the TFA in the lateral face. This subsequently leads to the obstruction of the central retinal artery, disrupting the blood supply to the optic nerve. The consequence is impaired vision and, in some unfortunate cases, potential blindness [13].

Several limitations are associated with the current study. Radiological imaging solely permits the assessment of hemodynamically proficient arteries. Consequently, this aspect could potentially introduce a considerable degree of bias when appraising anatomical deviations within the TFA and other arterial structures. Although not without limitations, our study attempts to thoroughly analyze the anatomy of the TFA using methods that meet the requirements of evidence-based anatomy.

## **CONCLUSIONS**

The most frequent origin of the TFA was from the STA; however, it also originated from the ECA in some cases. Moreover, the topographic relationships between the TFA and nearby arteries and anatomical landmarks were measured and analyzed. It is hoped that the results of

the present study can assist physicians in accurately localizing the TFA during various aesthetic and reconstructive procedures. Having a comprehensive understanding of the arterial blood supply of the lateral face can play a crucial role in minimizing the risk of iatrogenic injury to these vessels. ~~This knowledge is essential in ensuring better outcomes for patients undergoing medical procedures in this region.~~

**Abbreviations:** TFA - Transverse Facial Artery; ECA - External Carotid Artery; STA - Superficial Temporal Artery; CCA - Common Carotid Artery; MA - Maxillary Artery; OA - Occipital Artery; ICA - Internal Carotid Artery; PAA - Posterior Auricular Artery

### **Statements and Declarations:**

**Funding:** The author(s) received no financial support for the research, authorship, and/or publication of this article.

**Competing Interests:** The author(s) declare no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

**Data and Code Availability Statement:** The data that support the findings of this study are available from the corresponding author, upon reasonable request.

### **References**

1. Baçar R, Sargon MF, Tekdemir I, et al. Accurate course and relationships of the transverse facial artery in human cadavers. *Morphologie*. 2004; 88(283): 191–195, doi: 10.1016/S1286-0115(04)98148-9.
2. Chatrath V, Banerjee PS, Goodman GJ, et al. Soft-tissue Filler–associated Blindness: A Systematic Review of Case Reports and Case Series. *Plast Reconstr Surg Glob Open*. 2019; 7(4): e2173, doi: 10.1097/GOX.0000000000002173.
3. Chun BY, Kim SY. Acute visual loss after botulinum toxin A injection in the masseter muscle. *Int Ophthalmol*. 2018; 38(3): 1339–1342, doi: 10.1007/s10792-017-0569-4.
4. Cormack G, Lamberty B. *The Arterial Anatomy of Skin Flaps*. Churchill Livingstone, Edinburgh, London, Melbourne and New York 1989.

5. Hanneman K, Newman B, Chan F. Congenital Variants and Anomalies of the Aortic Arch. *RadioGraphics*. 2017; 37(1): 32–51, doi: 10.1148/rg.2017160033-
6. Koziej M, Bonczar M, Ostrowski P, et al. Termination points of the facial artery—A meta-analysis. *Clinical Anatomy*. 2022; 35(4): 469–476, doi: 10.1002/ca.23817-
7. Koziej M, Polak J, Wnuk J, et al. The transverse facial artery anatomy: Implications for plastic surgery procedures. *PLoS One*. 2019; 14(2): e0211974, doi: 10.1371/journal.pone.0211974-
8. Koziej M, Trybus M, Hołda M, et al. Anatomical Map of the Facial Artery for Facial Reconstruction and Aesthetic Procedures. *Aesthet Surg J*. 2019; 39(11), doi: 10.1093/asj/sjz028-
9. Moore KL, Dalley AF, Agur A. *Clinically oriented anatomy* (8th ed.). Lippincott Williams and Wilkins. 2017-
10. Olson C, Tan Y, Campbell M. A Unique Bilateral Anatomical Variation of the Transverse Facial Artery: A Case Report. *Cureus*. 2022, doi: 10.7759/cureus.30511-
11. Pompei B, Pollastri G, Molteni G, et al. The New Transverse–Facial Artery Musculomucosal Flap for Intraoral Reconstructions. *Plast Reconstr Surg Glob Open*. 2016; 4(3): e652, doi: 10.1097/GOX.0000000000000632-
12. Toure G, Nguyen T-M, Vlavanou S, et al. Transverse facial artery: Its role in blindness after cosmetic filler and botulinum toxin injections. *Journal of Plastic, Reconstructive & Aesthetic Surgery*. 2021; 74(8): 1862–1869, doi: 10.1016/j.bjps.2020.12.042-
13. Townshend A. Blindness After Facial Injection. *J Clin Aesthet Dermatol*. 2016; 9(12): E5–E7-
14. Tubbs RS, Salter EG, Oakes WJ. Unilateral agenesis of the facial artery with compensation by a giant transverse facial artery. *Folia Morphol (Warsz)*. 2005; 64(3): 226–8-
15. Yamakawa S, Suda S, Hayashida K. A new lower eyelid reconstruction using transverse facial artery perforator flap based on an anatomical study. *Journal of Plastic, Reconstructive & Aesthetic Surgery*. 2023; 77: 39–45, doi: 10.1016/j.bjps.2022.11.046-
16. Yang H-J, Gil Y-C, Lee H-Y. Topographical anatomy of the transverse facial artery. *Clinical Anatomy*. 2009; NA-NA, doi: 10.1002/ca.20880-
17. Yotova N, Novakov S, Fasova A, et al. Hypoplasia of The Facial Artery Substituted By Transverse Facial Artery. *Trakia Journal of Sciences*. 2019; 17(Suppl. 2): 79–81, doi: 10.15547/tjs.2019.s.02.018-
18. Zhang S, Zhao S, Song P, et al. Application of a Modified Perforator Flap of the Transverse Facial Artery for Zygomaticofacial Skin Defect Repair After Tumor Resection. *Journal of Craniofacial Surgery*. 2023; 34(5): 1532–1535, doi: 10.1097/SCS.00000000000009230-

19. Żytkowski A, Tubbs RS, Iwanaga J, et al. Anatomical normality and variability: Historical perspective and methodological considerations. *Translational Research in Anatomy*. 2021; 23: 100105, doi: 10.1016/j.tria.2020.100105-

**Table 1.** Qualitative results of the data analysis. TFA – Transverse Facial Artery. STA – Superficial Temporal Artery. ECA – External Carotid Artery.

<b>Category</b>	<b>N</b>	<b>Percentage</b>
<i>Patients' sex</i>		
Females	42	47.7%
Males	46	52.3%
<i>Patients' side</i>		
Left	44	50.0%
Right	44	50.0%
<i>TFA Origin Artery</i>		
STA	84	95.5%
ECA	4	4.5%

**Table 2.** Overall results of the measurements. The results were established in millimeters (mm / mm<sup>2</sup>). LQ – lower quartile. UQ – upper quartile. SD – standard deviation. TFA – Transverse Facial Artery. STA – Superficial Temporal Artery. CCA – Common Carotid Artery. ECA – External Carotid Artery. MA – Maxillary Artery. PAA - Posterior Auricular Artery. OA – Occipital Artery. FA – Facial Artery.

Category	Media n	LQ	UQ	Minimu m	Maximu m	Mean	SD
TFA Length	43.39	38.53	46.37	30.50	64.26	43.43	6.46
TFA Diameter (At Its Origin)	2.26	1.93	2.54	1.14	3.32	2.25	0.47
TFA Cross-Sectional Area (At Its Origin)	2.54	1.67	3.10	0.63	5.97	2.50	1.04
TFA Branching Angle [deg]	74.55	59.09	89.70	30.40	149.38	75.24	21.80
STA Diameter at the TFA Origin	3.14	2.76	3.49	1.56	4.00	3.08	0.51
STA Cross-Sectional Area at the TFA Origin	5.33	4.24	6.15	1.47	10.19	5.30	1.66
CCA Diameter (Just Before Bifurcation)	8.02	7.37	8.82	4.62	11.78	8.11	1.31
CCA Cross-Sectional Area (Just Before Bifurcation)	41.44	35.12	49.46	13.80	77.26	42.47	12.24
ECA Diameter (Just Before Bifurcation into STA and MA)	4.10	3.70	4.53	2.48	5.56	4.11	0.67
ECA Cross-Sectional Area (Just Before Bifurcation STA and MA)	10.02	8.09	11.62	3.45	21.32	10.19	3.31
ECA Diameter (Just Before Its First Branch)	5.15	4.59	5.72	3.77	7.57	5.19	0.89
ECA Cross-Sectional Area (Just Before Its First Branch)	15.89	12.47	19.99	8.58	34.21	16.94	5.76
MA Diameter (At Its Origin)	3.38	3.09	3.63	1.76	4.50	3.32	0.51
MA Cross-Sectional Area (At Its Origin)	6.25	5.28	7.58	1.34	11.22	6.29	1.79
PAA Diameter (At Its Origin)	2.55	2.28	2.91	1.42	3.73	2.54	0.47
PAA Cross-Sectional Area (At Its Origin)	3.22	2.47	4.13	1.25	5.97	3.27	1.12
OA Diameter (At Its Origin)	2.91	2.49	3.17	1.51	5.08	2.85	0.61
OA Cross-Sectional Area (At Its Origin)	4.25	3.34	5.62	1.13	8.46	4.43	1.54
FA Diameter (At Its Origin)	3.24	2.91	3.63	1.99	5.00	3.26	0.58
FA Cross-Sectional Area (At Its Origin)	6.00	4.59	7.31	2.19	14.17	6.12	2.10

Distance between the origin of the MA to the origin of the TFA (over the surface of the ECA/STA)	6.36	5.11	8.04	3.08	18.02	6.82	2.35
Distance between the origin of the TFA to the origin of the STA (over the surface of the ECA/STA)	51.81	35.90	65.06	22.20	103.33	52.11	18.38
Distance between the origin of the OA to the origin of the TFA (over the surface of the ECA/STA)	46.48	39.33	52.54	20.48	81.59	46.74	10.69
Distance between the origin of the FA to the origin of the TFA (over the surface of the ECA/STA)	47.74	43.59	53.36	31.14	66.78	48.30	7.78
Distance between the origin of the PAA to the origin of the TFA (over the surface of the ECA/STA)	23.14	17.22	27.14	7.90	44.21	22.82	6.67
Distance between the origin of the TFA to the lower border of the Zygomatic Arch	24.16	21.94	27.38	12.13	36.07	24.63	4.10

**Table 3.** Results of the measurements with respect to the patient's sex. The results were established in millimeters (mm / mm<sup>2</sup>). LQ – lower quartile. UQ – upper quartile. SD – standard deviation. TFA – Transverse Facial Artery. STA – Superficial Temporal Artery. CCA – Common Carotid Artery. ECA – External Carotid Artery. MA – Maxillary Artery. PAA - Posterior Auricular Artery. OA – Occipital Artery. FA – Facial Artery.

Category	Sex	Median	LQ	UQ	Minimum	Maximum	Mean	SD	p-value
TFA Length	Female	41.80	38.32	44.09	30.50	49.98	41.12	4.66	0.01



	Males	4.33	3.76	4.73	2.74	5.56	4.31	0.66	
ECA Cross-Sectional Area (Just Before Bifurcation STA and MA)	Females	9.40	7.81	11.21	3.45	15.27	9.25	2.71	0.02
	Males	10.75	8.58	13.14	4.81	21.32	11.05	3.60	
ECA Diameter (Just Before Its First Branch)	Females	4.84	4.34	5.37	3.83	7.52	4.94	0.83	0.01
	Males	5.50	4.74	5.99	3.77	7.57	5.41	0.89	
ECA Cross-Sectional Area (Just Before Its First Branch)	Females	14.97	11.59	17.23	8.76	30.80	15.35	5.12	0.00
	Males	17.95	13.61	22.04	8.58	34.21	18.38	5.98	
MA Diameter (At Its Origin)	Females	3.23	2.79	3.57	1.76	4.19	3.19	0.56	0.04
	Males	3.51	3.15	3.65	2.49	4.50	3.45	0.42	
MA Cross-Sectional Area (At Its Origin)	Females	6.23	4.78	7.51	1.34	9.12	6.02	1.91	0.40
	Males	6.28	5.63	7.65	3.22	11.22	6.54	1.64	
PAA Diameter (At Its Origin)	Females	2.51	2.22	2.69	1.50	3.54	2.47	0.45	0.10
	Males	2.60	2.47	2.98	1.42	3.73	2.60	0.48	
PAA Cross-Sectional Area (At Its Origin)	Females	2.68	2.10	3.68	1.25	5.97	2.93	1.12	0.00
	Males	3.67	3.01	4.26	1.57	5.38	3.57	1.05	
OA Diameter (At Its Origin)	Females	2.65	2.34	2.96	1.51	3.85	2.59	0.52	0.00
	Males	3.04	2.69	3.43	2.03	5.08	3.07	0.60	
OA Cross-Sectional Area (At Its Origin)	Females	3.68	3.06	4.52	1.13	6.51	3.74	1.29	0.00




	Males	4.97	3.88	5.78	2.31	8.46	5.04	1.50	
FA Diameter (At Its Origin)	Females	3.04	2.78	3.37	1.99	3.82	3.01	0.47	0.00
	Males	3.55	3.08	3.88	2.29	5.00	3.49	0.58	
FA Cross-Sectional Area (At Its Origin)	Females	5.53	4.00	6.20	2.19	7.36	5.17	1.43	0.00
	Males	6.72	5.11	8.08	3.32	14.17	6.97	2.26	
Distance between the origin of the MA to the origin of the TFA (over the surface of the ECA/STA)	Females	6.16	4.98	7.95	3.57	11.92	6.62	2.06	0.53
	Males	6.80	5.12	8.16	3.08	18.02	7.00	2.61	
Distance between the origin of the TFA to the origin of the STA (over the surface of the ECA/STA)	Females	46.90	29.68	59.07	22.20	79.97	46.63	16.45	0.02
	Males	54.34	42.52	70.47	23.25	103.33	56.74	18.83	
Distance between the origin of the OA to the origin of the TFA (over the surface of the ECA/STA)	Females	44.32	38.54	47.67	20.48	68.55	43.85	8.89	0.00
	Males	50.02	44.50	55.69	28.19	81.59	49.26	11.56	
Distance between the origin of the FA to the origin of the TFA (over the surface of the ECA/STA)	Females	45.13	40.35	48.65	31.14	57.53	45.16	6.59	0.00
	Males	51.26	45.93	56.52	32.56	66.78	51.16	7.73	
Distance between the origin of the PAA to the origin of the TFA (over the surface of the ECA/STA)	Females	21.81	16.96	26.55	12.15	44.21	22.82	6.96	0.65
	Males	23.30	18.01	27.14	7.90	34.59	22.82	6.52	
Distance between the origin of the TFA to the lower border of the Zygomatic Arch	Females	23.98	21.85	25.58	17.89	32.99	23.82	3.18	0.10

	Males	25.10	22.03	28.49	12.13	36.07	25.38	4.70	
--	-------	-------	-------	-------	-------	-------	-------	------	--

**Figure 1.** Scheme, presenting the arterial anatomy of the region in which the measurements were taken. TFA – Transverse Facial Artery. STA – Superficial Temporal Artery. CCA – Common Carotid Artery. ECA – External Carotid Artery. ICA – Internal Carotid Artery. MA – Maxillary Artery. PAA - Posterior Auricular Artery. OA – Occipital Artery. FA – Facial Artery. LA – Lingual Artery.

**Figure 2.** Three-dimensional reconstruction of one of the analyzed computed tomography angiographies. Some of the arteries were removed in order to provide the best possible quality of the image. TFA – Transverse Facial Artery. STA – Superficial Temporal Artery. PAA - Posterior Auricular Artery. OA – Occipital Artery. FA – Facial Artery.



STA

TFA

MA

PAA

OA

FA

ECA

LA

ICA

CCA

