

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.



ISSN: 0015-5659

e-ISSN: 1644-3284

The double-headed accessory iliacus muscle: a case report

Authors: Marta Pośnik, Nicol Zielinska, Piotr Łabętowicz, Mariola Głowacka, Łukasz Olewnik

DOI: 10.5603/fm.98029

Article type: Case report

Submitted: 2023-10-29

Accepted: 2024-01-01

Published online: 2024-04-03

This article has been peer reviewed and published immediately upon acceptance. It is an open access article, which means that it can be downloaded, printed, and distributed freely, provided the work is properly cited.

Articles in "Folia Morphologica" are listed in PubMed.

CASE REPORT

The double-headed accessory iliacus muscle: a case report

Marta Pośnik et al., Accessory iliacus muscle

Marta Pośnik¹, Nicol Zielinska¹, Piotr Łabętowicz², Mariola Głowacka³, Łukasz Olewnik¹

¹Department of Anatomical Dissection and Donation, Medical University of Lodz, Lodz, Poland

²Department of Normal and Clinical Anatomy, Medical University of Lodz, Lodz, Poland

³Nursing Department, Masovian Academy in Płock, Poland

Address for correspondence: Łukasz Olewnik, Department of Anatomical Dissection and Donation, Medical University of Lodz, Łódź, Poland, email: lukasz.olewnik@umed.lodz.pl

ABSTRACT

Background: Numerous accessory muscles are present in the human body, many of which are clinically significant. We present a case of an anomalous accessory iliacus composed of two heads, whose occurrence and anatomical location indicate a high probability of causing femoral nerve compression.

Materials and methods: During a routine dissection of the posterior abdominal wall of a 78-year-old cadaver, a double headed accessory muscle was noted, measured and photographed.

Results: In addition to the normal anatomy of dissected structures from the posterior abdominal wall, an accessory iliacus muscle composed of superficial and deep heads was identified. In addition, the inferior roots of the divided femoral nerve located between the heads was found to follow an unusual course.

Conclusions: It is important to be aware of morphological variability around structures such as double-headed accessory iliacus muscle presented in this study, due to their association with neurovascular bundles and hip joint. The reported atypical morphology is not widely known in the literature but might be of great clinical significance, therefore knowledge of such variability might be regarded in order to diagnose properly and introduce accurate treatment.

Keywords: anatomical variations, accessory iliacus, femoral nerve compression

INTRODUCTION

Occurrence of morphological variations of anatomical structures such as muscles and their tendons was mentioned numerous in medically-oriented literature [14, 15, 18, 22, 31]. However, its diversity still remains surprising and hip muscles are no exception.

The iliacus (IM) is a triangular muscle of the inner hip with a vast origin encompassing two-thirds of the iliac fossa, the inner lip of the iliac crest, lateral aspect of the sacrum, anterior sacroiliac ligament and iliolumbar ligament. It inserts onto the lesser trochanter of the femur [17]. Together with the psoas major and occasionally, if present, the psoas minor, this muscle forms the iliopsoas muscle complex, which is recognised as the strongest flexor of the hip joint. Numerous variations of iliopsoas muscle complex have been noted. One of those is the occurrence of the accessory iliacus muscle (AIM). In previously presented cases, this structure was usually described as a slim muscular slip arising from the middle third of inner lip of iliac crest; it was distinguishable from iliacus muscle mass and traced the iliopsoas muscle complex distally onto its site of insertion: the lesser trochanter of the femur [6, 26].

Variations such as the AIM are of great clinical importance due to its close relation to the neurovascular structures. The lumbar plexus (LP) is usually located posterior to, or between, the layers of the psoas major. One component of the LP is the femoral nerve (FN), a structure formed by the posterior division of the ventral rami of L2, L3, L4, and occasionally by L1 and/or L5 roots; it leaves the muscle mass via its lateral margin and descends through the anterior surface of the IM [17]. Additional structures might produce tension in the FN, which would most likely result in neuropathy, presenting as pain that refers from the hip towards the knee joint.

The present report presents the first description of the unusual occurrence of a double-headed AIM together with an atypical course of the FN that most likely resulted in FN entrapment.

CASE REPORT

Unusual muscle was noticed during the routine dissection of the right posterior abdominal wall of a male cadaver that 78 years old at death. The dissection was performed according to standard techniques for research and teaching purposes, at the Department of Anatomical Dissection and Donation, Medical University of Lodz, Poland. Briefly, the cadaver was positioned in supine position on the dissection table. The intestines were separated from the greater omentum, mesentery and fatty tissue in order to expose the posterior abdominal wall structures. Iliopsoas muscle complex and components of lumbar plexus were identified. An additional muscular structure covered by common fascia together with the IM was noted.

The accessory muscle was composed of two heads characterised by a muscular origin: a superficial head that originated from middle third of inner lip of iliac crest, and a deep head that originated from both middle third of inner lip of the iliac crest and muscular mass of the IM. The superficial head descended obliquely and medially over the deep head. At the level of the inguinal ligament, both heads merged together and were completely distinguishable from the IM mass. Inferior to the inguinal ligament, an additional muscle is present which developed a tendon, which descended through the femoral triangle between the laterally-located IM and medially-positioned psoas major tendons, before finally inserting into the lesser trochanter. At the insertion point, the tendons of iliopsoas muscle complex were completely separated from one another via the abnormal muscle tendon.

The FN left the psoas muscle mass as two disconnected roots created from the posterior ventral ramus: superior root formed of L2 and L3, and an inferior one formed from L4. The superior root descended through the anterior surface of the superficial head of the accessory muscle, while the inferior root was located between its heads. Both roots joined together and formed a common main trunk of the FN at the lateral border of the superficial head, after the inferior root passed between the accessory muscle heads. No other deviations from the typical course of FN were noted — Figs. 1, 2.

Accessory muscle was subjected to detailed morphometric measurements and photographic documentation. The measurements were taken twice with up to 0.1 mm accuracy using an electronic caliper (Mitutoyo Corporation, Kawasaki-shi, Kanagawa, Japan). Morphometric measurements are presented in Table 1 and Table 2. The anomalous muscle was identified as a double-headed AIM.

Table 1. Morphometric measurements of individual parts of the AIM's superficial head

Length	110.10 mm
Width	
Proximal attachment	17.93 mm
Point of fusion with deep head	6.68 mm
Thickness	
Proximal attachment	1.38 mm
Point of fusion with deep head	0.70 mm

Table 2. Morphometric measurements of individual parts of AIM's deep head

Length	
Muscle belly	136.51 mm
Muscle belly to the fusion with superficial head	95.61 mm
Tendon	91.30 mm
Width	
Proximal attachment	46.68 mm
Musculotendinous junction	14.33 mm
Distal attachment	8.67 mm
Thickness	
Proximal attachment	3.47 mm
Musculotendinous junction	4.28 mm
Distal attachment	2.65 mm

DISCUSSION

Numerous morphological variations of the iliopsoas muscle complex components have been described in the literature. These were classified by Aleksandrova et al. based on their

relationship with the FN [1]. This classification comprises type A, where the muscular mass of the IM from the anterior and middle part was replaced by fibrous slips [1]; type B, with complete separation of IM and psoas major [8, 13, 26]; type C, with complete fusion of IM and psoas major with typical course of FN [9]; type D, with highly originated muscular slips that include extremely rare reports of psoas tertius and psoas quartus [5, 11, 27]; type E, with an iliacus minor - a muscle arising from the anterior inferior iliac spine of the ilium and inserting onto the anterior trochanteric line, and the ilio-capularis, with a similar origin - anterior trochanteric line and insertion onto the ilio-femoral ligament [2, 29]; type F, with deep and superficial layers of IM [10, 16]; type G, with large muscular slips of iliacus [21]; type H, with one slip, separated from IM by FN [6]; type I, with two muscular slips (accessory psoas and accessory iliacus) that pierce the FN [10] and J, with a small muscular slip of IM piercing through the FN [1]. Some authors introduce variations as iliacus minimus or iliacus minor in order to describe additional muscular bands of IM that originate from the iliolumbar ligament or ala ossis sacri [1, 23].

A different classification was introduced by Philippon et al. [20], who classified iliopsoas tendon unit into three types: 1 — single tendon generated by fusion of IM and psoas major (prevalence 28.3%), 2 — double banded tendon composed of IM and psoas major tendons separately (64.2%) and 3 triple-banded tendon composed of a medially-located psoas major tendon, an intermediate IM tendon and an AIM tendon in the most lateral position (7.5%) [20].

Case presented in this study, or more specifically its superficial head, seems to most closely resemble type H of the Aleksandrova et al. classification [1]; however, it cannot be classified as such, due to the existence of the second head. Our identified double-headed AIM also does not completely resemble type 3 from Philippon et al. [20], since it has a different arrangement of tendons: in our case, the IM tendon is positioned most laterally, psoas major most medially and they are separated by the AIM.

The critical zone for femoral nerve entrapment is believed to be the fibro-muscular ring bounded superficially by the inguinal ligament; however, regions such as the space around the iliopsoas is still regarded as entrapment-risk in various FN entrapment syndromes [12, 24, 28]. Additional muscular bands of both the IM and psoas major, especially those that split the FM, were previously suspected as causes of FN compression [1, 4, 6, 28]. Interestingly, FN might be split by more than one variant muscle. Parker et al. [19] presented a study where FN

was split into three parts by two additional muscles identified as iliacus minor and psoas quartus.

As accentuated by Bayrón-Ho et al. [4], when FN passes under the muscle slip and continues its usual route under the inguinal ligament and into the thigh, nerve compression and irritation are major concerns due to the variant muscle and route increasing the tension on the FN [4]. The possibility of clinically significant compression seems even more likely when a divided FN courses between muscular heads.

The FN has both motor and sensory functions. Its superior muscular branches innervate the pectineus, iliacus and sartorius muscles, which participate in hip flexion. The inferior muscular branches innervate the quadriceps components that participate in knee straightening and hip joint stabilization, i.e the rectus femoris and the lateralis, medialis and intermedius vasti muscles [17]. The FN also innervates the anterior and medial parts of the thigh via its cutaneous branches. Compression of the FN at such high point on its course, i.e. between the AIM heads, could impair the function of the muscles innervated by the nerve resulting in numbness, tingling or paraesthesia. It is possible that AIM constriction would intensify the compression, which may occur during iliopsoas muscle function, i.e. hip flexion. Nonetheless, to confirm such suspicions, further studies are required.

Wong et al. [30] presented a study where muscle that originated from the anteromedial surface of the quadratus lumborum and fused with the tendon of the psoas major, identified as psoas quartus, split the FN into multiple parts. Authors emphasized that variation of the FN and variant within the iliopsoas muscle complex may interfere with numerous surgical approaches to the region, including lateral transpsoas interbody fusion procedures, lumbar plexus blockade, femoral artery manipulation for angiography or other pelvic region procedures [30]. AIM occurring in accordance with the description presented in this study also may interfere with mentioned procedures. It is important to note, that due to the high variability of anatomical parameters it is impossible to develop standard recommendations for surgical procedures without prior imaging evaluation, therefore pre-operation visualization study seems compulsory in order to prevent from complications caused by occurrence of anatomical variation.

It is also possible that the AIM presented in this study could interfere in snapping hip syndrome. Two types of the snapping hip syndrome might be distinguished. The lateral snap, that corresponds to the sudden slip of the iliotibial band over the bulge of the greater

trochanter with an abnormal insertion of the gluteus maximus muscle when there is coxa vara, dysplasia or bi-iliac narrowness [25], and anterior snapping which occurs most frequently when the hip moves from a position of flexion-abduction-external rotation to the neutral position [7]. During this movement, the medial part of the iliacus is restricted between iliacus and iliopsoas tendons. Such syndrome manifests itself as a sudden, painful and audible snapping of the hip and is usually seen among young ballet dancers [7].

Tatu et al. [25], who examined anatomy of the femoral portion of the iliopsoas muscle, described the iliopsoas tendon as a musculotendinous complex made up of a main tendon arising from the psoas major, an accessory tendon arising from the IM and proper muscular fibers belonging to the IM. Interestingly, snapping hip syndrome is treated via iliopsoas tendon release. Surgical techniques of release confirms the necessity of deep anatomical knowledge of the musculotendinous complex of the femoral portion of the iliopsoas muscle complex, especially since it is suspected that recurrence is connected with incomplete release, i.e. when only one tendon is released during the procedure; this seems probable when two or three tendons are present [20]. Such variations as abnormal tendon formation, like that presented in the study, is likely to disturb such treatment, not only by the occurrence of an additional tendon, but also due to its atypical presentation, which does not completely fit into existing classifications. However, further studies are needed to confirm this.

Battaglia et al. [3] reported co-presentation of unilateral FN and bilateral sciatic nerve variants in one cadaver. In this study FN emerged both lateral and deep to the psoas major between the psoas major and IM and then was pierced and divided into two separate divisions by an accessory slip of the IM [3]. In described study authors emphasize that both the course of the sciatic nerve and FN as well as the surrounding musculature may affect the results of nerve traction tests performed commonly to ascertain the presence of a lumbar disc herniation [3]. Reproduction of radicular leg pain in both sciatic and FN distributions with nerve traction testing is a common sign of lumbar disc herniation. Variation of the sciatic nerve or FN split via additional muscular structure may produce a clinical picture similar to that of a lumbar or lumbosacral radiculopathy [3]. Described variations together with variant from this study should be considered when a suspected lumbar spine disc herniation is refractory to care since recognition of those variants may lead to earlier intervention of physiologic testing and better outcomes of treatment [3].

CONCLUSIONS

The study reports a case of accessory iliacus muscle composed of superficial and deep heads connected with a variable course of femoral nerve and a high possibility of nerve entrapment. In the authors opinion, morphological variation with particular definition is significant not only for anatomists, but also for clinicians, who should be aware of atypical structures that might interfere with numerous surgical procedures or neurovascular functions.

Article information and declarations

Availability of data and materials

Please contact authors for data requests (Łukasz Olewnik PhD — email address: lukasz.olewnik@umed.lodz.pl).

Author contributions

Marta Pośnik — project development, data collection and management, data analysis and manuscript writing.

Nicol Zielinska — data collection, data analysis and manuscript editing.

Piotr Łabętowicz — data analysis and manuscript editing.

Mariola Głowacka — data analysis and manuscript editing.

Łukasz Olewnik data analysis and manuscript editing.

All authors have read and approved the manuscript.

Ethics statement

The cadavers belonged to the Department of Anatomical Dissection and Donation, Medical University of Lodz.

Funding

The authors have no financial or personal relationship with any third party whose interests could be positively or negatively influenced by the article's content. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of interest: The authors declare that they have no competing interests.

REFERENCES

1. Aleksandrova JN, Malinova L, JeleV L (2013) Variations of the iliacus muscle: report of two cases and review of the literature. *Int J Anat Var* 6:149–152
2. Babst D, Steppacher SD, Ganz R, Siebenrock KA, Tannast M (2011) The iliocapsularis muscle: An important stabilizer in the dysplastic hip. *Clin Orthop Relat Res* 469:1728–1734. doi: 10.1007/s11999-010-1705-x
3. Battaglia PJ, Scali F, Enix DE (2012) Co-presentation of unilateral femoral and bilateral sciatic nerve variants in one cadaver: A case report with clinical implications. *Chiropr Man Therap* 20:34. doi: 10.1186/2045-709X-20-34
4. Bayrón-Ho DI, Tavez WV, Valentín-Carro N (2023) Atypical Origin of the Saphenous Nerve and a Variation of the Right Iliacus Muscle: A Case Report. *International journal of cadaveric studies and anatomical variations* 4:44–50
5. Clarkson RD, Rainy H (1889) Clarkson RD, Rainy H. 1889. Unusual arrangement of the psoas muscle. *J Anat Physiol* 504
6. D'Costa S, Lakshmi A, Ramanathan A, Madhyastha S, Nayak SR, Prabhu LV (2008) An accessory iliacus muscle: a case report. *Romanian Journal of Morphology and Embryology* 49:407–409
7. Deslandes M, Guillin R, Cardinal E, Hobden R, Bureau NJ (2008) The snapping iliopsoas tendon: new mechanisms using dynamic sonography. *AJR Am J Roentgenol* 190:576–581
8. Le Double AF (1897) Muscles de la fosse lombo-iliaque. In: *Traité des Variations du Système Musculaire de l'Homme*. Tome I, Paris,. Schleicher Frères 325–332
9. Fabrizio PA (2011) Anatomic variation of the iliacus and psoas major muscles. *Int J Anat Var* 4:28–30
10. JeleV L, Shivarov V, Surchev L (2005) Bilateral variations of the psoas major and the iliacus muscles and presence of an undescribed variant muscle - Accessory iliopsoas muscle. *Ann Anat* 187:281–286. doi: 10.1016/j.aanat.2004.10.006
11. Khalid S, Iwanaga J, Loukas M, Tubbs RS (2017) Split Femoral Nerve Due to Psoas Tertius Muscle: A Review with Other Cases of Variant Muscles Traversing the Femoral Nerve. *Cureus*. doi: 10.7759/cureus.1555

12. Kokubo R KKITMD (2016) The impact of tarsal tunnel syndrome on cold sensation in the pedal extremities. . *World Neurosurg* 92:249–254
13. Kopsch F (1914) *Rauber's Lehrbuch der Anatomie des Menschen. Abt. 3: Muskeln, Gefäße*, 10th Ed. Leipzig, Verlag von Georg Thieme 153–154
14. Koptas K, Zielinska N, Tubbs RS, Olewnik Ł, Łabętowicz P (2022) A newly reported muscle: an accessory infraspinatus or a deep layer of the latissimus dorsi? *Surgical and Radiologic Anatomy* 44:617–620. doi: 10.1007/s00276-022-02917-8
15. Kurtys K, Gonera B, Olewnik, Karauda P, Polguy M (2020) A highly complex variant of the plantaris tendon insertion and its potential clinical relevance. *Anat Sci Int* 95:553–558. doi: 10.1007/s12565-020-00540-4
16. Macalister A (1875) Additional observations on muscular anomalies in human anatomy (third series): With a catalogue of the principal muscular variations hitherto published. . *Transactions of the Royal Irish Academy* 126–200
17. Moore KL, Dalley AF, R. AAM, Moore KL (2014) *Moore clinically oriented anatomy*. Philadelphia, PA: Wolters Kluwer Health/Lippincott Williams & Wilkins
18. Olewnik Ł, Gonera B, Kurtys K, Tubbs RS, Polguy M (2021) “Popliteofascial muscle” or rare variant of the tensor fasciae suralis? *Folia Morphol (Warsz)* 80:1037–1042. doi: 10.5603/fm.a2020.0138
19. Parker A, Olewnik Ł, Iwanaga J, Dumont AS, Tubbs RS (2022) Iliacus minor and psoas quartus muscles traversing the femoral nerve. *Morphologie* 106:307–309. doi: 10.1016/j.morpho.2021.10.001
20. Philippon MJ, Devitt BM, Campbell KJ, Michalski MP, Espinoza C, Wijdicks CA, Laprade RF (2014) Anatomic variance of the iliopsoas tendon. *Am J Sports Med* 42:807–811. doi: 10.1177/0363546513518414
21. Rao TR, Kanyan PS, Rao S (2008) Bilateral variation of iliacus muscle and splitting of femoral nerve. *Neuroanatomy* 7:72–75
22. Ruzik K, Waśniewska A, Olewnik Ł, Tubbs RS, Karauda P, Polguy M (2020) Unusual case report of seven-headed quadriceps femoris muscle. *Surg Radiol Anat* 42:1225–1229. doi: 10.1007/s00276-020-02472-0
23. Spratt JD, Logan BM, Abrahams PH (1996) Variant Slips of Psoas and Iliacus Muscles, With Splitting of the Femoral Nerve. *Clinical Anatomy* 9:401–404
24. Stafford MA PPH DA (2007) Sciatica: a review of history, epidemiology, pathogenesis, and the role of epidural steroid injection in management. . *Br J Anaesth* 99:461–473
25. Tatu L, Parratte B, Vuillier F, Diop M, Monnier G (2002) Descriptive anatomy of the femoral portion of the iliopsoas muscle. *Anatomical basis of anterior snapping of the hip. Surgical and Radiologic Anatomy* 23:371–374. doi: 10.1007/s00276-001-0371-9
26. Tsuyoshi S, Nagahiro T (2016) Abdominal Wall Muscles. In *Bergman's Comprehensive Encyclopedia of Human Anatomic Variation* (eds R.S. Tubbs, M.M. Shoja and M. Loukas). doi: <https://doi.org/10.1002/9781118430309.ch36>

27. Tubbs RS, Oakes WJ, Salter EG (2006) The psoas quartus muscle. *Clin Anat* 19:678–680. doi: 10.1002/ca.20288
28. Unat F, Sirinturk S, Cagimni P, Pinar Y, Govsa F, Nteli Chatzioglou G (2019) Macroscopic observations of muscular bundles of accessory iliopsoas muscle as the cause of femoral nerve compression. *J Orthop* 16:64–68. doi: 10.1016/j.jor.2018.12.009
29. Ward WT, Isabelle-De'sire'e Fleisch, Ganz R (2000) Anatomy of the Iliocapsularis Muscle Relevance to Surgery of the Hip. *Clin Orthop Relat Res* 374:278–285
30. Wong TL, Kikuta S, Iwanaga J, Tubbs RS (2019) A multiply split femoral nerve and psoas quartus muscle. *Anat Cell Biol* 52:208–210. doi: 10.5115/acb.2019.52.2.208
31. Zielinska N, Tubbs RS, Ruzik K, Olewnik Ł (2021) Classifications of the extensor hallucis longus tendon variations: Updated and comprehensive narrative review. *Ann Anat* 238. doi: 10.1016/j.aanat.2021.151762




Figure 1. Structures of the right posterior abdominal wall after dissection. PM psoas major muscle, IM iliacus muscle, sAIM superficial head of the accessory iliopsoas muscle, dAIM deep head of the accessory iliopsoas muscle, L2L3 superior root of the femoral nerve, L4 inferior root of the femoral nerve




Figure 2. Structures of the right posterior abdominal with visible course of the femoral nerve. PM psoas major muscle, IM iliacus muscle, sAIM superficial head of the accessory iliacus muscle, dAIM deep head of the accessory iliacus muscle, LFC lateral femoral cutaneous nerve, GF branches of the genitofemoral nerve, L2L3 superior root of the femoral nerve, L4 inferior root of the femoral nerve