

The venous system of the human foetal spinal cord. Scanning electron microscope of vascular corrosion casts

J. Zawiliński¹, K. Zagórska-Świeży¹, J. Składzień²

¹SEM Laboratory of the ENT Clinic, University Hospital in Krakow, Poland

²Department of Otorhinolaryngology, Jagiellonian University, Krakow, Poland

[Received 11 October 2013; Accepted 25 November 2013]

The investigation was carried out on 16 human foetal cadavers at the age of 17–23 weeks from the time of conception. The foetal vascular system was injected with the synthetic resin MERCOX CL-2R and analysed in scanning electron microscope. The vascular system of the foetal spinal cord was studied. The foetal vascular system was characterised by high variability concerning the number, course and localisation of blood vessels. It contained numerous anastomoses with the internal spinal venous plexuses, which included anterior and posterior radicular veins. Large arteries running on the surface of the spinal cord are accompanied by the homoname veins. The venous system of the investigated fetuses was divided into 2 categories of veins: internal veins responsible for the drainage of blood from the central area, that is central and peripheral veins coming radially to the surface of the spinal cord and external veins, which form the venous system of the surface of the spinal cord. The venous system of the foetal spinal cord was also examined as to the presence of the valves. (Folia Morphol 2014; 73, 2: 139–142)

Key words: veins, venous valves, foetal spinal cord

INTRODUCTION

The venous system of the human foetal spinal cord has not been paid much attention to. In available literature few papers have been found on this subject [1, 2, 4, 5, 12, 13] and scarce works in the form of albums or illustrations [14].

Although many authors investigating the arterial system of the spinal cord touched the subject of the veins, they were treated in the marginal way. The information was usually restricted to the statements that the veins run together with the arteries. None of the above-mentioned papers has been concerned with the venous system of the human foetal spinal cord.

Recently there have been several publications dealing with this problem, but their authors focused

mainly on the venous system of the animal spinal cord [8].

In such circumstances, present investigations due to the use of scanning microscope enabling almost unfaillingly to determine the type of blood vessel, cast more light on the problem of the veins of the human foetal spinal cord.

The aim of this work was the observation of the venous system of the human spinal cord during the foetal development, using the technique of vascular corrosion casts and scanning electron microscopy [9, 10], which gives high resolution pictures of the vascular network. Only few publications have been concerned with the arterial system of the spinal cord during the prenatal development. For that reason the authors of this paper have undertaken the investigation.

MATERIALS AND METHODS

The investigation of the vascular system of grey and white matter of the spinal cord was carried out on 16 human foetal cadavers ranging in crown-rump length 150–235 mm, which was about 17–23 weeks from the conception. After the foetal vascular system was rinsed with heparinised physiological saline and fixed with fixing agent (0.66% paraformaldehyde/0.08% glutaraldehyde), it was filled through the ascending aorta with the synthetic resin Mercox CL-2r/2B with an additive of methyl metacrylate monomer. After polymerisation of the resin at 55°C, the spinal columns with spinal cords and meninges were isolated. Then demineralisation of the soft tissues was carried out in 5% trichloro-acetic acid followed by maceration in 10–15% KOH alternating with distilled water. The obtained specimens were then rinsed thoroughly in distilled water, freeze-dried and coated with gold, and finally analysed under the scanning electron microscope JEOL JSM 35 CF.

RESULTS AND DISCUSSION

The venous system of the foetal spinal cord exhibits great variability as to the number, course and localisation of the vessels. It contains numerous anastomoses with internal spinal venous plexuses, which include anterior and posterior radicular veins. Large arteries running on the surface of the spinal cord are accompanied by homonymous veins. In case of large vascular trunks in the foetus, veins had usually larger diameter than the arteries (Figs. 1, 2). Similarly to an adult, the veins branched irregularly, joined with various anastomoses or formed large singular vascular trunks connected to the radicular veins (Figs. 3, 4). The veins of the spinal cord do not contain the valves, what is confirmed by the increase in the pressure of the cerebro-spinal fluid in the Valsalva's or Stookey's trials. However, some authors have described the presence of the venous valves, or rather a specific mechanism of valves including the formation of the internal fold in the site of anastomosis of the vessels [6]. These folds can occasionally be the cause of poor filling of the spinal cord veins in case of their injecting with filling mass. In the investigated foetal material the valves were not observed, however, in some foetuses difficulties were noted in the perfusion of the injected mass into the lumen of the vessels even at low density of the injected mass as in case of synthetic resins.

In the venous system of the investigated foetuses 2 categories of veins were distinguished:

- internal veins (venae spinales internae) responsible for the drainage of the blood from the central




Figure 1. Anterior spinal vessels; ASA — anterior spinal artery; VSA — anterior spinal vein; VL — lateral vein. Note the differences in the imprints of the endothelial nuclei of arteries and veins. 20 week's crown-rump length. Bar 100 μ m.



Figure 2. Anterior aspect of the spinal cord; ARA — anterior radicular artery; VRA — anterior radicular vein; VL — lateral vein; ASA — anterior spinal artery; VSA — anterior spinal vein. 23 week's crown-rump length. Bar 1000 μ m.




Figure 3. Posterior aspect of the spinal cord; ASPDx — right posterior spinal artery; TP — posterior trunk; VL — lateral vein. 21 week's crown-rump length. Bar 100 μ m.



Figure 4. Posterior aspect of the spinal cord; TP — posterior trunk; VRP — posterior radicular vein; VL — lateral vein. 22 week's crown-rump length. Bar 1000 μm .

area (grey matter of the anterior and lateral horns, intermediary white matter, basis of the posterior horns and deep regions of white matter), which include central veins and peripheral veins coming radially to the surface of the spinal cord (responsible for the drainage of the blood from the posterior horns, dorsal area of white matter of the posterior and lateral columns) (Figs. 5, 6); — external veins (venae spinales externae) located on the surface of the spinal cord, responsible for the drainage of the blood from the system of internal veins and the blood outflow to the internal spinal venous plexuses. Their layout accompanies the arterial trunks, hence their parallel course to the longitudinal axis of the spinal cord. They join the system of anterior and posterior radicular veins.

The anterior spinal cord vein ran accompanying laterally or ventrally the homoname artery. This is consistent with the observations on the adults [1–7, 11–13]. In the foetus, its diameter always exceeded by about 20–120 μm the diameter of the artery. It received the lateral and medial blood supply and then via the anastomoses with anterior radicular veins the blood was carried to the internal spinal venous plexuses.

The posterior veins of the spinal cord were frequently fused together into a single trunk located on the posterior surface of the spinal cord or formed venous plexuses rich in anastomoses, responsible for drainage of the blood from the area of the spinal cord supplied by the peripheral arterial system. This single trunk in the stage of formation (in the investigated material) had the greatest diameter in the lumbar region of the spinal cord, smaller in the cervical region and the smallest in the thoracic region (Fig. 7). Karmański et al. [8] in the studies on animal samples reported the greatest diameter in the sacral region.



Figure 5. Lateral aspect of the spinal cord; VSA — anterior spinal vein; VC — central veins. 20 week's crown-rump length. Bar 1000 μm .




Figure 6. Anterior aspect of the spinal cord; VC — central vein; Vv — venules and capillaries. 17 week's crown-rump length. Bar 100 μm .




Figure 7. Posterior aspect of the spinal cord; TP — posterior trunk; VR — radial veins; VL — lateral vein. 27 week's crown-rump length. Bar 100 μm .

They described the presence of a single venous trunk on the ventral side of the spinal cord similar to that on the dorsal side. In the analysed material, no such trunk

was found on the ventral side. The observed discrepancies might result from the differences between the species or from the assessment of the venous system at various stages of its development. The blood was carried from the external veins of the foetal spinal cord to the internal venous plexuses of the spinal cord. This plexus joining the outflow of the blood from the spinal cord and walls of the spinal canal exhibits better development on the ventral side in the adult. On the dorsal side, this plexus shows considerable variability as to the shape and levels of arrangement of the anastomoses [6]. The blood flow to the internal spinal cord plexuses is via the anterior and posterior radicular veins of various number and localisation. From the anterior plexus the blood flows to the baso-vertebral veins and partially, similarly as from the dorsal plexus, the blood outflows via superior and inferior intervertebral veins, situated in the area of intervertebral openings, to the external spinal venous plexuses. On both sides of the anterior surface of the dura mater there are longitudinal trunks, the so-called medial epidural vein and lateral epidural vein whose numerous anastomoses form anterior internal spinal plexus [14]. Next the outflow of the blood occurs through the system of azygos veins, in the sacral part to the internal iliac veins and their branches and in the cervical part to the spinal veins, internal cervical veins and the veins accompanying the arteries branching off the external cervical arteries. In the analysed samples, the internal spinal plexuses were well formed in all age groups. However, the epidural veins of longitudinal course, which were described in the adult, were not found in the investigated material.

CONCLUSIONS

1. Large arteries running on the surface of the spinal cord are accompanied by homoname veins. The diameter of the veins was usually greater by about 40 μm than that of the arteries and the veins located on the posterior surface of the spinal cord had greater diameter in all cases.
2. The venous system of the spinal cord in the investigated fetuses was divided into 2 categories of veins:
 - external veins, which form the venous system of the spinal cord surface;
 - internal veins, including the veins responsible for the drainage of the blood from the central area, that is the central and peripheral veins coming radially to the surface of the spinal cord.
3. In the investigated material no venous valves were found, although in case of some fetuses the difficulties in perfusion of the mass into the lumen of the veins were observed, which was probably related to the presence of the endothelial folds in the site of the vessel division.
4. The posterior spinal veins occasionally fused together forming a single trunk situated on the posterior surface of the spinal cord or formed characteristic venous plexuses on the surface of the spinal cord. On the anterior surface, no anterior trunk running independent of the artery, described in animals, was noted. On the whole length of the spinal cord, the anterior spinal artery was accompanied by 1 anterior spinal vein, even in case of 2 anterior spinal arteries.
5. In the investigated material no epidural veins of longitudinal course, described in the adult, were found.

REFERENCES

1. Clemens HJ (1961) Die Venensystem der Menschlichen Wirbelsäule. Morphologie und Funktionelle Bedeutung. Walter de Gruyter & Co. Berlin.
2. Clemens HJ (1961) Beitrag zur Histologie der Plexus venosi vertebrales interni. *Z Mokrosk Anat Forsch*, 67: 183–189.
3. Corliss CE (1984) Embryology of blood circulation. Blood vessels and organ systems. Academic Press Inc, Orlando, San Diego, San Francisco, New York, London, Toronto, Montreal, Sydney, Tokyo, Sao Paulo.
4. Craigie EH (1983) Vascular supply of the spinal cord. Igaku-Shoin Ltd, Tokyo.
5. Gillilan LA (1970) Veins of the spinal cord. Anatomic details; suggested clinical applications. *Neurology*, 20: 860–868.
6. Groen RJM, Groenewegen HJ, van Alphen HAM, Hoogland PVJM (1997) The morphology of the human internal vertebral venous plexus. A cadaver study after intravenous Araldite CY 221 injection. 11-th International Congress of Neurological Surgery, Amsterdam.
7. Kadyi H (1888) O naczyniach krwionośnych rdzenia pacierzowego ludzkiego. Drukarnia UJ, Krakow.
8. Karmański A, Mikusek J, Karmańska W, Łopata P (1995) External veins of the spinal cord in the wistar Rats. *Folia Morphol*, 54: 231–241.
9. Lametschwandtner A, Miodoński AJ, Simonsberger P (1980) On the prevention of specimen charging in scanning electron microscopy of vascular corrosion casts by attaching conductive bridges. *Microscopie*, 36: 270–273.
10. Lametschwandtner A, Lametschwandtner U, Weiger T (1990) Scanning electron microscopy of vascular corrosion casts: technique and application: updated review. *Scanning Microsc*, 4: 889–941.
11. Lazorthes G, Gouaze A, Djindjian R (1973) Vascularisation et circulation de la moelle epiniere. Anatomie, physiologie, pathologie, angiographie. Masson et Cie, Paris.
12. Renard M, Masson J-P, Larde D (1976) Les veines épidurales lombaires; Les veines du canal Sacré. *Bull Assoc Anat*, 60: 787–803.
13. Tadie M, Hemet J, Aaron C, Bianco C, Creissard P, Houdard P (1979) Le dispositif protecteur anti-reflux des veines de la moelle. *Neurochirurgie*, 25: 28–30.
14. Théron J, Moret J (1978) Spinal phlebography. Springer-Verlag, Berlin.