

# Protective effect of resveratrol on acrylamide-induced renal impairment

M.M. Nasralla<sup>1</sup>, S.M. Zaki<sup>1,2</sup>, R.A. Attia<sup>1</sup>

<sup>1</sup>Faculty of Medicine, Cairo University, Cairo, Egypt

<sup>2</sup>Fakeeh College for Medical Sciences, Jeddah, Saudi Arabia

<sup>3</sup>Department of Pharmacology, Fakeeh College for Medical Sciences, Jeddah, Saudi Arabia

[Received: 8 October 2020; Accepted: 19 October 2020; Early publication date: 30 October 2020]

**Background:** Acrylamide (ACR) has a wide range of uses. It possesses a renal impairment effect. The work aimed to study the possible protecting role of resveratrol (RVS) over the ACR-mediated renal impairment in rats. The suggested underlying mechanisms participating in such protection were investigated.

**Materials and methods:** Thirty Sprague-Dawley adult albino rats were divided into three groups: control, ACR, and RVS. After 4 weeks, the kidney was removed and prepared for histological, immunohistochemical, and biochemical studies. The activity of tissue oxidative (malondialdehyde [MDA]) and anti-oxidative (glutathione [GSH]) markers were assessed.

**Results:** Acrylamide induced glomerular renal affection in the form of shrinkage and distortion of the glomeruli with wrinkling of their basement membranes and widening of the urinary spaces. Degenerative tubular changes were markedly present in the proximal convoluted tubules. The necrotic tubular cells exhibited cytoplasmic vacuolation with desquamated epithelial cells within the tubular lumen. ACR increases the deposition of collagen fibres in the basement membrane of the glomerular capillaries and induced thickening of the basement membranes of the renal corpuscles and renal tubules. The administration of RVS affords high protection to the kidney. The glomeruli and renal tubules were nearly normal. The content of collagen fibres and the periodic acid Schiff reaction of the basement membrane of the renal tubules were 70% and 19% lower linked to the ACR group. The creatinine and urea levels decreased by 51% and 47%. RVS induced such a protective role through its antioxidant effect as the MDA level decreased by 45%, while the GSH level increased by 83% compared with the ACR group.

**Conclusions:** Acrylamide causes structural and functional disorders of the kidney. It induces kidney damage through oxidative stress and apoptosis. With the use of RVS, normal kidney architecture was preserved with little structural changes. Adding, functional kidney test became normal. RVS exerts its protective effect through its anti-apoptotic and antioxidant features. (Folia Morphol 2021; 80, 4: 985–993)

**Key words:** resveratrol, acrylamide, kidney

Address for correspondence: Dr. S.M. Zaki, Fakeeh College for Medical Sciences, Jeddah, Saudi Arabia, e-mail: zaky.sherif@yahoo.com

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

## INTRODUCTION

Acrylamide (ACR) is a well-known environmental pollutant that exerts a range of systemic toxic effects on people after both occupational and dietary exposure [2, 22]. It possesses a variety of harmful properties: carcinogenicity, genotoxicity, neurotoxicity, and reproductive toxicity [5, 7]. ACR and its analogues are widely used in various chemical and environmental applications and are produced by heating plant tissue-derived biological material [25]. ACR formation occurs during food processing due to exposure of carbohydrates to temperatures above 200°C [12]. High levels of ACR have been found in foodstuffs commonly consumed, in particular; potato chips and bread [31].

Acrylamide is absorbed from the gastrointestinal tract and dispersed widely in body fluids and stored in the liver and kidney [28]. The ACR is known to cause structural and functional changes in many organs. The renal tubular cells undergo degenerative vacuolar changes, inflammatory cell infiltration and periglomerular oedema [28]. Furthermore, ACR administration in rats raises the levels of serum urea, creatinine, uric acid, and renal proinflammatory cytokine [1].

The metabolism of ACR triggers the release of free radicals (ROS), which initiates oxidative stress leading to an imbalance in the development and degradation of ROS [19]. It also induces lipid peroxidation and DNA harm [1].

The effects of many antioxidants and anti-inflammatory compounds such as olive oil, vitamin E and 5-aminosalicylic acid were studied to prevent and treat ACR-induced renal impairment [9, 19].

Resveratrol (RVS) is a phytoalexin found in at least 72 species of plants, many of which are eaten by humans, including mulberries, peanuts, and grapes [8]. RVS has anti-inflammatory, antiplatelet, antioxidant, and anti-carcinogenic activity, as well as the ability to reduce kidney damage caused by chemical compounds [8, 10]. However, it is unclear whether RVS can defend against ACR-induced renal impairment or not. Therefore, we studied the oxidative and apoptotic damaging effect of the ACR over the kidney and investigated the protecting role of RVS over such renal impairment in rats.

## MATERIALS AND METHODS

### Animals

Thirty Sprague-Dawley adult albino rats weighing 170–200 g were encompassed in our study. The an-

imals were maintained in spacious wire mesh cages in a special room with direct daylight and natural ventilation. The rats had free access to standard rat chow and water.

All the animals were treated according to the standard guidelines for the care and use of laboratory animals. The study was permitted by the Ethics Committee, Faculty of Medicine, Cairo University, Egypt. The procedures followed were following the ethical standards of the responsible organization and with the Helsinki Declaration of 1975 as revised in 1983.

### Experimental design

The rats were distributed into three groups (10 in each group): control (given distilled water at a dose of 1 mL), ACR group, and RVS group (concomitant ACR + RVS).

### Chemicals

Acrylamide was obtained in a container of powder purchased by Biostain Company (United Kingdom) weighing 500 g. It was dissolved in distilled water, in a concentration of 10 g/L. It was given at a dose of 1 mL of distilled water containing 40 mg/kg/day orally via gastric gavage [9].

Resveratrol (purity, > 99%) was purchased from Sigma-Aldrich (St. Louis, MO, USA). It was dissolved in dimethyl sulfoxide and diluted in 0.9% physiological saline. It was given at a daily dose of 20 mg/kg/day orally via gastric gavage [18].

By the end of the experiment (after 4 weeks), each animal was weighed, and a blood sample was withdrawn from the tail vein using a fine heparinized capillary tube. The kidney was extracted, washed with saline, and left to dry on a plot paper.

### Light microscopic study

Kidney specimens were fixed in formalin 10%, dehydrated in ethyl alcohol, cleared in xylol, and embedded in paraffin wax. Sections of 5 µm thickness were cut and mounted on glass slides. Other sections were mounted on positively charged slides for immunohistochemistry. The sections were subjected to the following:

- haematoxylin and eosin (H&E) and Masson's trichrome stained sections were prepared according to Suvana et al. [24];
- histochemical evaluation: periodic acid Schiff (PAS) stain: PAS stained sections were prepared according to Suvana et al. [24];

— immunohistochemistry analysis of Bcl-2-associated X protein (BAX) [20].

Paraffin sections were prepared. Then, a suitable quantity of serum was added to the sections for 30 min. Endogenous peroxidase was inactivated with a methanol solution containing  $H_2O_2$  (1:50) for 10 min and washed with phosphate buffered saline (PBS). The tissue sections were blocked with 1.5% serum for 30 min. The sections were incubated with the primary antibody BAX (anti-human BAX protein, DakoCytomation, Denmark), followed by the secondary antibody (biotinylated link universal from the commercial kit LSAB: DakoCytomation, Denmark). Subsequently, samples were incubated with AB enzymes for 30 min and rinsed in PBS. Positive signals were detected using peroxidase chromogenic substrates. The negative control included PBS instead of the secondary antibody.

#### Image analysis and morphometric measurements

Using the Leica LAS V3.8 image analyser computer system (Switzerland), the following parameters were assessed: the diameters of the renal glomeruli and proximal convoluted tubules, the width of the renal space, and the height of the lining proximal tubular epithelium. Adding, the number of structurally altered glomeruli was assessed as a percentage of the total number of glomeruli. The content of the collagen fibres was also assessed.

In PAS-stained histological sections, the optical density of the basement membrane of the proximal convoluted tubules and the parietal layers of Bowman's capsules was additionally determined.

In the BCL2 immunohistochemical staining area percentage of the immune reaction was also measured.

#### Biochemical study

Blood samples withdrawn from the rats before sacrifice were used for biochemical assessment in the Biochemistry and Molecular Biology Department, Faculty of Medicine, Cairo University, Egypt.

The serum urea and creatinine levels were estimated by the conventional colorimetric method using Quanti Chrom TM Assay Kits (DIUR-500 and DICT-500), based on the improved Jung and Jaffe methods, respectively [32]. The mean values of these biochemical parameters were calculated and subjected to statistical analysis.

**Tissue level of malondialdehyde (MDA) and reduced glutathione (GSH).** The renal tissue was

homogenised in 5–10 mL cold buffer (50 mM potassium phosphate, pH 7.5. 1 mM EDTA) per gram tissue then it was centrifuged at 100,000 g for 15 min at 4°C. The supernatant was removed for assay and stored on ice.

Malondialdehyde assay was performed with thiobarbituric acid (TBA) test in the supernatant, according to the method suggested by Buege and Aust [4]. MDA reacts with TBA to give a red compound absorbing at 535 nm.

Measurement of GSH was based on the reduction of 5,5-dithiobis (2-nitrobenzoic acid) (DTNB) with reduced GSH to produce a yellow compound [6].

#### Statistical analysis

The mean values of relative kidney weight, histomorphometric measurements, and biochemical levels were analysed using SPSS version 22. Statistical estimation was done using ANOVA followed by Bonferroni pairwise comparisons.

## RESULTS

#### Light microscopic evaluation


The examination of the control group showed intact architecture of the renal cortex. The renal cortex is formed of renal corpuscles, and proximal and distal convoluted tubules (Fig. 1A).

The examination of the renal cortex of the ACR group revealed marked structural changes. The renal glomeruli showed moderate to marked shrinkage, distortion, segmentation, and vacuolation with widened urinary spaces. The renal tubules were dilated with marked diminution of their epithelial height and widening of their lumina. Their lining cells showed cytoplasmic vacuolation, cellular fragmentation, and intraluminal cast formation were observed in many convoluted tubules. The interstitium showed congested blood vessels with intimal thickening and massive cellular infiltration could be also seen (Fig. 1B–D).

Examination of the RVS group revealed the almost normal appearance of most of the glomeruli and the tubules. Minimal interstitial inflammatory cellular infiltration was encountered (Fig. 1E).

#### Content of collagen fibres

The content of the fibres was minimal in the control group. The content increased around the parietal layers of Bowman's capsules and the basement membranes of the renal tubules in ACR (9-fold) and RVS



**Figure 1.** A. Normal renal architecture in the control group showing normal architecture with normal renal glomeruli (G), Bowman's spaces (S), proximal convoluted tubules (PCT) (P), and distal convoluted tubules (DCT) (D); B–D. Shrinkage and distortion of renal glomeruli (G) with glomerular congestion and interstitial cellular infiltration (arrow) in the acrylamide group. Note vacuolation and degenerated nuclei of the PCT (P); E. Normal appearance of renal glomeruli (G), Bowman's spaces (S), PCT (P), and DCT of the resveratrol group. Note minimal interstitial cellular infiltration (arrow) (haematoxylin and eosin;  $\times 400$ ); d — distal convoluted tubules; BC — blood capillary.

(2-fold) groups when related to the control group. The content in the RVS group was 70% lower when related to the ACR group (Fig. 2A–C, Table 1).

### Histochemistry of the kidney

In the control group, the basement membranes of the renal tubules and the parietal layers of Bowman's capsules demonstrated a weak PAS reaction. Adding, the apical brush borders of the proximal convoluted tubules (PCT) were intact and partially occluding the tubular lumina (Fig. 3A).


The basement membranes of the renal tubules and the parietal layers of Bowman's capsules of the ACR group showed an intense PAS reaction (42% higher than the control group). Adding, the apical brush borders of the PCT were attenuated (Fig. 3B). With the use of RVS, the PAS reaction became comparable to the control group and was 19% lower than the ACR group (Fig. 3C, Table 1).

### Immunohistochemical staining BAX

BAX showed a weak reaction in the control group (Fig. 4A). The area percentage of BAX immunopositive cells increased 4.5-fold in the ACR group matched to the control group (Fig. 4B, C, Table 1). With the use of RVS, the area percentage of the immunopositive cells decreased 56% equated to the ACR group; however, the area percentage in the RVS group was 1.4-fold higher than the control group (Fig. 4D, Table 1).

### Biochemical and oxidative/antioxidative markers

The serum creatinine and urea levels increased in the ACR group by 2.75-fold, 1.9-fold linked to the control group. With the concomitant use of RVS, the levels of creatinine and urea decreased by 51%, 47% allied to the ACR. Nevertheless, the levels of creatinine and urea in the RVS group were 83%, 55% higher than the control group (Table 1).



**Figure 2.** **A.** The control group with normal basement membranes of renal tubules (arrows) and parietal layers of Bowman's capsules (arrow); **B.** Acrylamide group with thickened basement membranes of the renal tubules (arrows) and parietal layer of Bowman's capsules (arrowheads). Note augmented interstitial collagen fibres (arrow); **C.** Resveratrol group with normal appearance of renal tubules (arrows) and parietal layer of Bowman's capsules (arrow). Note a slight increase of interstitial collagen fibres (Masson's trichrome;  $\times 400$ ).

**Table 1.** Content of collagen fibres, periodic acid Schiff (PAS) reaction, apoptotic, biochemical and oxidative/antioxidative markers


Group		Content of collagen fibres	Optical density of PAS reaction	Area percentage of BAX immunopositive cells	Serum creatinine [mg/dL]	Serum urea [mg/dL]	Malondialdehyde [nmol/g protein]	Glutathione [ $\mu$ mol/g protein]
Control	Mean $\pm$ SD	3.51 $\pm$ 0.46	0.47 $\pm$ 0.01	1.35 $\pm$ 0.15	0.12 $\pm$ 0.01	36.1 $\pm$ 2.0	17.52 $\pm$ 2.4	0.28 $\pm$ 0.02
Acrylamide group	Mean $\pm$ SD	30.46 $\pm$ 0.50	0.67 $\pm$ 0.01	7.45 $\pm$ 0.37	0.45 $\pm$ 0.03	106.6 $\pm$ 2.2	42.18 $\pm$ 3.6	0.12 $\pm$ 0.02
	Versus control group	< 0.001*	< 0.001*	< 0.001*	< 0.001*	< 0.001*	< 0.001*	< 0.001*
	Versus resveratrol group	0.003*	0.005*	< 0.001*	< 0.001*	< 0.001*	< 0.001*	< 0.001*
Resveratrol group	Mean $\pm$ SD	9.63 $\pm$ 0.60	0.54 $\pm$ 0.01	3.24 $\pm$ 0.21	0.22 $\pm$ 0.03	56.6 $\pm$ 2.4	23.16 $\pm$ 1.8	0.22 $\pm$ 0.01
	Versus control group	0.002*	0.144	0.052	0.007*	< 0.001*	< 0.001*	0.259
	Versus acrylamide group	0.003*	0.005*	< 0.001*	< 0.001*	< 0.001*	< 0.001*	< 0.001*

\*p-value significant; SD — standard deviation; BAX — Bcl-2-associated X protein


The values of the oxidative marker (MDA) in the ACR group increased by 1.4-fold, while the anti-oxidative marker (GSH) decreased by 1.3-fold compared with the control group. With the use of RVS, the MDA level decreased by 45%, while the GSH level increased by 83% compared with the ACR group. Still, the level of both markers was away from the control group (Table 1).

#### Morphometric glomerular and PCT changes

The percentage of the affected glomeruli in the ACR group increased 8.8-fold matched to the control group. With the use of RVS, the per cent of the affected glomeruli decreased 71% equated to the ACR group; however, the percentage in the RVS group was 1.6-fold higher than the control group (Table 2).



**Figure 3.** **A.** Apical brush borders (arrowheads) of proximal convoluted tubules (PCT) of the control group occluding the tubular lumina. Note weak periodic acid Schiff (PAS) reaction of parietal layers of Bowman’s capsules and basement membranes of PCT; **B.** Intense PAS reaction of parietal layers of Bowman’s capsules and the basement membranes of the renal tubules (arrows) of the acrylamide group; **C.** Moderate PAS reaction of the resveratrol group (PAS;  $\times 400$ ).



**Figure 4.** **A.** A negative immunoreactivity of the control group; **B.** A positive immunoreactivity of the acrylamide group; **D.** A negative immunoreactivity of the resveratrol group (BAX;  $\times 400$ ).

**Table 2.** Morphometric glomerular and proximal convoluted tubules (PCT) changes

Group		Percentage of the affected glomeruli	Glomerular diameter [ $\mu\text{m}$ ]	Width of urinary space [ $\mu\text{m}$ ]	PCT diameter [ $\mu\text{m}$ ]	PCT epithelial height [ $\mu\text{m}$ ]
Control	Mean $\pm$ SD	<b>5.23 <math>\pm</math> 1.78</b>	<b>353.29 <math>\pm</math> 1.39</b>	<b>24.93 <math>\pm</math> 1.6</b>	<b>180.38 <math>\pm</math> 1.24</b>	<b>134.59 <math>\pm</math> 2.21</b>
Acrylamine group	Mean $\pm$ SD	<b>49.15 <math>\pm</math> 2.7</b>	<b>146.45 <math>\pm</math> 2.09</b>	<b>83.40 <math>\pm</math> 2.05</b>	<b>237.13 <math>\pm</math> 1.64</b>	<b>50.80 <math>\pm</math> 2.77</b>
	Versus control group	< 0.001*	< 0.001*	< 0.001*	< 0.001*	< 0.001*
	Versus resveratrol group	< 0.001*	< 0.001*	< 0.001*	< 0.001*	< 0.001*
Resveratrol group	Mean $\pm$ SD	<b>13.2 <math>\pm</math> 2.2</b>	<b>330.74 <math>\pm</math> 1.12</b>	<b>35.69 <math>\pm</math> 1.38</b>	<b>193.46 <math>\pm</math> 1.35</b>	<b>120.64 <math>\pm</math> 3.01</b>
	Versus control group	0.003*	0.01*	0.01*	< 0.001*	< 0.001*
	Versus acrylamide group	< 0.001*	< 0.001*	< 0.001*	< 0.001*	< 0.001*

\*p-value significant; SD — standard deviation

In the ACR group, the glomerular diameter decreased 58% with an increase in the width of the urinary space 2.4-fold matched to the control group. With the use of RVS, the glomerular diameter increased 1.26-fold with a decrease in the width of the urinary space 57% equated to the ACR group. The glomerular diameter and width of urinary space in RVS and control groups were alike (Table 2).

In the ACR group, the diameter of the PCT increased 31%, while the height of their lining epithelium decreased 62% compared to the control group. With the use of RVS, the diameter of the PCT decreased 18%, while the height of their lining epithelium increased 1.4-fold equated to the ACR group. Both parameters were comparable in RVS and control groups (Table 2).

## DISCUSSION

Acrylamide induced glomerular renal affection in the form of shrinkage and distortion of the glomeruli with wrinkling of their basement membranes and widening of the urinary spaces. Adding, degenerative tubular changes were markedly present in the PCT. The necrotic tubular cells exhibited cytoplasmic vacuolation with desquamated epithelial cells within the tubular lumen. Adding, ACR induced massive inflammatory cellular infiltration with congestion of glomerular blood vessels.

Acrylamide induced fibrosis that was established by a 9-fold increase deposition of collagen fibres in the basement membrane of the glomerular capillaries. Such collagen fibres can be the result of the epidermal growth factor that stimulates fibroblast proliferation and collagen synthesis [14].

The basement membranes of the renal corpuscles and renal tubules of the ACR group showed an intense

PAS reaction (42% higher than the control group). Thickening of the tubular basement membrane is a common feature of atrophy and may be associated with hyalinosis [14]. Thickening is also a possible cause of reduced active transport in PCT causing micro-albuminuria [14].

The brush border of the PCT in the ACR group was interrupted. Loss of the brush border is the earliest morphological sign of impaired proximal tubular function [17, 27]. Furthermore, loss of the brush border affects the reabsorptive power of the PCT with loss of glucose, salts, and large amounts of water in urine [17, 27]. This mostly explains the observed serological changes (the elevated serum levels of urea and creatinine). The diameter of the PCT increased 31% in the ACR group which mostly is a compensatory mechanism to conserve the renal function [15].

Oxidative stress is the main pathogenic mechanism through which ACR induces renal damage. Oxidative stress is a shift in the balance between oxidants and antioxidants in favour of oxidants [3]. Many researchers proved the oxidative stress role of ACR over the kidney [23]. The values of the oxidative marker (MDA) in the ACR group increased by 1.4-fold, while the anti-oxidative marker (GSH) decreased by 1.3-fold. Oxidative stress creates oxygen free radical (ROS) that reacts with numerous biomolecules in the cell, leading eventually to oxidative damage [16]. ROS is scavenged by several cellular defence mechanisms involving non-enzymatic (GSH). GSH peroxidase proteins convert hydrogen peroxide to water and lipid peroxides to their respective alcohols [30]. The prolonged use of ACR decreased the activities GSH. This consequence in augmented production of the  $\text{O}_2^-$  and  $\text{H}_2\text{O}_2$  that outcomes the production of  $\text{OH}^-$  [11]. Many researchers believed that MDA's level is sufficient

proof of oxidative stress [13] and higher value of MDA revealed an increase in lipid peroxidation.

Apoptosis is also another pathogenic mechanism through which ACR induced renal affection [23]. BAX reaction increased 4.5-fold in the ACR group. BAX exerts proapoptotic activity [29].

Resveratrol, as one of the flavonoids, affords a high protection to the kidney as the glomeruli and renal tubules were nearly normal. Compared to ACR group, the content of collagen fibres and the PAS reaction of the renal tubules decreased by 70, 19%. Adding, the levels of creatinine and urea decreased by 51, 47%.

Many researchers recorded the exogenous antioxidant protective effect of RVS over the kidney [21]. RVS induced such protective role through its antioxidant effect as the MDA level decreased by 45%, while the GSH level increased by 83% compared with the ACR group. RVS prevents superoxide production from uncoupled endothelial nitric oxide synthase and increases the expression of various antioxidant enzymes [30]. The antioxidant activity of many flavonoids is due to direct scavenging of oxygen-free radicals or excited oxygen species as well as inhibition of oxidative enzymes producing ROS [26].

Another protective mechanism to RVS is through its anti-apoptotic effect. With the use of RVS, the area percentage of the immunopositive BAX cells decreased 56% equated to the ACR group.

## CONCLUSIONS

In conclusion, ACR causes structural and functional disorders of the kidney. It induces kidney damage through oxidative stress and apoptosis. With the use of RVS, the normal kidney architecture was preserved with little structural changes. RVS exerts its protection through its anti-apoptotic and antioxidant features.

**Conflict of interest:** None declared

## REFERENCES

1. Abdel-Daim MM, Abd Eldaim MA, Hassan AGA. Trigonella foenum-graecum ameliorates acrylamide-induced toxicity in rats: Roles of oxidative stress, proinflammatory cytokines, and DNA damage. *Biochem Cell Biol.* 2015; 93(3): 192–198, doi: [10.1139/bcb-2014-0122](https://doi.org/10.1139/bcb-2014-0122), indexed in Pubmed: [25607344](https://pubmed.ncbi.nlm.nih.gov/25607344/).
2. Baskar G, Aiswarya R. Overview on mitigation of acrylamide in starchy fried and baked foods. *J Sci Food Agric.* 2018; 98(12): 4385–4394, doi: [10.1002/jsfa.9013](https://doi.org/10.1002/jsfa.9013), indexed in Pubmed: [29572830](https://pubmed.ncbi.nlm.nih.gov/29572830/).
3. Birben E, Sahiner UM, Sackesen C, et al. Oxidative stress and antioxidant defense. *World Allergy Organ J.* 2012; 5(1): 9–19, doi: [10.1097/WOX.0b013e3182439613](https://doi.org/10.1097/WOX.0b013e3182439613), indexed in Pubmed: [23268465](https://pubmed.ncbi.nlm.nih.gov/23268465/).
4. Buege J, Aust S. [30] Microsomal lipid peroxidation. *Methods Enzymol.* 1978: 302–310, doi: [10.1016/s0076-6879\(78\)52032-6](https://doi.org/10.1016/s0076-6879(78)52032-6).
5. Carere A. Genotoxicity and carcinogenicity of acrylamide: a critical review. *Ann Ist Super Sanita.* 2006; 42(2): 144–155, indexed in Pubmed: [17033134](https://pubmed.ncbi.nlm.nih.gov/17033134/).
6. Crowley C, Gillham B, Thorn MB. A direct enzymic method for the determination of reduced glutathione in blood and other tissues. *Biochem Med.* 1975; 13(3): 287–292, doi: [10.1016/0006-2944\(75\)90087-3](https://doi.org/10.1016/0006-2944(75)90087-3), indexed in Pubmed: [1203068](https://pubmed.ncbi.nlm.nih.gov/1203068/).
7. Dearfield KL, Abernathy CO, Ottley MS, et al. Acrylamide: its metabolism, developmental and reproductive effects, genotoxicity, and carcinogenicity. *Mutat Res.* 1988; 195(1): 45–77, doi: [10.1016/0165-1110\(88\)90015-2](https://doi.org/10.1016/0165-1110(88)90015-2), indexed in Pubmed: [3275881](https://pubmed.ncbi.nlm.nih.gov/3275881/).
8. Dudka J, Gieroba R, Korga A, et al. Different effects of resveratrol on dose-related Doxorubicin-induced heart and liver toxicity. *Evid Based Complement Alternat Med.* 2012; 2012: 606183, doi: [10.1155/2012/606183](https://doi.org/10.1155/2012/606183), indexed in Pubmed: [23258992](https://pubmed.ncbi.nlm.nih.gov/23258992/).
9. Ghorbel I, Elwej A, Fendri N, et al. Olive oil abrogates acrylamide induced nephrotoxicity by modulating biochemical and histological changes in rats. *Ren Fail.* 2017; 39(1): 236–245, doi: [10.1080/0886022X.2016.1256320](https://doi.org/10.1080/0886022X.2016.1256320), indexed in Pubmed: [27846768](https://pubmed.ncbi.nlm.nih.gov/27846768/).
10. Gülçin İ. Antioxidant activity of food constituents: an overview. *Arch Toxicol.* 2012; 86(3): 345–391, doi: [10.1007/s00204-011-0774-2](https://doi.org/10.1007/s00204-011-0774-2), indexed in Pubmed: [22102161](https://pubmed.ncbi.nlm.nih.gov/22102161/).
11. Halliwell B, Gutteridge JM. Free radicals and antioxidant protection: mechanisms and significance in toxicology and disease. *Hum Toxicol.* 1988; 7(1): 7–13, doi: [10.1177/096032718800700102](https://doi.org/10.1177/096032718800700102), indexed in Pubmed: [3278973](https://pubmed.ncbi.nlm.nih.gov/3278973/).
12. Kjuus H, Hansteen IL, Ryberg D, et al. Chromosome aberrations in tunnel workers exposed to acrylamide and N-methylolacrylamide. *Scand J Work Environ Health.* 2005; 31(4): 300–306, doi: [10.5271/sjweh.886](https://doi.org/10.5271/sjweh.886), indexed in Pubmed: [16161713](https://pubmed.ncbi.nlm.nih.gov/16161713/).
13. Kurt A, Tumkaya L, Turut H, et al. Protective effects of infliximab on lung injury induced by methotrexate. *Arch Bronconeumol.* 2015; 51(11): 551–557, doi: [10.1016/j.arbres.2015.03.018](https://doi.org/10.1016/j.arbres.2015.03.018), indexed in Pubmed: [26071367](https://pubmed.ncbi.nlm.nih.gov/26071367/).
14. Liu Y. Epithelial to mesenchymal transition in renal fibrogenesis: pathologic significance, molecular mechanism, and therapeutic intervention. *J Am Soc Nephrol.* 2004; 15(1): 1–12, doi: [10.1097/01.asn.0000106015.29070.e7](https://doi.org/10.1097/01.asn.0000106015.29070.e7), indexed in Pubmed: [14694152](https://pubmed.ncbi.nlm.nih.gov/14694152/).
15. McCance KL, Huether SE. *Pathophysiology: the biologic basis for disease in adults and children.* 7th ed. Elsevier, St. Louis, Missouri 2014.
16. McCord JM. Human disease, free radicals, and the oxidant/antioxidant balance. *Clin Biochem.* 1993; 26(5): 351–357, doi: [10.1016/0009-9120\(93\)90111-i](https://doi.org/10.1016/0009-9120(93)90111-i), indexed in Pubmed: [8299205](https://pubmed.ncbi.nlm.nih.gov/8299205/).
17. Nangaku M. Chronic hypoxia and tubulointerstitial injury: a final common pathway to end-stage renal failure. *J Am Soc Nephrol.* 2005; 17(1): 17–25, doi: [10.1681/asn.2005070757](https://doi.org/10.1681/asn.2005070757).

18. Qiao Y, Gao Ke, Wang Y, et al. Resveratrol ameliorates diabetic nephropathy in rats through negative regulation of the p38 MAPK/TGF- $\beta$ 1 pathway. *Exp Ther Med*. 2017; 13(6): 3223–3230, doi: [10.3892/etm.2017.4420](https://doi.org/10.3892/etm.2017.4420), indexed in Pubmed: [28588674](https://pubmed.ncbi.nlm.nih.gov/28588674/).
19. Rajeh NA, Al-Dhaheri NM. Antioxidant effect of vitamin E and 5-aminosalicylic acid on acrylamide induced kidney injury in rats. *Saudi Med J*. 2017; 38(2): 132–137, doi: [10.15537/smj.2017.2.16049](https://doi.org/10.15537/smj.2017.2.16049), indexed in Pubmed: [28133684](https://pubmed.ncbi.nlm.nih.gov/28133684/).
20. Ramos-Vara JA, Kiupel M, Baszler T, et al. Suggested guidelines for immunohistochemical techniques in veterinary diagnostic laboratories. *J Vet Diagn Invest*. 2008; 20(4): 393–413, doi: [10.1177/104063870802000401](https://doi.org/10.1177/104063870802000401), indexed in Pubmed: [18599844](https://pubmed.ncbi.nlm.nih.gov/18599844/).
21. Rašić D, Micek V, Klarić MS, et al. Oxidative stress as a mechanism of combined OTA and CTN toxicity in rat plasma, liver and kidney. *Hum Exp Toxicol*. 2019; 38(4): 434–445, doi: [10.1177/0960327118819049](https://doi.org/10.1177/0960327118819049), indexed in Pubmed: [30545268](https://pubmed.ncbi.nlm.nih.gov/30545268/).
22. Semla M, Goc Z, Martiniaková M, et al. Acrylamide: a common food toxin related to physiological functions and health. *Physiol Res*. 2017; 66(2): 205–217, doi: [10.33549/physiolres.933381](https://doi.org/10.33549/physiolres.933381), indexed in Pubmed: [27982682](https://pubmed.ncbi.nlm.nih.gov/27982682/).
23. Sengul E, Gelen V, Yildirim S, et al. The effects of selenium in acrylamide-induced nephrotoxicity in rats: roles of oxidative stress, inflammation, apoptosis, and DNA damage. *Biol Trace Elem Res*. 2021; 199(1): 173–184, doi: [10.1007/s12011-020-02111-0](https://doi.org/10.1007/s12011-020-02111-0), indexed in Pubmed: [32166561](https://pubmed.ncbi.nlm.nih.gov/32166561/).
24. Suvarna SK, Layton C, Bancroft JD. Bancroft's theory and practice of histological techniques. 8th ed. Elsevier, Oxford 2019.
25. Tareke E, Rydberg P, Karlsson P, et al. Acrylamide: a cooking carcinogen? *Chem Res Toxicol*. 2000; 13(6): 517–522, doi: [10.1021/tx9901938](https://doi.org/10.1021/tx9901938), indexed in Pubmed: [10858325](https://pubmed.ncbi.nlm.nih.gov/10858325/).
26. Terao J. Dietary flavonoids as antioxidants. *Forum of Nutr*. 2009; 87–94, doi: [10.1159/000212741](https://doi.org/10.1159/000212741).
27. Tirapelli LF, Barione DF, Trazzi BFM, et al. Comparison of two models for evaluation histopathology of experimental renal ischemia. *Transplant Proc*. 2009; 41(10): 4083–4087, doi: [10.1016/j.transproceed.2009.09.061](https://doi.org/10.1016/j.transproceed.2009.09.061), indexed in Pubmed: [20005345](https://pubmed.ncbi.nlm.nih.gov/20005345/).
28. Totani N, Yawata M, Ojiri Y, et al. Effects of trace acrylamide intake in Wistar rats. *J Oleo Sci*. 2007; 56(9): 501–506, doi: [10.5650/jos.56.501](https://doi.org/10.5650/jos.56.501), indexed in Pubmed: [17898518](https://pubmed.ncbi.nlm.nih.gov/17898518/).
29. Tsujimoto Y. Role of Bcl-2 family proteins in apoptosis: apoptosomes or mitochondria? *Genes Cells*. 1998; 3(11): 697–707, doi: [10.1046/j.1365-2443.1998.00223.x](https://doi.org/10.1046/j.1365-2443.1998.00223.x), indexed in Pubmed: [9990505](https://pubmed.ncbi.nlm.nih.gov/9990505/).
30. Xia N, Daiber A, Förstermann U, et al. Antioxidant effects of resveratrol in the cardiovascular system. *Br J Pharmacol*. 2017; 174(12): 1633–1646, doi: [10.1111/bph.13492](https://doi.org/10.1111/bph.13492), indexed in Pubmed: [27058985](https://pubmed.ncbi.nlm.nih.gov/27058985/).
31. Zhang Y. Study on reduction of acrylamide in fried bread sticks by addition of antioxidant of bamboo leaves and extract of green tea. *Asia Pac J Clin Nutr*. 2007; 16(Suppl 1): 131–136.
32. Zhang L, Yang J, Eastwood GM, et al. Extended daily dialysis versus continuous renal replacement therapy for acute kidney injury: a meta-analysis. *Am J Kidney Dis*. 2015; 66(2): 322–330, doi: [10.1053/j.ajkd.2015.02.328](https://doi.org/10.1053/j.ajkd.2015.02.328), indexed in Pubmed: [25843704](https://pubmed.ncbi.nlm.nih.gov/25843704/).