

A rare occurrence of persistent hypoglossal artery and its clinical significance

S. Yasin¹ , R. Tasdemir² , O.F. Cihan³ 

¹Department of Neurology, Faculty of Medicine, Gaziantep University, Gaziantep, Turkey

²Department of Anatomy, Faculty of Medicine, Gaziantep Islam, Science and Technology University, Gaziantep, Turkey

³Department of Anatomy, Faculty of Medicine, Gaziantep University, Gaziantep, Turkey

[Received: 16 September 2022; Accepted: 6 December 2022; Early publication date: 22 December 2022]

Persistent hypoglossal artery (PHA) is an embryological vascular variation mostly originating from the internal carotid artery. The presence of PHA has been associated with the incidence of some diseases such as cerebral ischaemia, atherosclerosis, and aneurysm. Here, a very rare case of PHA that was discovered incidentally by digital subtraction angiography in Turkey is reported. Endovascular stenting was not performed for this patient. Also, its clinical importance is discussed. (Folia Morphol 2023; 82, 4: 953–956)

Key words: persistent hypoglossal artery, digital subtraction angiography, rare variation, vertebrobasilar insufficiency

INTRODUCTION

Presegmental arteries are the carotid-vertebrobasilar anastomoses that supply blood from the internal carotid artery (ICA) to the vertebrobasilar system in the embryonic period. Four pairs of presegmental arteries, which are named for neighbouring structures, arise from the primitive ICA: the trigeminal, otic, hypoglossal, and proatlantal intersegmental arteries (Fig. 1). After the posterior communicating arteries develop and the vertebral and basilar arteries join, these temporary collateral vessels disappear at approximately 5 weeks of gestation [8]. The first to regress is the otic artery, followed by the hypoglossal artery, the trigeminal artery, and then the proatlantal intersegmental arteries [7]. Rarely, primitive carotid-vertebrobasilar anastomoses do not disappear and they persist into adult life. Persistent hypoglossal artery (PHA) is the second most common anastomosis with an incidence of 0.03–0.09% [9]. It has been reported that PHA is more common in females and on the left side [3]. PHA usually originates from the ICA at the C1–C3 level and rarely from the external carotid artery.




Figure 1. Persistent carotid vertebo-basilar anastomoses. Black and green arrows show trigeminal artery, blue arrow otic artery, yellow arrow hypoglossal artery and purple arrow indicates proatlantal artery. Srinivas et al. (2016) [9] adapted from his work.

CASE REPORT

A 57-year-old male patient presented to our clinic with complaints of numbness of the left arm and left leg, impaired balance, and severe dizziness. On neurological

Address for correspondence: Dr. R. Tasdemir, Department of Anatomy, Faculty of Medicine, Gaziantep Islam, Science and Technology University, Gaziantep, Turkey, tel: +90 505 854 8754, e-mail: rabiatsdmr@gmail.com

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.




Figure 2. Red arrow shows severe stenosis of the persistent hypoglossal artery (PHA), proximal to the right internal carotid artery, and distal to the bulb on the right side of the patient on digital subtraction angiography imaging; yellow arrow indicates PHA. Green arrow shows common carotid artery, blue arrow external carotid artery and purple arrow proximal right internal carotid artery.




Figure 4. Left vertebral artery appears hypoplastic compared to right side persistent hypoglossal artery (PHA). Yellow arrow indicates left vertebral artery, green arrow left PHA, blue arrow right internal carotid artery and white arrow left subclavian artery.




Figure 3. Red arrow shows stenosis of the left internal carotid artery on digital subtraction angiography imaging. Green arrow shows common carotid artery and blue arrow external carotid artery.




Figure 5. Yellow arrows show diffuse hyperintense millimetric ischaemic gliotic foci in the supratentorial white matter on brain magnetic resonance imaging T2-FLAIR sequence.

examination, the patient showed no motor function loss and his cerebellar tests were normal. The patient reported that dizziness increased when standing up and partially improved after lying down. Doppler ultrasonography of the carotid vertebral artery showed the presence of PHA on the right side. Upon detecting 70–99% stenosis of the proximal right ICA and 70% stenosis of the proximal left ICA along with PHA, imaging with digital subtraction angiography (DSA) was planned. DSA revealed PHA on the right side and 95% stenosis of the PHA, proximal to the right ICA and distal to the bulb (Fig. 2). The patient

also had atherosclerotic stenosis causing 75% stenosis in the proximal left ICA (Fig. 3, Suppl. Video 1). The left vertebral artery appeared hypoplastic compared to the right-sided PHA (Fig. 4). In addition, widespread ischaemic gliotic foci were observed in the supratentorial white matter on magnetic resonance images of the patient (Fig. 5). Endarterectomy was planned for 95% stenosis on the right side of the patient because of anatomical incompatibility. Elective endovascular stenting for 70% stenosis proximal to the left ICA was scheduled following endarterectomy at our centre.

DISCUSSION

Four criteria have been determined for the diagnosis of PHA: 1) the hypoglossal artery originates from the ICA as a robust branch at the C1–C3 level, 2) enters the posterior cranial fossa through the hypoglossal canal, 3) joins the basilar artery just beyond its origin near the pontomedullary junction, 4) the posterior communicating artery is either absent or not visible on angiography [1]. Vertebral arteries may be hypoplastic or aplastic [7]. It was reported that the diameter of the anterior condylar foramen is normally 6–7 mm, which can increase up to 8 mm in the presence of PHA [4].

Begg (1961) [see 4] first demonstrated the presence of PHA on angiographic imaging. PHA is usually detected incidentally on imaging studies and is often asymptomatic. However, considering the fact that anterior and posterior cerebral circulation depends on the blood supplied by the ICA, it is crucial to identify such a variation before deciding how to perform surgical or endovascular interventions [6].

Although PHA is considered an anatomical variation of embryological origin, cerebral pathologies should also be considered because any aneurysm may be accompanied by vascular diseases or atherosclerosis [7]. Presence of PHA may be associated with abnormal vessel wall structure. Accordingly, as reported by former studies, exposure of the basilar artery to excessive haemodynamic stress may cause the development of aneurysms and arteriovenous malformations. Cases associated with ruptured aneurysms and arteriovenous malformations have been reported in the literature [6].

Persistent hypoglossal artery has also been associated with atherosclerotic cerebrovascular disease. Atherosclerotic plaque may develop because PHA originates from the ICA, causing haemodynamics similar to that of the carotid bulb. The presence of a plaque in this region is extremely important in patients with carotid and vertebrobasilar ischaemia because PHA generally supplies most of the posterior circulation [6]. Additionally, there are studies in the literature showing the presence of posterior cerebral artery fenestrations and cerebral infarction with PHA [7, 10, 11]. As a matter of fact, areas of atherosclerosis and cerebral infarction were observed in the present study, which is consistent with the literature data.

Angiography is required before performing carotid endarterectomy and surgical procedures to the skull base region. This is even more important in the

absence of the vertebral artery, since in that case the ICA will be the sole source of cerebral blood supply and it will be difficult to maintain cerebral perfusion during surgery to this region.

Although the incidence of PHA varies in different parts of the world, only 7 cases were reported in Turkey based on our literature search [2, 4, 5, 7, 8, 11, 12]. The fact that so few cases were reported in Turkey makes the present case report even more noteworthy.

CONCLUSIONS

While dizziness has many possible causes, it should be borne in mind that a rare variation such as PHA and vertebrobasilar insufficiency as well as carotid artery stenosis, as emerges in this case, may also cause dizziness. DSA may be considered for the diagnosis of PHA in patients with unexplained dizziness and impaired balance. We believe that this case is representative of a very rare variation and will make a valuable contribution to the literature since there are few cases reported from Turkey.

Acknowledgements

This study was conducted in accordance with the ethical standards and the 1964 Declaration of Helsinki and its later amendments. Written informed consent was obtained from the patient.

All authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest or non-financial interest in the subject matter or materials discussed in this case report.

Conflict of interest: None declared

REFERENCES

1. Albay S, Kastamoni Y, Koyuncu E. Embriyonal Kalıntı Arterler. *SDÜ Tıp Fakültesi Dergisi*. 2012; 19: 62–67.
2. Aysel T, Tülay Ö, Turhan C, et al. Multidetector CT angiography of the persistent primitive hypoglossal artery. *Eur J Radiol Extra*. 2008; 68(2): e63–e66, doi: [10.1016/j.ejrex.2008.06.013](https://doi.org/10.1016/j.ejrex.2008.06.013).
3. Coulier B. Persistent hypoglossal artery. *J Belg Soc Radiol*. 2018; 102(1): 28, doi: [10.5334/jbsr.1481](https://doi.org/10.5334/jbsr.1481), indexed in Pubmed: [30039040](https://pubmed.ncbi.nlm.nih.gov/30039040/).
4. Memiş A, Güney B. Persistan hipoglossal arterin MR anjiyografi ile görüntülenmesi. *Türk J Diagn İntervent Radiol*. 2001; 7: 248–251.
5. Numan F. A successful treatment with carotid arterial stenting for symptomatic severe internal carotid artery stenosis with ipsilateral persistent primitive hypoglossal artery and aplasia of the A1 segment of anterior cerebral artery: a case report. *Turkish J Thoracic Cardiovasc Surg*. 2013; 21(2): 440–444, doi: [10.5606/tgkdc.dergisi.2013.5162](https://doi.org/10.5606/tgkdc.dergisi.2013.5162).

6. Paraskevas GK, Tsitsopoulos PPh, Papaziogas B, et al. Persistent primitive hypoglossal artery: an incidental autopsy finding and its significance in clinical practice. *Folia Morphol.* 2007; 66(2): 143–147, indexed in Pubmed: [17594674](#).
7. Pasaoglu L, Hatipoglu HG, Vural M, et al. Persistent primitive hypoglossal artery and fenestration of posterior cerebral artery: CT and MR angiography. *Neurocirugia (Astur)*. 2009; 20(6): 563–566, doi: [10.1016/s1130-1473\(09\)70137-x](#), indexed in Pubmed: [19967323](#).
8. Petik B, Colak D, Sirik M, et al. Rare variants of carotid-vertebrobasilar anastomoses. *J Belg Soc Radiol.* 2016; 100(1): 74, doi: [10.5334/jbr-btr.1167](#), indexed in Pubmed: [30151473](#).
9. Srinivas MR, Vedaraju KS, Manjappa BH, et al. Persistent primitive hypoglossal artery (PPHA): a rare anomaly with literature review. *J Clin Diagn Res.* 2016; 10(1): TD13–TD14, doi: [10.7860/JCDR/2016/15556.7116](#), indexed in Pubmed: [26894148](#).
10. Tse GH, Martin A, Dyde RA, et al. Persistent hypoglossal artery aneurysm: case report and qualitative systematic review. *Interv Neuroradiol.* 2019; 25(2): 164–171, doi: [10.1177/1591019918809087](#), indexed in Pubmed: [30394836](#).
11. Uysal E, Velioglu M, Kara E, et al. Persistent hypoglossal artery associated with a ruptured ipsilateral posterior inferior cerebellar artery aneurysm. A case report. *Neuroradiol J.* 2007; 20(5): 570–573, doi: [10.1177/197140090702000516](#), indexed in Pubmed: [24299948](#).
12. Yilmaz E, Ilgit E, Taner D. Primitive persistent carotid-basilar and carotid-vertebral anastomoses: a report of seven cases and a review of the literature. *Clin Anat.* 1995; 8(1): 36–43, doi: [10.1002/ca.980080107](#), indexed in Pubmed: [7697511](#).