

The evaluation of vasculature in post-mortem angio-computed tomography for anatomy research purposes: method description based on coeliac trunk analysis

Jakub Batko¹ , Radosław Chmiel¹, Aleksiej Juszcak¹, Jerzy A. Walocha¹, Artur Moskała², Michał Bonczar¹, Artur Pasternak¹, Krzysztof Woźniak²

¹Department of Anatomy, Jagiellonian University Medical College, Krakow, Poland

²Department of Forensic Medicine, Jagiellonian University Medical College, Krakow, Poland

[Received: 31 May 2022; Accepted: 5 July 2023; Early publication date: 4 August 2023]

Background: Anatomical research based on deceased body specimens is a time-consuming process that requires a great deal of skill and time to perform correctly. Three-dimensional medical image analysis is an excellent tool for anatomic evaluation, but it often includes patients with comorbidities in the study group, which can skew the results. The purpose of this study was to develop and evaluate methods for anatomic research based on postmortem contrast-enhanced computed tomography angiography 3D reconstruction of the coeliac trunk.

Materials and methods: Postmortem contrast-enhanced computed tomography angiography of 105 (28.6% female, age 50.8 ± 18.7) decedents without abdominal trauma or tumour was analysed. The abdominal portion of the aorta and the coeliac trunk with its branches were reconstructed and evaluated. The type of coeliac trunk was evaluated. The results were analysed.

Results: The coeliac trunk, splenic artery, and common hepatic artery were visualized in all cases. The left gastric artery was visible in 97.1% of cases. The dorsal pancreatic artery was visualized in 61.0% of cases. The most common type of coeliac trunk was 1 (88.6%), and the rarest types were 2, 3, and 6 (1.0%). We observed 4 morphologies of the truncus coeliacus that did not fit the classification presented previously.

Conclusions: This study has demonstrated that three-dimensional reconstruction of postmortem contrast-enhanced computed tomography is an excellent tool for performing accurate morphometric analyses for anatomic research purposes. This method can serve as a source for anatomic studies in the healthy population. (Folia Morphol 2024; 83, 2: 343–347)

Keywords: post-mortem CT, coeliac trunk, post-mortem 3D visualisation, RadiAnt

Address for correspondence: Artur Pasternak, Department of Anatomy, Jagiellonian University Medical College, ul. Kopernika 12, 31–034 Krakow, Poland; e-mail: artur.pasternak@uj.edu.pl

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

INTRODUCTION

Anatomical research has long been associated with dissection. In this form of research, the researcher is exposed to fumes of formaldehyde (formalin) for a long time, which can lead to serious health complications, including the development of cancer [2, 3]. On the other hand, dissection is a time-consuming process that is highly dependent on manual skills. If performed incorrectly, it cannot be recovered. To avoid these problems, modern anatomy research is supported by studies based on medical imaging and its three-dimensional reconstruction [1, 5, 8]. However, in order to perform a correct analysis, patients should not have comorbidities, which is a challenge because medical imaging is usually performed as part of the diagnostic process. The data collected during such an examination can have great clinical implications for the surgical treatment of various diseases and the diagnosis of certain disorders. To solve the above problems, postmortem contrast-enhanced computed tomography angiography can be considered a valuable source for anatomical research. It is performed in selected departments of forensic medicine as a valuable tool to support autopsy in the search for cause of death and other important information [4, 14]. However, the visualization method could also be used in anatomy departments to provide medical images of donors for future research and teaching purposes.

Therefore, the aim of this study is to evaluate the usefulness of postmortem contrast-enhanced computed tomography angiography for anatomic studies, develop the research method, and discuss its limitations. To evaluate this method in practice, we performed studies on a previously well-described region of the truncus coeliacus.

MATERIALS AND METHODS

Study population

Postmortem contrast-enhanced computed tomography angiography (CTA) scans (16-detector row computed tomography Somatom Emotion, Siemens AG, Hanau, Germany) of 105 (28.6% female, age 50.8 ± 18.7) human bodies imaged between 2012 and 2017 were retrospectively evaluated. Scanning parameters were 130 kVp, 50 mAs, and 240 mAs, reconstructed slice thickness of 0.75 mm, collimation 16×0.6 mm. The oily liquid contrast agent: 6% Angiofil (Fumedica, Muri, Switzerland) kerosene oil solution via unilateral

or bilateral access to the femoral vessels according to the procedure used by the Technical Working Group Postmortem Angiography Methods [4]. Results were obtained using RadiAnt DICOM Viewer three-dimensional reconstruction software (Medixant. RadiAnt DICOM Viewer [Software]. Version 2022.1. Feb 10, 2022. URL: <https://www.radiantviewer.com>).

Reconstructions for the present work were based on arterial phase acquisition results. Three-dimensional volume rendering reconstructions were performed and evaluated. The quality of imaging for research purposes was graded: Images were classified as high quality if all coeliac main branches with their branches were fully visible, medium quality if one coeliac branch was not fully visible, and low quality if one of the coeliac main branches was not visible or more than two coeliac branches were not visible.

Analysis of the coeliac trunk

The types of the coeliac trunk were taken from the research of Adachi [4] and determined by two different researchers. If the evaluation of the type differed between the researchers, the evaluation was repeated and reported as the final result.

The diameter of the aorta, coeliac trunk and each of its branches was measured. The distances from the origin of the coeliac trunk to each branch, the angle of origin, and the location were measured and evaluated. The distance between the origin of the coeliac trunk and the origin of the abdominal aorta and the aortic bifurcation was measured.

Statistical analyses

Data were analysed using IBM SPSS Statistics 28.0 (Predictive Solutions, Pittsburgh, PA, USA). Categorical variables are presented as number (n) or percentages. Quantitative variables are presented as mean with standard deviation. Normal distribution was examined using the Shapiro-Wilk test. Group comparisons between quantitative variables were performed using the U-Mann Whitney test. The p value < 0.05 was considered significant.

Ethical statement

This study was approved by the Bioethical Committee of the Jagiellonian University, Cracow, Poland (No. 1072.6120.241.2021 approved in 2021). The study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki.



Figure 1. Visualization of abdominal aorta branches. Good quality visualization of coeliac trunk. 1 — coeliac trunk; 2 — common hepatic artery; 3 — left gastric artery; 4 — splenic artery; 5 — superior mesenteric artery; 6 — left renal artery; 7 — right renal artery; 8 — aorta bifurcation.

RESULTS

According to the computed tomography protocol, 61% of the images were classified as very good quality and 39.0% as intermediate quality. There were no poor quality images in this study. The main cause of poor to moderate quality was visualization of the dorsal pancreatic artery, which was poorly visualized in these cases. In 2.9% of cases, intermediate quality was determined because of poor visualization of the branches of the left gastric artery.

All coeliac branches were reconstructed and visualized. Semiautomated reconstruction was challenging in the pancreatic region. In the other regions, the terminal branches of each coeliac trunk were captured in 97.1% of cases. (Fig. 1).

The most common coeliac trunk type was 1 (88.6%). Types 2, 3, and 6 were present in only one case. Type 5 was present in 5 cases (4.8%). 4 cases (3.8%) did not meet the classification criteria (in 3 cases, we observed multiple origins of the left gastric artery, and in one case, we observed a common origin of the hepatic artery from the superior mesenteric artery and an additional trunk for the left and right gastric artery). The diameter of the coeliac trunk was 7.2 ± 1.7 mm. The distance between the coeliac trunk and the origin of the abdominal aorta was 7.4 ± 2.9 cm




Figure 2. Coeliac trunk origin distance to abdominal aorta origin and to aorta bifurcation.

and between the coeliac trunk and the aortic bifurcation was 13.2 ± 1.5 cm (Fig. 2).

The diameter of the splenic artery was 6.1 ± 1.3 mm. The length from the origin of the coeliac trunk to the

splenic artery was 20.0 ± 6.6 mm, the angle of origin was $106.5 \pm 28.1^\circ$, and the origin was most often anterior in 35.2% of cases and least often posterior in 1.0%. Pancreatic branches of the splenic artery were visible in 61% of cases.

The left gastric artery was visualized in 97.1% of cases. The diameter of the left gastric artery was 3.8 ± 1.1 mm. The length from the coeliac trunk to the origin of the left gastric artery was $14.7 \text{ mm} \pm \pm 5.4$ mm, the angle of origin was $126.4 \pm 31.2^\circ$, and the site of origin was upward in 80.8% of cases and downward in 1.0% of cases. The terminal branches of the left gastric artery were visualized in 100% of the cases in which the left gastric artery was visible.

The diameter of the common hepatic artery was 5.1 ± 1.2 mm. The length from the origin of the coeliac trunk to the common hepatic artery was $12.4 \pm \pm 9.2$ mm, the angle of origin was $121.2 \pm 30.1^\circ$, and the site of origin was right in 49.5% of cases and left and downward in 1.0% of cases. The terminal branches of the common hepatic artery were visible in all cases. Detailed coeliac trunk branches morphometry comparison can be found in the Table 1.

In 14.3% of cases, we observed that the diaphragmatic arteries originated from the truncus coeliacus. In 2 decedents (1.9%), we found the origin of the gastroduodenal artery in the coeliac trunk.

DISCUSSION

This study describes a new useful research method for anatomical studies. It is based on postmortem CT visualization, which allows detailed visualization of anatomical structures. Significant differences between postmortem and normal low dose CT should be mentioned. In postmortem CT, higher radiation doses are used for better imaging quality so as not to harm the patient. In addition, a hydrophobic paraffin-based contrast agent is used, as opposed to the hydrophilic iodine contrast agent used in low-dose radiation CT patients. These features provide more detailed anatomical im-

ages compared to steady-state visualization methods. It should be considered especially for studies of small anatomic structures that cannot be visualized by other means. Postmortem CTA can be a useful tool for anatomical research. It is a very sufficient source that could be performed more frequently for scientific purposes. Collaboration between the departments of anatomy and forensic science could also be a part of the collection of imaging data for scientific and didactic purposes. However, there should be clear inclusion criteria for ethical consultation during data collection so that donors accept the proposed analysis. The process of three-dimensional reconstruction is reproducible, simpler, and faster compared with classical anatomical dissection, and it allows the analysis of anatomical structures in their correct localization. This method can increase the quality and quantity of anatomical examinations and improve safety because researchers are not exposed to formalin. All of these factors can lead to more reliable anatomical studies based on imaging of younger, deceased human bodies, with no or well-defined clinical or disease biases. The collection of computed tomography scans and multicentre collaboration may lead to the development of a large database of computed tomography scans, which can greatly improve anatomic research. Our study revealed some rare variations of the truncus coeliacus. Similar and other rare cases have been previously reported in the literature [6, 10]. Postmortem CTA could be a suitable tool to study rare variations of the truncus coeliacus in a representative, large population. Our results are comparable to those of other studies based on dissection, medical image analysis, and reconstruction, which further demonstrates the usefulness and quality of the proposed method [1, 5, 7, 9, 11–13].

The main limitations of using this method are access to CT scanners and qualified technicians, similar to the contrast injection method. In addition, as in the present case, rapidly decomposing parts of the human body, such as the pancreas, and their surroundings may be difficult to visualize.

Table 1. Coeliac trunk branches morphometry comparison

	Common hepatic artery		Left gastric artery		Splenic artery		P
	Mean	SD	Mean	SD	Mean	SD	
Artery origin angle [°]	121.2	30.1	126.4	31.2	106.5	28.1	< 0.001
Distance from CT origin [mm]	20.6	8.5	14.7	5.4	20	6.6	< 0.001
Artery origin diameter [mm]	5.1	1.2	3.8	1.1	6.1	1.3	< 0.001

CT — coeliac trunk.

CONCLUSIONS

Three-dimensional reconstruction of postmortem contrast-enhanced computed tomography is an excellent tool for performing precise morphometric analyses for anatomic research purposes. Visualization of vessels near rapidly decomposing organs can be challenging. Three-dimensional reconstruction analysis with contrast-enhanced computed tomography after death may be an alternative for dissection studies.

ARTICLE INFORMATION AND DECLARATIONS

Ethical statement

This study was approved by the Bioethical Committee of the Jagiellonian University, Cracow, Poland (No. 1072.6120.241.2021 approved in 2021). The study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki.

Funding

This study was supported by the National Science Centre (NCN), grant No. K/MNT/000184.

Funding sources had no involvement in study design, collection, analysis and interpretation of data, in writing of the report and in the decision to submit the article for publication.

Conflict of interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

1. Araujo Neto SA, Franca HA, de Mello Júnior CF, et al. Anatomical variations of the celiac trunk and hepatic arterial system: an analysis using multidetector computed tomography angiography. *Radiol Bras.* 2015; 48(6): 358–362, doi: [10.1590/0100-3984.2014.0100](https://doi.org/10.1590/0100-3984.2014.0100), indexed in Pubmed: [26811552](https://pubmed.ncbi.nlm.nih.gov/26811552/).
2. Cogliano VJ, Grosse Y, Baan RA, et al. Working Group for Volume 88. Meeting report: summary of IARC monographs on formaldehyde, 2-butoxyethanol, and 1-tert-butoxy-2-propanol. *Environ Health Perspect.* 2005; 113(9): 1205–1208, doi: [10.1289/ehp.7542](https://doi.org/10.1289/ehp.7542), indexed in Pubmed: [16140628](https://pubmed.ncbi.nlm.nih.gov/16140628/).
3. Dimenstein I. A pragmatic approach to formalin safety in anatomical pathology: table 1. *Lab Med.* 2009; 40(12): 740–746, doi: [10.1309/lmq1hdfd4un0wwbp](https://doi.org/10.1309/lmq1hdfd4un0wwbp).
4. Grabherr S, Heinemann A, Vogel H, et al. Postmortem CT angiography compared with autopsy: a forensic multicenter study. *Radiology.* 2018; 288(1): 270–276, doi: [10.1148/radiol.2018170559](https://doi.org/10.1148/radiol.2018170559), indexed in Pubmed: [29714682](https://pubmed.ncbi.nlm.nih.gov/29714682/).
5. Juszcak A, Czyżowski J, Mazurek A, et al. Anatomical variants of coeliac trunk in Polish population using multi-detector computed tomography angiography. *Folia Morphol.* 2021; 80(2): 290–296, doi: [10.5603/FM.a2020.0051](https://doi.org/10.5603/FM.a2020.0051), indexed in Pubmed: [32394417](https://pubmed.ncbi.nlm.nih.gov/32394417/).
6. Juszcak A, Czyżowski J, Mazurek A, et al. Unusual variations in the branching pattern of the coeliac trunk and their clinical significance. *Folia Morphol.* 2021; 80(2): 283–289, doi: [10.5603/FM.a2020.0067](https://doi.org/10.5603/FM.a2020.0067), indexed in Pubmed: [32639575](https://pubmed.ncbi.nlm.nih.gov/32639575/).
7. Juszcak A, Mazurek A, Walocha JA, et al. Coeliac trunk and its anatomic variations: a cadaveric study. *Folia Morphol.* 2021; 80(1): 114–121, doi: [10.5603/FM.a2020.0042](https://doi.org/10.5603/FM.a2020.0042), indexed in Pubmed: [32301103](https://pubmed.ncbi.nlm.nih.gov/32301103/).
8. Marco-Clement I, Martinez-Barco A, Ahumada N, et al. Anatomical variations of the celiac trunk: cadaveric and radiological study. *Surg Radiol Anat.* 2016; 38(4): 501–510, doi: [10.1007/s00276-015-1542-4](https://doi.org/10.1007/s00276-015-1542-4), indexed in Pubmed: [26267305](https://pubmed.ncbi.nlm.nih.gov/26267305/).
9. Marco-Clement I, Martinez-Barco A, Ahumada N, et al. Anatomical variations of the celiac trunk: cadaveric and radiological study. *Surg Radiol Anat.* 2016; 38(4): 501–510, doi: [10.1007/s00276-015-1542-4](https://doi.org/10.1007/s00276-015-1542-4), indexed in Pubmed: [26267305](https://pubmed.ncbi.nlm.nih.gov/26267305/).
10. Mazurek A, Juszcak A, Walocha JA, et al. Rare combined variations of the coeliac trunk, accessory hepatic and gastric arteries with co-occurrence of double cystic arteries. *Folia Morphol.* 2021; 80(2): 460–466, doi: [10.5603/FM.a2020.0052](https://doi.org/10.5603/FM.a2020.0052), indexed in Pubmed: [32459367](https://pubmed.ncbi.nlm.nih.gov/32459367/).
11. Olewnik Ł, Wysiadecki G, Polguy M, et al. Types of coeliac trunk branching including accessory hepatic arteries: a new point of view based on cadaveric study. *Folia Morphol.* 2017; 76(4): 660–667, doi: [10.5603/FM.a2017.0053](https://doi.org/10.5603/FM.a2017.0053), indexed in Pubmed: [28612916](https://pubmed.ncbi.nlm.nih.gov/28612916/).
12. Pinal-Garcia DF, Nuno-Guzman CM, Gonzalez-Gonzalez ME, et al. The celiac trunk and its anatomical variations: a cadaveric study. *J Clin Med Res.* 2018; 10(4): 321–329, doi: [10.14740/jocmr3356w](https://doi.org/10.14740/jocmr3356w), indexed in Pubmed: [29511421](https://pubmed.ncbi.nlm.nih.gov/29511421/).
13. Punja R, Punja D. Exploration of the coeliac trunk for its morphometry and positional relationships using cadaveric and radiological methods. *Transl Res Anat.* 2021; 24: 100116, doi: [10.1016/j.tria.2021.100116](https://doi.org/10.1016/j.tria.2021.100116).
14. Woźniak K, Rzepecka-Woźniak E, Moskała A, et al. Weapon identification using antemortem computed tomography with virtual 3D and rapid prototype modeling — a report in a case of blunt force head injury. *Forensic Sci Int.* 2012; 222(1-3): e29–e32, doi: [10.1016/j.forsciint.2012.06.012](https://doi.org/10.1016/j.forsciint.2012.06.012), indexed in Pubmed: [22748480](https://pubmed.ncbi.nlm.nih.gov/22748480/).