

# Coexistence of the 677C>T and 1298A>C MTHFR polymorphisms and its significance in the population of Polish women

Znaczenie współwystępowania polimorfizmów 677C>T oraz 1298A>C genu MTHFR u kobiet w populacji polskiej

Hubert Wolski<sup>1,2</sup>, Maria Kocięcka<sup>1,3</sup>, Aleksandra E. Mrozikiewicz<sup>4</sup>, Magdalena Barlik<sup>1,3</sup>, Grażyna Kurzawińska<sup>1,3</sup>

<sup>1</sup> Division of Perinatology and Women's Diseases, Poznan University of Medical Sciences, Poland

<sup>2</sup> Division of Gynecology and Obstetrics, Podhale Multidisciplinary Hospital, Nowy Targ, Poland

<sup>3</sup> Laboratory of Molecular Biology in the Division of Perinatology and Women's Diseases, Poznan University of Medical Sciences, Poznan, Poland

<sup>4</sup> Student Scientific Association of Department of Gynecological Endocrinology, Poznan University of Medical Sciences, Poland

## Abstract

**Objectives:** The aim of the study was to evaluate the frequency of the 677C>T and 1298A>C polymorphisms of the methylenetetrahydrofolate reductase (MTHFR) gene, as well as the coexistence of both these genetic variants in women from the Polish population.

**Material and methods:** A total of 662 women from the Polish population were enrolled in the study group. The frequency of the investigated genotypes of the 677C>T and 1298A>C polymorphisms of the MTHFR gene was analyzed with the use of PCR/RFLP methods.

**Results:** The frequency of the 677CC, 677CT and 677TT genotypes in the studied population of women was 50.60%, 39.88% and 9.52%, respectively. As to the 1298AA, 1298AC and 1298CC genotypes, the obtained results were as follows: 42.75%, 47.88% and 9.37%, respectively (Tables II and III). Simultaneous analysis revealed the most frequent coexistence of 677CC/1298AC (28.85%), 677CT/1298AA (20.85%) and 677CT/1298AC (19.03%) genotypes. The coexistence of 677CC/1298AA (12.39%), 677CC/1298CC (9.37%) and 677TT/1298AA (9.51%) genotypes was observed less frequently. In the studied population of Polish women, the coexistence of 677CT/1298CC, 677TT/1298AC and 677TT/1298CC genotypes has been not observed.

**Conclusions:** The frequency and coexistence of genotypes of the 677C>T and 1298A>C MTHFR gene polymorphisms in the studied population of Polish women is similar to other North-European populations. Women carriers of the mutated variants of both, 677C>T and 1298A>C polymorphisms of the MTHFR gene should receive special perinatal care in order to prevent fetal defects and thrombosis-related complications during pregnancy. It is vital to emphasize the significance of proper education of folate supplementation, especially in pregnant patients and women of reproductive age.

Key words: **MTHFR / genetic polymorphism / general population /**

## Corresponding author:

Hubert Wolski  
Division of Perinatology and Women's Disease  
Poznan University of Medical Sciences  
Tel. 618419223, fax. 618514746  
e-mail: hubertwolski@wp.pl

Otrzymano: 03.03.2015  
Zaakceptowano do druku: 01.04.2015

Hubert Wolski et al. Coexistence of the 677C&gt;T and 1298A&gt;C MTHFR polymorphisms and its significance in the population of Polish women.

## Streszczenie

**Cel:** Określenie częstości występowania polimorfizmów 677C>T i 1298A>C genu kodującego reduktazę metylenotetrahydrofolianową (MTHFR) oraz ocena współwystępowania obydwu wariantów genetycznych u kobiet w populacji polskiej.

**Materiał i metoda:** Do grupy badawczej włączono 662 kobiety z populacji polskiej. Częstość występowania genotypów polimorfizmów 677C>T oraz 1298A>C genu MTHFR została zbadana metodą PCR/RFLP.

**Wyniki:** W badanej grupie częstość występowania genotypów 677CC, 677CT oraz 677TT wynosiła odpowiednio 50,60%, 39,88% oraz 9,52%. Natomiast genotypy 1298AA, 1298AC oraz 1298CC występowały z częstością 42,75%, 47,88% oraz 9,37%. Jednocześnie analiza współwystępowania genotypów pozwoliła na określenie najczęściej występujących genotypów – 677CC/1298AC (28,85%), 677CT/1298AA (20,85%), 677CT/1298AC (19,03%). Współwystępowanie genotypów 677CC/1298AA (12,39%), 677CC/1298CC (9,37%), 677TT/1298AA (9,51%) było obserwowane rzadziej. W badanej populacji nie zaobserwowano współwystępowania polimorfizmów 677CT/1298CC, 677TT/1298AC, 677TT/1298CC.

**Wnioski:** Częstość występowania i współwystępowania polimorfizmów 677C>T i 1298A>C u kobiet w populacji polskiej jest porównywalna do częstości występowania tych wariantów w innych populacjach północnoeuropejskich. Kobiety, nosicielki zmutowanych genotypów polimorfizmów 677C>T i 1298A>C genu kodującego MTHFR, powinny zostać objęte specjalną opieką perinatalną, z powodu wzrostu ryzyka wystąpienia wad płodu oraz powikłań zakrzepowych w czasie ciąży. Należy zwrócić uwagę na właściwą edukację i suplementację folianami kobiet, szczególnie w wieku rozrodczym oraz w czasie ciąży.

Słowa kluczowe: **MTHFR / polimorfizm genetyczny / populacja ogólna /**

## Introduction

Proper function of the folate cycle is directly related to optimal human development and functioning. Adequate folate concentration prevents hyperhomocysteinemia, deficit of the synthesis of purine and pyrimidine, and methylation failure. Folate deficiency may lead to cardiovascular diseases, neurodegenerative and mental disorders, and even some cancers [1, 2, 3, 4, 5]. One of the key-enzymes in the folate cycle is methylenetetrahydrofolate reductase (MTHFR). Proper functioning of that enzyme is especially important in pregnant women [6, 7, 8].

In recent years, a few MTHFR gene polymorphisms have been described, including the 677C>T and 1298A>C as the most common. The 677C>T polymorphism concerns transition of cytosine into thymine, resulting in substitution of alanine for valine in the catalytic region of the enzyme (position 222) [9]. The MTHFR activity is 60-70% lower in carriers of the homozygotic 677TT genotype, which significantly increases the risk of hyperhomocysteinemia [10]. The 1298A>C genetic variant is the effect of substitution of adenine for cytosine in exon 7, which results in the conversion of glutamine to alanine (position 429), and also decreases the MTHFR activity [11].

The differences in the frequency of the 677C>T and 1298A>C polymorphisms have been demonstrated to depend on race and ethnicity [12,13,14]. The mutated 677TT variant concerns several to 15% percent of the Caucasian race. The mutated 1298CC genotype is present in up to 10% of the Caucasian population (frequency of the 1298C allele ranges from 27 to 36%) [12,13]. It has been shown that 15-20% of the Caucasian population are double heterozygotes (677CT/1298AC), and several percent are 677TT/1298AA carriers [11]. The coexistence of 677TT/1298AA and 677CT/1298AC genotypes lowers MTHFR activity to a great extent, by 60-70% and 40-50%, respectively) (Table I) [11, 15].

In recent years, it has been shown that disturbances of the folate metabolism could be connect with decreased MTHFR

activity and presence of the MTHFR polymorphisms. The consequences of the impaired folate metabolism in the general population are cardio-vascular, neurological, and psychiatric diseases [2,3,5]. In pregnant women, disturbed folate cycle constitutes a risk factor for neural tube defects, lip and palate cleft in the fetus, congenital heart and urinary tract defects, as well as Down syndrome. Also, it is the reasons for increased homocysteine level, leading to endothelium injury, thrombosis of the spiral arteries, and placental insufficiency. It may give raise to numerous obstetrical complications, including preeclampsia, recurrent miscarriages, fetal hypotrophy, preterm placental abruption, and intrauterine fetal death [6, 7, 8]. A correlation of the 677C>T and 1298A>G MTHFR polymorphisms with recurrent miscarriages has been suggested. It is also clear that the coexistence of both genetic polymorphisms has significant influence on the occurrence of the abovementioned disorders.

## Objectives

The aim of the study was to evaluate the frequency of the 677C>T and 1298A>C polymorphisms of the MTHFR gene, as well as the coexistence of both genetic variants in Polish women.

## Material and methods

### Study group

A total of 662 healthy Polish women from the general population (Wielkopolska region) were enrolled. Mean patient age was 30.12±4.52 years, range 18-43 years. Women with cardiovascular, renal, and hepatic diseases, as well as with cancers or genetic disorders were excluded from the study. The analysis was performed at the Laboratory of Molecular Biology, Department of Perinatology and Women's Diseases, Poznan University of Medical Sciences, between 20011-2013. Written informed consent was obtained from all participants. Local Ethics Committee approved of the study.

Hubert Wolski et al. Coexistence of the 677C>T and 1298A>C MTHFR polymorphisms and its significance in the population of Polish women.

**Table I.** MTHFR activity [%] according to the presence of 677C>T and 1298A>C polymorphisms [Put, Weisberg].

Genotypes		677C>T MTHFR		
		CC (%)	CT (%)	TT (%)
1298A>C MTHFR	AA (%)	100	60-70	30-40
	AC (%)	70-80	50-60	-
	CC (%)	50-60	-	-

**Table II.** The frequency of genotypes and alleles of 677C>T MTHFR gene polymorphism.

Polymorphism	Genotypes	Observed value (%)	Expected value (%)
MTHFR 677C>T	CC	335 (50.60)	49.76
	CT	264 (39.88)	41.56
	TT	63 (9.52)	8.68
	Total	662 (100.00)	100.00
	Alleles		
	C	934 (70.54)	
	T	390 (29.46)	
	Total	1324 (100.00)	

**Table III.** The frequency of genotypes and alleles of 1298A>C MTHFR gene polymorphism.

Polymorphism	Genotypes	Observed value (%)	Expected value (%)
MTHFR 1298 A>C	AA	283 (42.75)	44.48
	AC	317 (47.88)	44.43
	CC	62 (9.37)	11.09
	Total	662 (100.00)	100.00
	Alleles		
	A	883 (66.69)	
	C	441 (33.31)	
	Total	1324 (100.00)	

**Table IV.** The frequency of coexistence of 677C>T and 1298A>C polymorphisms.

		677C>T MTHFR			
		CC	CT	TT	Total
1298A>C MTHFR	AA	82 (12.39)	138 (20.85)	63 (9.51)	283 (42.75)
	AC	191 (28.85)	126 (19.03)	0 (0.00)	317 (47.88)
	CC	62 (9.37)	0 (0.00)	0 (0.00)	62 (9.37)
	Total	335 (50.61)	264 (39.88)	63 (9.51)	662 (100.00)

### Genetic analysis

Genetic analysis was conducted with the use of the polymerase chain reaction/ restriction fragments length polymorphism (PCR/RFLP) method. Genomic DNA was extracted from blood leucocytes using QIAamp DNA Blood Mini Kit (QIAGEN Inc., Germany). In order to determine the genotypes of the 677C>T and 1298A>C genetic variants, the previously described methodologies were used [9, 16].

Genomic DNA was amplified with primers (for the 677C>T polymorphism: F5'-TGA AGG AGA AGG TGT CTG CGG GA-3', R5'-AGG ACG GTG CGG TGA GAG TG-3' and for the 1298A>C polymorphism: F 5' CTT CTA CCT gAA gAg

CAA gTC3' and R5' CAT gTC CAC AgC ATg gAg 3'). After amplification, the products were hydrolyzed with a *Hin*I restriction enzyme (EURx, Poland) for the 677C>T polymorphism and with *Mbo*II (EURx, Poland) for the 1298A>C polymorphism.

The analysis of the digested fragments was conducted on 2% agarose gel by electrophoresis. The 677CC genotype was identified in the presence of 198 base pairs (bp) long band, the heterozygous 677CT genotype in the presence of 198, 175, 23 bp bands, and the mutated 677TT genotype – of 175, 23 bp bands. The following genotypes were identified for the 1298A>C polymorphism: AA 176, 30, 28, 22 pz, AC 204, 176, 30, 28, 22 pz, CC 204, 30, 22 pz.

Hubert Wolski et al. Coexistence of the 677C&gt;T and 1298A&gt;C MTHFR polymorphisms and its significance in the population of Polish women.

**Table V.** Co-existence of 677C>T and 1298A>C polymorphisms in different populations.

Author	Population	Study group	n	MTHFR 677C>T/MTHFR 1298A>C								
				N (%)								
				CC/AA	CC/AC	CC/CC	CT/AA	CT/AC	CT/CC	TT/AA	TT/AC	TT/CC
Isotalo i wsp., 2000	Canadian	newborns	119	17 (14.30)	42 (35.30)	9 (7.60)	14 (11.70)	23 (19.30)	-	12 (10.10)	2 (1.70)	-
van der Put i wsp., 1998	Dutch	women/men	403	62 (15.30)	105 (26.10)	38 (9.40)	81 (20.10)	81 (20.10)	-	36 (9.00)	-	-
De Re i wsp., 2010	Italian	women/men	454	24 (5.20)	90 (19.80)	38 (8.40)	98 (21.60)	137 (30.20)	3 (0.70)	57 (12.60)	7 (1.5)	-
		men	315	19 (6.0)	58 (18.40)	23 (7.30)	70 (22.20)	95 (30.20)	1 (0.30)	44 (14.00)	5 (1.60)	-
		women	139	5 (3.60)	32 (23.00)	15 (10.80)	28 (20.10)	42 (30.20)	2 (1.40)	13 (9.30)	2 (1.40)	-
Kumar Rai i wsp., 2005	Hindu	women	159	33 (20.80)	59 (37.10)	22 (13.80)	21 (13.20)	14 (8.80)	4 (2.50)	2 (1.30)	3 (1.90)	1 (0.60)
Rady i wsp., 1999	Ashkenazi/ Caucasian	women/men	186	33 (17.70)	50 (26.90)	20 (10.80)	19 (10.20)	39 (21.00)	-	25 (13.40)	-	-
		women/men	148	12 (8.10)	23 (15.50)	12 (8.10)	29 (19.60)	33 (22.30)	-	39 (26.40)	-	-
Sazci i wsp., 2005	Turkish	women/men	1684	178 (10.60)	401 (23.80)	146 (8.70)	411 (24.40)	363 (21.60)	17 (1.00)	146 (8.70)	22 (13.00)	-
Bodurogulu i wsp., 2005	Turkish	children	93	11 (11.80)	13 (14.00)	7 (7.50)	28 (30.10)	26 (28.00)	-	8 (8.60)	-	-
		fathers	72	3 (4.20)	10 (13.90)	10 (13.90)	28 (38.90)	27 (37.50)	-	2 (2.80)	-	-
		mothers	80	4 (5.00)	17 (21.25)	4 (5.00)	17 (21.25)	23 (28.75)	-	7 (8.75)	-	-
Ergul i wsp., 2003	Turkish	women	193	35 (18.10)	45 (23.30)	14 (7.30)	48 (24.90)	35 (18.10)	4 (2.10)	7 (3.60)	5 (2.60)	-

Frequencies of the genotypes were compared by the chi-square test. The expected genotype frequencies were calculated from allele frequencies with the use of the Hardy-Weinberg equilibrium.

## Results

The frequency of the 677CC, 677CT and 677TT genotypes in the studied population of Polish women was 50.60%, 39.88% and 9.52%, respectively. As for the 1298AA, 1298AC and 1298CC genotypes, the obtained results were as follows: 42.75%, 47.88% and 9.37% (Tables II and III). Simultaneous analysis revealed the most frequent coexistence of the 677CC/1298AC (28.85%), 677CT/1298AA (20.85%) and 677CT/1298AC (19.03%) genotypes. The coexistence of the 677CC/1298AA (12.39%), 677CC/1298CC (9.37%) and 677TT/1298AA (9.51%) genotypes was observed less frequently. To the best of our knowledge, the coexistence of the 677CT/1298CC, 677TT/1298AC and 677TT/1298CC genotypes has been not observed so far in studies on the population of Polish women (Table IV).

## Discussion

Numerous studies have investigated the frequency of the genotypes and alleles of the 677C>T and 1298A>C MTHFR gene polymorphisms in various races and ethnic groups [14, 17, 18, 19, 20].

However, only a few analyses have focused on the coexistence of these genetic variants in different populations, and their correlation with MTHFR activity and folate level changes [21].

One of the first researches in this field was performed by van der Put et al., in the Dutch population. Coexistence of the 677TT/1298AA and 677CT/1298AC genotypes was observed in 9% and 20.1% of the studied population, respectively (50% decrease of MTHFR activity). That study also revealed a significant decrease of the MTHFR activity in case of coexistence of the 677TT/1298AA genotypes (by approximately 70%). Other genotype combinations, such as 677CC/1298CC (9.4% of the population) and 677CT/1298AA (20.1% of the population) correlated with 40% less MTHFR activity. In the studied Dutch population, there was no coexistence of the 677CT/1298CC, 677TT/1298AC and 677TT/1298CC genotypes [11].

Other results concerning the Italian population were obtained by de Re et al., in their study on 315 males and 139 females. These authors observed a higher frequency of the 677CT/1298AC (30.2%) and 677TT/1298AA (12.60%) genotypes, which correlated with decreased MTHFR activity [22].

A significant number of studies on the MTHFR gene polymorphisms were performed in the Turkish population. Sazci et al., performed an analysis involving 1004 women and 680 men. The coexistence of the 677CT/1298AC genotypes was

Hubert Wolski et al. Coexistence of the 677C>T and 1298A>C MTHFR polymorphisms and its significance in the population of Polish women.

noted in 21.6% of the cases [23]. Also, Ergul et al., demonstrated a coexistence of the 677CT/1298AC genotypes in 21.6% of the investigated Turkish population [24]. In a study by Boduroglu et al., the coexistence of the 677CT/1298AC genotypes was observed in 28.75% of the members of the Turkish population (the study group consisted of 245 subjects: 80 mothers, 72 fathers, 93 children) [25].

Another ethnic group which was analyzed was a population of 159 Hindu women. The coexistence of the 677CC/1298AC genotypes was observed in 37.1%, whereas the coexistence of 677TT/1298CC genotypes was significantly less frequent, only in 0.6% of the cases [26].

One of the most interesting investigations on that subject was performed by Rady et al. Their study involved two populations: Ashkenazi and Caucasian population from Texas, USA. In the Ashkenazi population, the most frequent genotype coexistence was 677CC/1298AC and, surprisingly, a much less frequent 677CT/1298AA, while in the population from Texas high frequency of the 677TT/1298AA genotypes was observed (26.4%) [27].

In various studies on many European populations, the coexistence of mutated genotypes (four mutated alleles, 677TT/1298CC), as well as the coexistence of homozygous and heterozygous genotypes (three mutated alleles, 677CT/1298CC and 677TT/1298AC) has been not observed [11], with the exception of the Turkish population, where a coexistence of 677CT/1298CC as well as 677TT/1298AC has been noted [23, 24]. Also, carries of the 677CT/1298CC genotypes were found in the Hindu population (2.5%) [25]. Interestingly, the coexistence of three mutated 677TT/1298AC alleles was also detected in the Canadian population (1.70%) [28]. The results of the abovementioned studies are presented in Table V. These results are consistent with those obtained in our analysis of Polish women. The frequency of the coexistence of the genotypes of the 677C>T and 1298A>C polymorphisms was comparable to those observed in the Dutch population [11]. As in their study, we also did not observe the coexistence of three or four mutated alleles of the investigated MTHFR polymorphisms in the Polish population. Contrary to our results, coexistence of the 677CT/1298AC genotypes was observed most frequently (30.2%) in the Italian population (10% more as compared to the frequency observed in Polish women from the Wielkopolska region). Noteworthy, a coexistence of as many as three mutated alleles of the 677CT/1298CC and 677TT/1298AC polymorphisms in the population from the south of Italy was reported [22].

Our results suggested that environmental factors should be also considered in such analysis. The observed higher frequency of the coexistence of genotypes with a higher number of the mutated alleles suggested a significant role of diet rich in folate in the south of Europe (Italy, Turkey). Higher folate intake could promote survival of individuals with mutated genotypes of the MTHFR polymorphisms. Polymorphism distribution is also influenced by ethnic origin and folic acid demand. Increased seasonal availability of folate-rich food compensates for the negative effects of the 677T allele. Proper folate intake is believed to increase the prevalence of the 677T allele in the general population. High survivability of the 677T embryos might result in an increased number of cardiovascular and neurodegenerative diseases [29].

Some authors also demonstrate the influence of latitude and ultraviolet radiation on the frequency of the 677C>T polymorphism in many populations. Prolonged radiation time is correlated with increased frequency of the mutated alleles of the 677C>T polymorphism. Elongation of the environmental UV radiation time in the periconceptional period decreases folate concentration, because ultraviolet light type A induces oxidative degradation of 5-methylenetetrahydrofolate (5-MTHF) in the skin, which results in decreased 5,10-MTHF cell concentration. In such cases, the 5,10-MTHF is more intensively used for DNA synthesis in 677T-MTHFR embryos than in 677CC embryos. As a result, 677T-MTHFR embryos are in the group with a great chance of survival and the frequency of 677T alleles increases [30].

The frequency of the mutated MTHFR gene variants and coexistence of the MTHFR genotypes are influenced by the interaction of the genetic-environmental natural selection [30,31]. All these facts show that increased folate intake should be recommended in female carriers of both, the mutated genotypes of 677C>T and the 1298A>C polymorphisms (677TT, 1298CC, 677CT/1298AC and 677TT/1298AA). Both these genetic variants are correlated with changes in the MTHFR activity, higher homocysteine, and lower folate concentration.

## Conclusions

1. The frequency of genotype coexistence of the 677C>T and 1298A>C MTHFR gene polymorphisms in the studied population of Polish women is similar to findings about other North European populations.
2. Female carriers of the mutated variants of both, 677C>T and 1298A>C polymorphisms of the MTHFR gene should receive special perinatal care in order to prevent fetal defects and thrombosis-related complications during pregnancy. It is vital to emphasize the significance of proper education about folate supplementation, especially in pregnant patients and women of reproductive age.

## Oświadczenie autorów

1. Hubert Wolski – autor koncepcji i założeń pracy, przygotowanie anuskryptu i piśmiennictwa, ostateczna korekta i akceptacja manuskryptu – autor zgłaszający i odpowiedzialny za manuskrypt.
2. Maria Kocięska – zebranie materiału, analiza statystyczna wyników, przygotowanie manuskryptu.
3. Aleksandra E. Mrozkiewicz – współautor tekstu pracy, korekta i aktualizacja literatury
4. Magdalena Barlik – analiza i interpretacji wyników, współautor tekstu pracy.
5. Grażyna Kurzawińska – współautor tekstu pracy, analiza statystyczna wyników.

## Źródło finansowania:

badania statutowe Kliniki Perinatologii i Chorób Kobięcych UM w Poznaniu – nr: 502-01-02218344-0003344

## Konflikt interesów:

Autorzy nie zgłaszają konfliktu interesów oraz nie otrzymali żadnego wynagrodzenia związanego z powstawaniem pracy

Hubert Wolski et al. Coexistence of the 677C&gt;T and 1298A&gt;C MTHFR polymorphisms and its significance in the population of Polish women.

## References

- Chiang FF, Huang SC, Wang HM, [et al.]. High serum folate might have a potential dual effect on risk of colorectal cancer. *Clin Nutr*. 2014 Nov 8. pii: S0261-5614(14)00264-7. doi: 10.1016/j.clnu.2014.10.011.
- McGarel C, Pentieva K, Strain JJ, McNulty H. Emerging roles for folate and related B-vitamins in brain health across the lifecycle. *Proc Nutr Soc*. 2014, 5, 1-10.
- Catena C, Colussi G, Nait F, [et al.]. Elevated homocysteine levels are associated with the metabolic syndrome and cardiovascular events in hypertensive patients. *Am J Hypertens*. 2014 Dec 12. pii: hpu248.
- Mandaviya PR, Stolk L, Heil SG. Homocysteine and DNA methylation: A review of animal and human literature. *Mol Genet Metab*. 2014, 113, 243-252. doi: 10.1016/j.ymgme.2014.10.006.
- Scazzone C, Bono A, Tornese F, [et al.]. Correlation between low folate levels and hyperhomocysteinemia, but not with vitamin B12 in hypertensive patients. *Ann Clin Lab Sci*. 2014, 44, 286-290.
- Seremak-Mrozikiewicz A. Znaczenie metabolizmu folianów w rozwoju powikłań u kobiet ciężarnych. *Ginekol Pol*. 2013, 84, 377-384.
- Seremak-Mrozikiewicz A. Implikacje kliniczne metabolizmu folianów. *Ginekol Pol*. 2014, 7, 65-71.
- Iacobazzi V, Infantino V, Castegna A, Andria G. Hyperhomocysteinemia: related genetic diseases and congenital defects, abnormal DNA methylation and newborn screening issues. *Mol Genet Metab*. 2014, 113, 27-33. doi: 10.1016/j.ymgme.2014.07.016.
- Frosst P, Blom HJ, Milos R, [et al.]. A candidate genetic risk factor for vascular disease: a common mutation methylenetetrahydrofolate reductase. *Nat Genet*. 1995, 10, 111-113.
- Rozen R. Molecular genetics of Methylenetetrahydrofolate reductase deficiency. *J Inher Metab Dis*. 1996, 19, 589-594.
- van der Put NM, Gabreëls F, Stevens EM, [et al.]. A second common mutation in the methylenetetrahydrofolate reductase Gene: An Additional Risk Factor for Neural-Tube Defects? *Am J Hum Genet*. 1998, 62, 1044-1051.
- Hughes LB, Beasley TM, Patel H, [et al.]. Racial or ethnic differences in allele frequencies of single-nucleotide polymorphisms in the methylenetetrahydrofolate reductase gene and their influence on response to methotrexate in rheumatoid arthritis. *Ann Rheum Dis*. 2006, 65, 1213-1218.
- Robien K, Ulrich CM. 5,10-methylenetetrahydrofolate reductase polymorphisms and leukemia risk: a HuGE minireview. *Am J Epidemiol* 2003, 157, 571-582.
- Binia A, Contreras AV, Canizales-Quinteiros S, [et al.]. Geographical and ethnic distribution of single nucleotide polymorphism within genes of the folate/homocysteine pathway metabolism. *Genes Nutr*. 2014, 9, 421. doi: 10.1007/s12263-014-0421-7.
- Weisberg I, Tran P, Chritensen B, [et al.]. A second genetic polymorphism in methylenetetrahydrofolate reductase (MTHFR) associated with decreased enzyme activity. *Mol Genet Metab*. 1998, 64, 169-172.
- Hanson NQ, Aras O, Yang F, Tsai MY. C677T and A1298C polymorphisms of the methylenetetrahydrofolate reductase gene: incidence and effect of combined genotypes on plasma fasting and postmethionine load homocysteine in vascular disease. *Clin Chem*. 2001, 47, 661-666.
- Zappacosta B, Graziano M, Persichilli S, [et al.]. 5,10-Methylenetetrahydrofolate reductase (MTHFR) C677T and A1298C polymorphisms: genotype frequency and association with homocysteine and folate levels in middle-southern Italian adults. *Cell Biochem Funct*. 2014, 32, 1-4.
- Yang B, Liu Y, Li Y, [et al.]. Geographical distribution of MTHFR C677T, A1298C and MTRR A66G gene polymorphisms in China: findings from 15357 adults of Han nationality. *PLoS One*. 2013;8(3):e57917. doi: 10.1371/journal.pone.0057917.
- Antonio-Véjar V, Del Moral-Hernández O, Alarcón-Romero LC, [et al.]. Ethnic variation of the C677T and A1298C polymorphisms in the methylenetetrahydrofolate-reductase (MTHFR) gene in southwestern Mexico. *Genet Mol Res*. 2014, 13, 7950-7957.
- Ozen F, Sen M, Ozdemir O. Methylenetetrahydrofolate reductase gene germ-line C677T and A1298C SNPs are associated with colorectal cancer risk in the Turkish population. *Asian Pac J Cancer Prev*. 2014, 15, 7731-7735.
- Guéant-Rodriguez RM, Guéant JL, Debard R, [et al.]. Prevalence of methylenetetrahydrofolate reductase 677T and 1298C alleles and folate status: a comparative study in Mexican, West African, and European populations. *Am J Clin Nutr*. 2006, 83, 701-707.
- de Re V, Cannizzaro R, Canzonieri V, [et al.]. MTHFR polymorphisms in gastric cancer and in first-degree relatives of patients with gastric cancer. *Tumor Biol*. 2010, 31, 23-32.
- Sazci A, Ergul E, Kaya G, Kara I. Genotype and allele frequencies of the polymorphic methylenetetrahydrofolate reductase gene in Turkey. *Cell Biochem Funct*. 2005, 23, 51-54.
- Ergul E, Sazci A, Utkan N, Canturk Z. Polymorphisms in the MTHFR gene are associated with breast cancer. *Tumour Biol*. 2003, 24, 286-290.
- Boduroğlu K, Alanay Y, Aikaşifoğlu M, [et al.]. Analysis of MTHFR 1298A>C in addition to MTHFR 677C>T polymorphism as a risk factor for neural tube defects in the Turkish population. *Turk J Pediatr*. 2005, 47, 327-333.
- Kumar Rai A, Singh S, Mehta S, [et al.]. MTHFR C677T and A1298C polymorphisms are risk factors for Down's syndrome in Indian mothers. *J Hum Genet*. 2006, 51, 278-283.
- Rady P, Tyring S, Hudnall D, [et al.]. Methylenetetrahydrofolate reductase (MTHFR): the incidence of mutations C677T and A1298C in the Ashkenazi Jewish population. *Am J Med Genet*. 1999, 86, 380-384.
- Isotalo PA, Wells GA, Donnelly JG. Neonatal and Fetal Methylenetetrahydrofolate Reductase Genetic Polymorphisms: An Examination of C677T and A1298C Mutations. *Am J Hum Genet*. 2000, 67, 986-990.
- Lucock M, Yates Z. Folic acid - vitamin and panacea or genetic time bomb? *Nat Rev Genet*. 2005, 6, 235-240.
- Lucock M, Gianville T, Ovadia L, [et al.]. Photoperiod at conception predicts C677T-MTHFR genotype: A novel gene-environment interaction. *Am J Hum Biol*. 2010, 22, 484-489. doi: 10.1002/ajhb.21022.
- Yafei W, Lijun P, Jinfeng W, Xiaoying Z. Is the prevalence of MTHFR C677T polymorphism associated with ultraviolet radiation in Eurasia? *J Hum Genet*. 2012, 57, 780-786.