

# The impact of preoperative anxiety on pain and analgesia consumption in women undergoing vaginal hysterectomy with general anesthesia and spinal anesthesia

Hasan Berkan Sayal<sup>1</sup> , Hasan Ali İnal<sup>1</sup> , Zeynep Ozturk Inal<sup>2</sup> 

<sup>1</sup>Antalya Training and Reseach Hospital, Health Science University, Varlık Mah. Kazım Karabekir Cd. Muratpasa, Antalya, Türkiye

<sup>2</sup>Department of Obstetrics and Gynecology, Konya Training and Research Hospital, Türkiye

## ABSTRACT

**Objectives:** To assess the impact of preoperative anxiety on pain and analgesic consumption in patients undergoing vaginal hysterectomy (VH) with general and spinal anesthesia.

**Material and methods:** A total of 200 participants, including 100 undergoing vaginal hysterectomy with general anesthesia (group 1) and 100 with spinal anesthesia (group 2), were enrolled. A visual analog scale (VAS) was used for the postoperative pain intensity.

**Results:** The 1<sup>st</sup> hour, 6<sup>th</sup> hour, 12<sup>th</sup> hour, and 18<sup>th</sup> hour VAS scores were higher in vaginal hysterectomy with general anesthesia than in vaginal hysterectomy with spinal anesthesia.

**Conclusions:** Although participants undergoing VH with spinal anesthesia (preoperative state anxiety inventory score > 45) had lower pain intensity scores in the first 18 hours compared to those undergoing VH with general anesthesia, their postoperative analgesic requirements were similar.

**Keywords:** anxiety; general anesthesia; pain; spinal anesthesia; vaginal hysterectomy

Ginekologia Polska 2024; 95, 8: 601–606

## INTRODUCTION

Today, minimal invasive surgical methods are preferred for hysterectomy, which is the most common gynecologic surgical operation performed worldwide [1, 2]. Vaginal hysterectomy (VH), one of these methods, is widely performed, especially in indications such as uterovaginal prolapse, uterine leiomyoma, and abnormal uterine bleeding [1, 3]. However, the size and accessibility of the uterus, additional gynecologic surgery, the availability of appropriate surgical equipment, and the technological infrastructure of the hospital are important in the choice of this method [1, 4].

Unfortunately, postoperative pain after surgical interventions remains an important problem despite modern anesthesia techniques and medication [5, 6]. The phenomenon of postoperative pain, which varies from person to

person, is directly affected by preoperative anxiety [5]. If this preoperative anxiety can be minimized with pharmacologic and other alternative medicine methods, postoperative pain will also be reduced [5, 7]. As a result, patients with reduced postoperative pain will be able to mobilize earlier and the risk for thromboembolic events will be reduced because hypercoagulability will be reduced [6, 8].

There are contradictory results related to preoperative anxiety and postoperative pain in the literature; some studies show that preoperative anxiety increases postoperative pain and analgesic requirement [7, 9], and other studies [5, 10] claimed that there was no relationship. This study aimed to assess whether preoperative anxiety has any effect on postoperative pain and analgesic consumption in participants undergoing VH.

### Corresponding author:

Hasan Berkan Sayal

Antalya Training and Reseach Hospital, Health Science University, Aarlık, Kazım Karabekir Cd. Muratpasa, 07100 Antalya, Türkiye

e-mail: drberkansayal@yahoo.com

Received: 16.03.2023 Accepted: 19.07.2023 Early publication date: 15.04.2024

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

## MATERIAL AND METHODS

### Setting

The study was approved by institutional review board (reference number: 2016/10-12) and a total of 200 VHs performed were included.

### Study population

The inclusion criteria were benign uterine leiomyoma, menometrorrhagia resistant to medical treatment, and uterovaginal prolapse. The exclusion criteria were as follows: having any chronic illness (e.g. diabetes mellitus, systemic vascular disease, cardiac and pulmonary disease), any malignant tumor, any mental disorder, mental retardation or cognitive disabilities, a chronic pain history, and drug and alcohol abuse.

### Study design

A total of 200 participants (aged 45–65 years) requiring VH were divided consecutively into two groups: 100 patients underwent general anesthesia (group 1) and 100 underwent spinal anesthesia (group 2). Two senior gynecologists and two anesthesiologists performed all the procedures. The age, parity, indications, surgical time, pre and postoperative serum hemoglobin (Hb) levels, blood loss, hospital stay, analgesic needs, living area, and educational level, were recorded and compared.

### Preoperative anxiety assessment

Three independent questionnaires [State Anxiety Inventory (SAI), Trait Anxiety Inventory (TAI) [11], and Somatosensory Amplification Scale (SASS) [12]] were completed 24 hours before surgery by each patient.

State Anxiety Inventory and TAI, which are used to measure the patient's anxiety level, consist of 20 items graded from 1 (not all) to 4 (very much). The current level of anxiety is measured using the SAI, and whether a person is generally anxious is measured using the TAI. It has been shown that both measurement scales are useful in evaluating anxiety [13, 14], and a score of > 45/80 is an indicator of high anxiety [14].

Barsky et al. [12] developed SASS and was translated into Turkish and validated by Gulec and Sayar [15]. Somatosensory amplification, which is important in clinical situations, is disproportionately correlated with somatic symptoms in significant organ pathologies [12].

### Pre-surgery preparation

Patients scheduled for VH were hospitalized one day before surgery and each underwent simultaneous abdominal and transvaginal ultrasonography to determine the uterine size, along with a standard preoperative evaluation

(laboratory test and premedication) required by anesthesia. A ward nurse administered cefazolin 2 g via intravenous routes for prophylaxis to all participants, a bladder catheter was placed before the surgery, and it was withdrawn by the ward nurse after 6-8 hours when the patient was mobilized. Low-molecular-weight heparin was administered for antithrombotic prophylaxis.

### Anesthesia protocol

Anesthesia type selection was performed by the anesthesiologist and was terminated when the target number for each group was reached. An intravenous line was opened, preferably in the left arm, and 500 mL of Ringer's lactate solution was given as a liquid bolus and the patient's vital signs were monitored before the surgery.

After the surgery was completed, the patients were taken to the postoperative care unit to monitor their vital signs and were transferred to the ward within 1 hour.

### Surgical procedure for VH


Vaginal hysterectomy was performed in the Trendelenburg position. After disinfection and sterile closure, the portio cervix was held using by two forceps. An annular incision was performed in the cervix, the bladder was separated, and the vesico-uterinum spatium was opened. Then, both the ligg. Sacrouterinae and Cardinale were grasped, cut, and ligated, and then the parametria were separated. The annexes were ligated separately. The uterus was removed, and finally, the peritoneum was closed.

### Assessment of pain perception and analgesic needs

A VAS was used to assess pain perception. The patients marked their pain on the VAS (0 = no pain and 10 = worst pain).

### Statistical analysis

All the variables were analyzed using the SPSS 15.0 software (Statistical Package for the Social Sciences, SPSS Inc., Chicago, IL, USA). Kolmogorov-Smirnov tests were used to determine normal or non-normal distribution for the continuous variables. Independent samples t-tests or Mann-Whitney U test were used for continuous variables. Mean  $\pm$  standard deviation (SD) was preferred for normally distributed variables, and those not distributed normally were presented as median and 25<sup>th</sup> and 75<sup>th</sup> ranges. Evaluation of the effect of VAS scores on the rates of patients undergoing VH was performed using logistic regression analysis. The categorical data was assessed by Chi-square test or Fisher's exact test was used for. A p value of 0.05 was accepted as statistical significance.



**Figure 1.** Enrollment and follow-up of the study subjects

A minimum of 100 participants was necessary to show a 10% difference at  $\alpha = 0.05$  and  $\beta = 0.20$  [16] (Research Sample Size Calculation Program). This difference of 10% was predicted both from a pilot study and from our clinical trials.

## RESULTS

A total of 200 participants were categorized as general anesthesia ( $n=100$ ) and spinal anesthesia ( $n = 100$ ) (Fig. 1).

Table 1 summarizes the sociodemographic characteristics of the study participants. Age, body mass index (BMI), parity, indications, surgical time, pre and postoperative serum Hb levels, blood loss, hospital stay, analgesic needs, levels of serum urea, alanine transaminase (ALT), aspartate aminotransferase (AST), and cholinesterase, living area, and educational level were comparable ( $p > 0.05$ ).

Table 2 lists anxiety scores, pain intensity measurements, analgesic consumptions, and VAS scores of the study participants. Although significant differences were obtained in the postoperative pain levels between the groups at the 1<sup>st</sup> hour, 6<sup>th</sup> hour, 12<sup>th</sup> hour, and 18<sup>th</sup> hour, there was no differences in the 24<sup>th</sup>-hour measurements ( $p > 0.05$ ).

Table 3 describes analgesic consumption and pain intensity based on SAI Index scores of the participants. The VAS scores at the 1<sup>st</sup> hour, 6<sup>th</sup> hour, 12<sup>th</sup> hour, and 18<sup>th</sup> hour

were difference between the groups with SAI scores  $\geq 45$ , but no difference was seen in the 24<sup>th</sup>-hour scores.

No difference was noted between the groups in surgical procedures performed concomitantly with VHs (oophorectomy 86.0% vs 81.0%,  $p = 0.446$ ; anterior colporaphy 67.0% vs 72.0%,  $p = 0.539$ , posterior colporaphy 69.0% vs 63.0%,  $p = 0.456$ , paravaginal repair 7.0% vs 11.0%,  $p = 0.522$ , vaginal vault suspension 29.0% vs 24.0%,  $p = 0.522$ , and anal sphincteroplasty 7.0% vs 4.0%,  $p = 0.537$ ).

## DISCUSSION

The pain severity scores in participants undergoing VH with general anesthesia were higher in the first 18 hours postoperatively compared to those undergoing VH with spinal anesthesia. It has been shown that preoperative anxiety scores before cesarean section are associated with higher postoperative pain severity and intensity in patients who delivered by cesarean section under general anesthesia compared with patients who delivered with spinal anesthesia, especially in the first 12 hours [6].

Although abdominal hysterectomy (AH) has been a widely used gynecologic surgery for many years, nowadays, VH, which has a shorter surgical time and is more cost-effective, is preferred especially in the presence of a small uterus, a history of vaginal delivery, and the absence of any adnexal pathology [2]. Therefore, the type of anesthesia used in postoperative pain management is important in increasing VH procedures. Although there is no clear consensus in the literature on the type of anesthesia that should be administered to patients who will undergo VH surgery, spinal anesthesia provides many advantages such as the continuation of spontaneous breathing during surgery, the patient's awakening, and the preservation of protective reflexes such as coughing, as well as early mobilization, minimal lung complications, the continuation of analgesia, and a short hospital stay in the postoperative days [6].

Pinto et al. [10] stipulated that preoperative anxiety did not affect postoperative pain and analgesic consumption in their study, which included 185 patients who underwent AH for benign indications. On the contrary, another study [7] reported that preoperative anxiety scores were directly positively related to analgesic consumption after AH in which 60 patients were evaluated. Additionally, it has been claimed that preoperative anxiety increases analgesic consumption in the acute period and is associated with pain even at 4 months postoperatively.

Kain et al. [5] found that although preoperative trait anxiety had no direct effect on postoperative pain in patients who underwent AH surgery, state anxiety was a direct positive predictor. Similarly, Aouad et al. [7] reported that only preoperative state anxiety affected postoperative pain

	<b>Group 1 (n = 100)</b>	<b>Group 2 (n = 100)</b>	<b>p value</b>
<b>Age [years]</b>	56.32 ± 7.20	57.44 ± 6.28	0.174
<b>BMI [kg/m<sup>2</sup>]</b>	25.63 ± 2.29	25.94 ± 2.51	0.308
<b>Parity</b>	3.0 (2.0–4.0)	3.0 (2.0–3.0)	0.374
<b>Indications [(%)]</b>			
Meno-metroragia resistant to medical treatment	55	49%	0.548
Cronic pelvic pain	8	12%	
Uterovaginal prolapse	37	39%	
<b>Duration of operation [min]</b>	73.40 ± 10.16	71.36 ± 11.02	0.189
<b>Hb (preoperative) [gr/L]</b>	11.79 ± 1.86	12.04 ± 1.76	0.346
<b>Hb (postoperative) [gr/L]</b>	10.29 ± 1.82	10.54 ± 1.76	0.315
<b>Blood loss [mL]</b>	232.68 ± 83.09	216.61 ± 58.27	0.115
<b>Length of hospital stay [days]</b>	2.69 ± 0.77	2.52 ± 0.69	0.103
<b>Analgesic needs [days]</b>	3.88 ± 1.11	3.63 ± 1.07	0.106
<b>Living area [%]</b>			
Village	11	17	0.369
Town	14	10	
City	75	73	
<b>Educational level [%]</b>			
Primary school	55	54	0.176
Secondary school	22	32	
University	23	14	

BMI — body mass index; Hb — hemoglobin

	<b>Group 1 (n = 100)</b>	<b>Group 2 (n = 100)</b>	<b>Odds ratio</b>	<b>95% confidence interval</b>	<b>p value</b>
<b>MSAI score</b>	40.03 ± 8.82	38.57 ± 9.14	–	–	0.253
<b>MTAI score</b>	43.97 ± 7.11	42.89 ± 7.21	–	–	0.149
<b>SAS score</b>	31.47 ± 6.77	29.96 ± 7.03	–	–	0.125
<b>The dosage of diclofenac consumption [mg]</b>	108.76 ± 51.54	117.74 ± 44.29	–	–	0.186
<b>The dosage of pethidine consumption [mg]</b>	23.01 ± 15.12	20.51 ± 15.61	–	–	0.619
<b>SAI score ≥ 45 [%]</b>	33	31	–	–	0.416
<b>VAS</b>					
1. hour	4.12 ± 1.80	3.16 ± 1.94	0.766	0.658–0.895	< 0.001*
6. hour	3.78 ± 1.96	2.82 ± 1.54	0.707	0.592–0.844	< 0.001*
12. hour	3.58 ± 2.05	2.54 ± 1.65	0.736	0.628–0.862	< 0.001*
18. hour	3.02 ± 1.56	2.47 ± 1.59	0.798	0.662–0.962	0.015*
24. hour	2.19 ± 1.07	2.32 ± 1.43	1.191	0.933–1.521	0.361
<b>MSAI score</b>	40.03 ± 8.82	38.57 ± 9.14	–	–	0.252

\*Statistically significant; MSAI — Mean State Anxiety Inventory; MTAI — Mean Trait Anxiety Inventory; SAS — Somato-sensory amplification Scale; SAI — State Anxiety Inventory

**Table 3.** Analgesic consumption and pain intensity of the participants with and without State Anxiety Index scores of  $\geq 45$ 

	SAI Score < 45			SAI Score $\geq 45$		
	General Anesthesia (Group 1) (n = 68)	Spinal Anesthesia (Group 2) (n = 70)	p value	General Anesthesia (Group 1) (n = 32)	Spinal Anesthesia (Group 2) (n = 30)	p value
The dosage of diclofenac consumption [mg]	118.02 $\pm$ 43.58	110.37 $\pm$ 43.72	0.304	117.18 $\pm$ 46.42	105.01 $\pm$ 67.07	0.408
The dosage of pethidine consumption [mg]	22.15 $\pm$ 16.75	16.92 $\pm$ 11.88	0.375	28.14 $\pm$ 12.02	25.01 $\pm$ 11.47	0.743
<b>VAS</b>						
1. hour	4.02 $\pm$ 1.81	3.63 $\pm$ 2.15	0.162	4.37 $\pm$ 1.81	3.05 $\pm$ 2.14	< 0.001*
6. hour	3.65 $\pm$ 2.02	3.18 $\pm$ 1.46	0.102	4.14 $\pm$ 1.78	2.79 $\pm$ 1.37	< 0.001*
12. hour	3.61 $\pm$ 2.04	3.06 $\pm$ 1.86	0.217	3.58 $\pm$ 2.08	2.48 $\pm$ 1.82	< 0.001*
18. hour	3.02 $\pm$ 1.47	2.83 $\pm$ 1.38	0.358	2.96 $\pm$ 1.72	2.02 $\pm$ 1.03	< 0.001*
24. hour	2.32 $\pm$ 1.89	2.07 $\pm$ 1.06	0.413	2.16 $\pm$ 1.94	1.88 $\pm$ 0.94	0.218

\*Statistically significant; SAI — State Anxiety Inventory; VAS — Visual analog scale

band analgesic consumption in AH, whereas trait anxiety had no such effect. On the other hand, Carvalho et al. [17] claimed that there was no correlation between anxiety scores and postoperative pain scores in hysterectomy surgery. It has been observed that psychosocial education and support before hysterectomy reduces preoperative anxiety and therefore this condition reduces postoperative pain severity and intensity [9].

Abdominal hysterectomy performed with a Pfannenstiel's incision in the abdomen may result in higher postoperative pain scores than VH [5, 7, 18]. Especially in open abdominal surgeries, the prolongation of surgical time causes extended pain stimulation and this causes an increase in postoperative pain scores [19]. Postoperative pain scores are lower in VH because it has a shorter surgical time and is less invasive than AH. Unfortunately, we did not compare the severity and intensity of preoperative anxiety and postoperative pain between AH and VH in our study.

The conflicting results between preoperative anxiety and postoperative pain severity and intensity in the literature may be due to the inhomogeneity of the groups in the studies, differences in surgical procedures, non-standardized anesthesia methods, different anesthetic agents, non-standardized pain scale scales, and insufficient sample sizes [5, 7, 10]. The possible confounding factors such as age, BMI, parity, surgical time, and educational level were similar. Additionally, VH was performed by specialist physicians experienced in surgical interventions and anesthesia procedures.

In experimental pain models, it has been observed that invasive procedures such as intravenous catheter insertion

before surgery increase the amount and duration of postoperative analgesic consumption [20, 21]. This situation was not evaluated in our study.

Control of postoperative pain is very important, especially in the first 24 hours after surgical interventions, because postoperative pain is more intense and analgesic consumption is at a maximum level in this period [6, 22]. If pain control cannot be achieved within this period, the first mobilization period of the patient will be later in the postoperative period, which will increase the susceptibility to thromboembolic events and will also create a risk factor for chronic pelvic pain [7, 8].

There are no data about postoperative pain severity and intensity, and also analgesic requirements after VH procedures. Generally, the data in the literature belong to cesarean section and AH procedures as mentioned above. To our knowledge, our study is the first in this topic.

The limitation was the use of a subjective method instead of an objective method in the assessment of anxiety and pain intensity.

## CONCLUSIONS

In conclusion, although participants undergoing VH with spinal anesthesia (Preoperative SAI score  $> 45$ ) had lower pain intensity scores in the first 18 hours compared to those undergoing VH with general anesthesia, their postoperative analgesic requirements were similar. Preoperative anxiety should be alleviated with psychiatric support to reduce the severity and intensity of postoperative pain. Further randomized controlled studies with larger participant numbers are necessary to support our findings.

## Article information and declarations

### Funding

None.

### Acknowledgements

None.

### Conflict of interest

The authors declared that they have no conflict of interest.

## REFERENCES

- Committee Opinion No 701: Choosing the Route of Hysterectomy for Benign Disease. *Obstetrics & Gynecology*. 2017; 129(6): e155–e159, doi: [10.1097/aog.0000000000002112](https://doi.org/10.1097/aog.0000000000002112).
- Inal ZO, Inal HA. Comparison of abdominal, vaginal, and laparoscopic hysterectomies in a tertiary care hospital in Turkey. *Ir J Med Sci*. 2018; 187(2): 485–491, doi: [10.1007/s11845-017-1660-6](https://doi.org/10.1007/s11845-017-1660-6), indexed in PubMed: [28726032](https://pubmed.ncbi.nlm.nih.gov/28726032/).
- Inal ZO, Inal HA, Kucukosmanoglu I, et al. Assessment of Endometrial Sampling and Histopathological Results: Analysis of 4,247 Cases. *Eurasian J Med*. 2017; 49(1): 44–47, doi: [10.5152/eurasianjmed.2017.16269](https://doi.org/10.5152/eurasianjmed.2017.16269), indexed in PubMed: [28416932](https://pubmed.ncbi.nlm.nih.gov/28416932/).
- Inal HA, Kaplan PB, Usta U, et al. Neuromuscular morphometry of the vaginal wall in women with anterior vaginal wall prolapse. *NeuroUrol Urodyn*. 2010; 29(3): 458–463, doi: [10.1002/nau.20779](https://doi.org/10.1002/nau.20779), indexed in PubMed: [19714736](https://pubmed.ncbi.nlm.nih.gov/19714736/).
- Kain ZN, Sevarino F, Alexander GM, et al. Preoperative anxiety and postoperative pain in women undergoing hysterectomy. A repeated-measures design. *J Psychosom Res*. 2000; 49(6): 417–422, doi: [10.1016/s0022-3999\(00\)00189-6](https://doi.org/10.1016/s0022-3999(00)00189-6), indexed in PubMed: [11182434](https://pubmed.ncbi.nlm.nih.gov/11182434/).
- Ozturk Inal Z, Gorkem U, Inal HA. Effects of preoperative anxiety on postcesarean delivery pain and analgesic consumption: general versus spinal anesthesia. *J Matern Fetal Neonatal Med*. 2020; 33(2): 191–197, doi: [10.1080/14767058.2018.1487948](https://doi.org/10.1080/14767058.2018.1487948), indexed in PubMed: [29886798](https://pubmed.ncbi.nlm.nih.gov/29886798/).
- Aouad MT, Kanazi GE, Malek K, et al. Predictors of postoperative pain and analgesic requirements following abdominal hysterectomy: an observational study. *J Anesth*. 2016; 30(1): 72–79, doi: [10.1007/s00540-015-2090-0](https://doi.org/10.1007/s00540-015-2090-0), indexed in PubMed: [26499321](https://pubmed.ncbi.nlm.nih.gov/26499321/).
- Perkins FM, Kehlet H. Chronic pain as an outcome of surgery. A review of predictive factors. *Anesthesiology*. 2000; 93(4): 1123–1133, doi: [10.1097/0000542-200010000-00038](https://doi.org/10.1097/0000542-200010000-00038), indexed in PubMed: [11020770](https://pubmed.ncbi.nlm.nih.gov/11020770/).
- Erdoğan E, Demir S, Çalışkan BB, et al. Effect of psychological care given to the women who underwent hysterectomy before and after the surgery on depressive symptoms, anxiety and the body image levels. *J Obstet Gynaecol*. 2020; 40(7): 981–987, doi: [10.1080/01443615.2019.1678574](https://doi.org/10.1080/01443615.2019.1678574), indexed in PubMed: [31795799](https://pubmed.ncbi.nlm.nih.gov/31795799/).
- Pinto PR, McIntyre T, Fonseca C, et al. Pre- and post-surgical factors that predict the provision of rescue analgesia following hysterectomy. *Eur J Pain*. 2013; 17(3): 423–433, doi: [10.1002/j.1532-2149.2012.00205.x](https://doi.org/10.1002/j.1532-2149.2012.00205.x), indexed in PubMed: [22893516](https://pubmed.ncbi.nlm.nih.gov/22893516/).
- Spielberger CD, Gorsuch RL, Lushene RE. *State-Trait Anxiety inventory for adults*. Palo Alto, CA: Mind Garden 1983.
- Barsky AJ, Wyshak G, Klerman GL. The somatosensory amplification scale and its relationship to hypochondriasis. *J Psychiatr Res*. 1990; 24(4): 323–334, doi: [10.1016/0022-3956\(90\)90004-a](https://doi.org/10.1016/0022-3956(90)90004-a), indexed in PubMed: [2090830](https://pubmed.ncbi.nlm.nih.gov/2090830/).
- Orbach-Zinger S, Ginosar Y, Elliston J, et al. Influence of preoperative anxiety on hypotension after spinal anaesthesia in women undergoing Caesarean delivery. *Br J Anaesth*. 2012; 109(6): 943–949, doi: [10.1093/bja/aes313](https://doi.org/10.1093/bja/aes313), indexed in PubMed: [22964265](https://pubmed.ncbi.nlm.nih.gov/22964265/).
- Kuo SY, Chen SR, Tzeng YL. Depression and anxiety trajectories among women who undergo an elective cesarean section. *PLoS One*. 2014; 9(11): e86653, doi: [10.1371/journal.pone.0086653](https://doi.org/10.1371/journal.pone.0086653), indexed in PubMed: [24466190](https://pubmed.ncbi.nlm.nih.gov/24466190/).
- Güleç H, Sayar K. Reliability and validity of the Turkish form of the Somatosensory Amplification Scale. *Psychiatry Clin Neurosci*. 2007; 61(1): 25–30, doi: [10.1111/j.1440-1819.2007.01606.x](https://doi.org/10.1111/j.1440-1819.2007.01606.x), indexed in PubMed: [17239035](https://pubmed.ncbi.nlm.nih.gov/17239035/).
- DSS Research. *Researcher's toolkit*. <http://www.dssresearch.com/toolkit/sscalc>.
- Carvalho B, Zheng M, Aiono-Le Tagaloa L. A prospective observational study evaluating the ability of prelabor psychological tests to predict labor pain, epidural analgesic consumption, and maternal satisfaction. *Anesth Analg*. 2014; 119(3): 632–640, doi: [10.1213/ANE.0000000000000357](https://doi.org/10.1213/ANE.0000000000000357), indexed in PubMed: [25029661](https://pubmed.ncbi.nlm.nih.gov/25029661/).
- Johnson N, Barlow D, Lethaby A, et al. Methods of hysterectomy: systematic review and meta-analysis of randomised controlled trials. *BMJ*. 2005; 330(7506): 1478, doi: [10.1136/bmj.330.7506.1478](https://doi.org/10.1136/bmj.330.7506.1478), indexed in PubMed: [15976422](https://pubmed.ncbi.nlm.nih.gov/15976422/).
- Latremliere A, Woolf CJ. Central sensitization: a generator of pain hypersensitivity by central neural plasticity. *J Pain*. 2009; 10(9): 895–926, doi: [10.1016/j.jpain.2009.06.012](https://doi.org/10.1016/j.jpain.2009.06.012), indexed in PubMed: [19712899](https://pubmed.ncbi.nlm.nih.gov/19712899/).
- Werner MU, Mjöbo HN, Nielsen PR, et al. Prediction of postoperative pain: a systematic review of predictive experimental pain studies. *Anesthesiology*. 2010; 112(6): 1494–1502, doi: [10.1097/ALN.0b013e3181dcd5a0](https://doi.org/10.1097/ALN.0b013e3181dcd5a0), indexed in PubMed: [20460988](https://pubmed.ncbi.nlm.nih.gov/20460988/).
- Janssen KJM, Kalkman CJ, Grobbee DE, et al. The risk of severe postoperative pain: modification and validation of a clinical prediction rule. *Anesth Analg*. 2008; 107(4): 1330–1339, doi: [10.1213/ane.0b013e31818227da](https://doi.org/10.1213/ane.0b013e31818227da), indexed in PubMed: [18806049](https://pubmed.ncbi.nlm.nih.gov/18806049/).
- Ranta PO, Ala-Kokko TI, Kukkonen JE, et al. Incisional and epidural analgesia after caesarean delivery: a prospective, placebo-controlled, randomised clinical study. *Int J Obstet Anesth*. 2006; 15(3): 189–194, doi: [10.1016/j.ijoa.2006.02.003](https://doi.org/10.1016/j.ijoa.2006.02.003), indexed in PubMed: [16798442](https://pubmed.ncbi.nlm.nih.gov/16798442/).