



# Prenatal diagnosis of isolated total anomalous pulmonary venous connection (TAPVC) to coronary sinus

Anna Wojtowicz<sup>1</sup>, Beata Zaluska-Pitak<sup>2</sup>, Magdalena Juszcak<sup>3</sup>,  
Hubert Huras<sup>1</sup>, Sebastian Goreczny<sup>2</sup>

<sup>1</sup>Department of Obstetrics and Perinatology, Jagiellonian University Medical College, Cracow, Poland

<sup>2</sup>Department of Pediatric Cardiology, University Children's Hospital, Faculty of Medicine, Jagiellonian University Medical College, Cracow, Poland

<sup>3</sup>Department of Gynecology and Obstetrics with Gynecology Oncology, L. Rydygier Hospital, Cracow, Poland

Total anomalous pulmonary venous connection (TAPVC) is the fifth most common cyanotic heart disease, affecting 5.9–7.1 per 100,000 live births [1, 2].

The main feature is that all four pulmonary veins (PV) are not connected to the left atrium (LA) but to the systemic veins, coronary sinus (CS), or right atrium (RA). Four TAPVC types are distinguished based on the PV connect sites: supracardiac, cardiac, infracardiac, and a mixed type. TAPVC can occur as part of complex cardiac malformations, especially in heterotaxy syndrome, or in isolation. Prenatal assessment of the fetal venous system and left venoatrial junction is part of the four-chamber-view evaluation, as stipulated in national and international fetal cardiac screening guidelines [3, 4], however assessment is challenging, and reports indicate that only 1.9% of TAPVC cases are diagnosed prenatally [5].

Prenatal diagnosis of TAPVC is crucial, as it explains postnatal cyanotic heart defect symptoms and enables prompt, sometime lifesaving, treatment planning. The presence of an obstruction at any level of the venous route is the most critical factor affecting outcome and management of the defect, making the defect one of the few true neonatal cardiac emergencies. Prognosis is very good if detected prenatally and corrected during the neonatal period [2].

The study aim was to identify characteristic sonographic features of TAPVC where there was PV drainage into the CS.

A 35-year-old primigravida at 26 weeks of pregnancy was referred to our department because PV identification by imaging was impossible. Examination revealed normal visceral situs, normal heart size, drainage of the systemic veins into the RA, and concordant atrioventricular connection. The space between the spine and the LA drew our attention as the four PV drained through a collector to the enlarged CS and then to the RA (Fig. 1A, Suppl. video). Imaging also identified the enlarged width of the CS (Fig. 1B). In colour and PW Doppler modes, the blood flow spectrum was normal (Fig. 1C). The outflow tracts and upper mediastinum were normal. During the third trimester, the RA and RV became enlarged, and the LA and LV volumes were smaller than normal, however the LV depth was normal and formed the apex of the heart. Trivial tricuspid regurgitation was also detected. Coronary sinus width was 4.9 mm. With TAPVC suspected, examination by a pediatric cardiologist confirmed the diagnosis. The neonate was born at 39 weeks of gestation with vitals of body weight 3300 g, Apgar score 10, venous blood pH = 7.321, and satO<sub>2</sub> = 90%.

Postnatal echocardiography and cardiac angiography confirmed that all PV formed a collector behind the LA, and the collector drained into the CS with its kinking (Fig. 1D, Fig. S1). At the level of the CS kinking, a flow gradient of max 16 and mean 12 mm Hg was revealed. The right atrium and ventricle were significantly enlarged (Fig. 1E, Fig. S2), with severe hypertrophy of the right ventricle free wall but normal contractility. Trivial tricuspid valve insufficiency with a retrograde systolic maximal gradient of 34–43 mm Hg, normal right ventricle outflow tract, and widened right and left pulmonary arteries (RPA = 5.2 mm, LPA = 4.7 mm) were observed. Pulmonary blood flow with hypertension symptoms was diagnosed. Atrial septal defect type II of 6 mm with right to left shunt was detected. Left ventricle output was 2.2 L/min/m<sup>2</sup>, with normal ejection 75% and shortening fraction 40%. The mitral valve was competent, and the aortic

Corresponding author:


Anna Wojtowicz

Department of Obstetrics and Perinatology, Jagiellonian University Medical College, Kopernika 23 St., 31–501 Cracow, Poland

e-mail: anna.3.wojtowicz@uj.edu.pl

Received: 9.07.2023 Accepted: 31.08.2023 Early publication date: 12.10.2023

This article is available in open access under Creative Commons Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.



**Figure 1.** **A.** Four chamber view at 26 weeks of gestation; the pulmonary veins join a collector visible behind the posterior wall of the left atrium; **B.** Enlarged coronary sinus at 26 weeks of gestation; **C.** 26 weeks of gestation; colour Doppler mode; unrestricted blood flow through the collector; **D.** Postnatal suprasternal view: right and left pulmonary veins draining to pulmonary vein collector. See arrows; **E.** Postnatal modified 4-chamber view: enlarged right atrium (RA) and right ventricle (RV), small left atrium (LA), compressed left ventricle (LV), CS indicates the point where pulmonary collector connects to coronary sinus, the red arrow shows atrial septal defect position (ASD); Ao — aorta; CS — coronary sinus; PV — pulmonary vein

valve was tricommisural and trileaflet, and functioning normally. The coronary arteries originated normally. Imaging identified a normal left aortic arch with normal wide isthmus.

Total correction was done on the 10th day of life: CS drainage was converted from the right to the left atrium. The post-surgery course was not complicated. Follow-up examination proved non-restrictive blood flow through the established coronary sinus to the LA connection. The ventricular systolic function and heart output were normal. The neonate was discharged on the 15th day of life in good general condition, with arterial SatO<sub>2</sub> 92–95%. Out-patient follow up indicated the child was healthy and developing well.

In conclusion, though difficult, TAPVC can be diagnosed using antenatal ultrasound. In our case, 2D and colour Doppler were chosen for diagnosis. In-utero diagnosis enabled a multidisciplinary approach and optimal prenatal and postnatal patient counselling and treatment.

#### Article information and declarations

##### Ethics statement

The institutional review board waived the requirement for a separate ethical approval for this clinical vignette, since the sonographic evaluations were performed as integral parts of routine clinical care, for which informed consent had been previously given by the patient. Data were anonymized.

##### Author contributions

Anna Wojtowicz — concept, analysis and interpretation of data, article draft, literature review, corresponding author, Beata Zaluska-Pitak — collecting data, article draft, literature review, Mgadalena Juszcak — literature review, Hubert Huras — literature review, Sebastian Goreczny — verification, literature review.

##### Acknowledgments

None.

##### Conflict of interest

The authors declare no conflict of interest.

##### Supplementary material

Supplementary video and Figures S1–S2 available on [https://journals.viamedica.pl/ginekologia\\_polska/article/view/gpl.96430](https://journals.viamedica.pl/ginekologia_polska/article/view/gpl.96430)

#### REFERENCES

- Hoffman JIE, Kaplan S, Liberthson RR. Prevalence of congenital heart disease. *Am Heart J.* 2004; 147(3): 425–439, doi: [10.1016/j.ahj.2003.05.003](https://doi.org/10.1016/j.ahj.2003.05.003), indexed in Pubmed: [14999190](https://pubmed.ncbi.nlm.nih.gov/14999190/).
- Seale AN, Uemura H, Webber SA, et al. British Congenital Cardiac Association. Total anomalous pulmonary venous connection: morphology and outcome from an international population-based study. *Circulation.* 2010; 122(25): 2718–2726, doi: [10.1161/CIRCULATIONAHA.110.940825](https://doi.org/10.1161/CIRCULATIONAHA.110.940825), indexed in Pubmed: [21135364](https://pubmed.ncbi.nlm.nih.gov/21135364/).
- Borowski D, Pietryga M, Basta P, et al. Practice guidelines of the Polish Society of Gynecologists and Obstetricians - Ultrasound Section for ultrasound screening in uncomplicated pregnancy - 2020. *Ginekol Pol.* 2020; 91(8): 490–501, doi: [10.5603/GP.2020.0110](https://doi.org/10.5603/GP.2020.0110), indexed in Pubmed: [32902850](https://pubmed.ncbi.nlm.nih.gov/32902850/).
- Carvalho JS, Allan LD, Chaoui R, et al. International Society of Ultrasound in Obstetrics and Gynecology. ISUOG Practice Guidelines (updated): sonographic screening examination of the fetal heart. *Ultrasound Obstet Gynecol.* 2013; 41(3): 348–359.
- Seale AN, Carvalho JS, Gardiner HM, et al. British Congenital Cardiac Association. Total anomalous pulmonary venous connection: impact of prenatal diagnosis. *Ultrasound Obstet Gynecol.* 2012; 40(3): 310–318, doi: [10.1002/uog.11093](https://doi.org/10.1002/uog.11093), indexed in Pubmed: [22262371](https://pubmed.ncbi.nlm.nih.gov/22262371/).