

Comparison of the clinical characteristics and prognosis between clear cell carcinomas and high-grade serous ovarian carcinomas

Shixiang Dong^{1,2} , Fengsheng Yu¹ , Yutong Liu^{1,2} , Xiao Yu¹ , Xin Sun¹ ,
Wenjie Wang¹ , Yankui Wang^{1,2} 

¹Department of Obstetrics and Gynecology, Affiliated Hospital of Qingdao University, Qingdao, China

²Qingdao University, Qingdao, China

ABSTRACT

Objectives: To compare the clinical characteristics and prognosis of women with clear cell versus high-grade serous ovarian carcinoma.

Material and methods: Retrospective analysis of the clinical data of 50 cases patients with ovarian clear cell carcinoma (OCCC) and 103 cases with high-grade serous ovarian carcinoma (HGSOC), who were initially treated and completed standardized therapy in Affiliated Hospital of Qingdao University from January 2013 to December 2017.

Results: There were significant differences in age, gravidity ($G > 1$), chief complaint, with ovarian endometriosis, tumor diameter, unilateral or bilateral, cystic and solid tumor, CA125, HE4, CA199, lactate dehydrogenase (LDH), and FIGO stage between the two groups. The differences in the prognosis between OCCC patients and HGSOC patients with early stage (FIGO I–II) were not statistically significant. The 5-year overall survival and progression-free survival of OCCC patients were significantly worse than those of HGSOC patients with advanced stage (FIGO III–IV) ($p < 0.05$). FIGO stage and non-R0 resection were independent risk factors affecting the prognosis of patients with ovarian clear cell carcinoma, screening by Cox regression analysis. FIGO stage, the lowest value of CA125, and non-R0 resection were independent risk factors affecting the prognosis of patients with high-grade serous ovarian cancer.

Conclusions: The clinical characteristics and prognosis of OCCC are different from those of HGSOC. Ovarian clear cell carcinoma (OCCC) patients have a significantly worse prognosis than those with HGSOC in the advanced stage (FIGO III–IV). Satisfactory tumor resection is an essential factor related to the prognosis of patients with OCCC and HGSOC.

Keywords: ovarian clear cell carcinoma; high-grade serous ovarian carcinoma; prognosis; risk factors; clinical characteristics

Ginekologia Polska 2023; 94, 10: 792–798

INTRODUCTION

Epithelial ovarian cancer (EOC) is one of the most lethal malignancies in gynecological tumors. According to 2014 WHO diagnostic criteria [1], the primary tissue types of EOC include seven types (serous carcinoma, endometrioid carcinoma, clear cell carcinoma, mucinous carcinoma, serous mucinous carcinoma, malignant Brenner tumor, and undifferentiated carcinoma), each of which has its unique clinical characteristics and biological behavior. Ovarian clear cell carcinoma (OCCC) is a kind of EOC with unusual biological behavior. Although most OCCC can be detected in

the early stage, it still has a high tendency of recurrence after chemotherapy [2]. Compared with high-grade serous ovarian carcinoma (HGSOC), OCCC has different clinical characteristics and prognoses. A series of large clinical studies on patients with EOC have also excluded patients with clear cells in recent years. As a result, there has been little progress in the treatment of OCCC in the last 30 years. This paper retrospectively analyzed the clinical and follow-up data of 50 patients with OCCC and 103 cases with HGSOC. They were initially treated and completed standardized therapy in the Affiliated Hospital of Qingdao University from

Corresponding author:

Yankui Wang

Department of Obstetrics and Gynecology, Affiliated Hospital of Qingdao University, 16 Jiang-su Road, Qingdao 266073, China

e-mail: qdwwangyk@163.com

Received: 9.06.2022 Accepted: 25.09.2022 Early publication date: 6.12.2022

This article is available in open access under Creative Commons Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

January 2013 to December 2017. We compared the differences in clinical characteristics and prognosis between the two groups. And we evaluated the risk factors affecting the prognosis. We wish we could provide some specific clinical basis for the further study of patients with OCCC.

MATERIAL AND METHODS

Patients

From January 2013 to December 2017, 99 patients with OCCC and 194 patients with HGSOC were registered and treated by the Affiliated Hospital of Qingdao University. The detailed clinical data were collected by reviewing medical charts and clinical follow-up visits. Patients were eligible if they fulfilled the following: (1) had histologically confirmed pure-type OCCC or HGSOC, (2) operations were performed at the Affiliated Hospital of Qingdao University, (3) standard treatment was completed, (4) did not complicate with other primary malignancies or serious complications that affect survival, (5) postoperative follow-up was standardized and complete. Finally, 50 patients with OCCC and 103 patients with HGSOC were enrolled in the current study. Figure 1 shows the data acquisition process. All the patients included had provided written informed consent for the use of their clinical data. The study was approved by the ethics committee at the Affiliated Hospital of Qingdao University in Shandong, China (Ethical approval number: QYFY WZLL 26500).

All the operation was performed by the gynecologic oncologist of Affiliated Hospital of Qingdao University to achieve optimal cytoreduction as far as possible, which was defined as no residual macroscopical lesion after primary debulking surgery. Staging was performed according to the International Federation of Gynecology and Obstetrics (FIGO) staging system (FIGO, 2014). Most of the patients received platinum-based chemotherapy regimens as the postoperative first-line treatment.

Follow-up

All patients were routinely followed up for disease progression by telephone or outpatient examinations until December 2020 or mortality. The Overall Survival (OS) was defined as the time from the diagnosis to death or the last follow-up time. Progression-Free Survival (PFS) was defined as the time from therapy initiation to the time of disease progression, relapse, or the last follow-up time. Disease progression or relapse was defined according to the Gynecologic Cancer Intergroup, GCG [3]. Patients with ovarian cancer after the treatment initiation appeared the following conditions: (1) elevated CA125; (2) a clinically palpable mass; (3) a mass found based on computed tomography, magnetic resonance imaging, or ultrasound; (4) hydrothorax or ascites; (5) Ileus with unknown causes; if 2 out of 5 items are satisfied, it is considered as clinical recurrence; If the patient has only




Figure 1. Flow chart of data selection; HGSOC — high-grade serous ovarian carcinoma; OCCC — ovarian clear cell carcinoma

elevated CA125 without clinical manifestations and radiographic evidence, it is a biochemical recurrence. Elevated CA125 was defined as two consecutive occasions when CA125 was greater than the normal upper limit (35 U/mL) in patients with normal CA125 or in patients whose values were normalized during treatment. For patients who never normalized CA125, it was greater than twice the nadir on two consecutive measurements.

Statistical analysis

Statistical Package for Social Science (SPSS) statistical software (Version 20.0, SPSS Inc, Chicago, Ill) was used for all analyses. T-test was used for the measurement data following a normal distribution. A nonparametric test was used for data that did not conform to normal distribution. The Pearson's chi-squared test, Yate's correction for continuity, or Fisher exact test was used to analyze the enumeration data. Kaplan–Meier curves were used to analyze the distribution of OS and PFS by groups. The log-rank test examined the comparison of patient survival among subgroups. Multivariate survival analysis was performed using the Cox regression model, including prognostic factors that were significant in univariate analysis by Logistic regression. All of the p values reported were 2-sided, and a value of $p < 0.05$ was considered statistically significant.

RESULTS

Patients' characteristics

The patients' characteristics between the two groups are summarized in Table 1. Among the patients with malignant

Characteristics	OCCC (N = 50)	HGSOC (N = 103)	P
Age, mean (SD) [years]	51.3 (7.8)	56.1 (8.9)	0.001†
≤ 50, years n (%)	24 (48.0)	31 (30.1)	0.030‡
>50 years, n (%)	26 (52.0)	72 (69.9)	
Menopausal status, n (%)			
No	21 (42.0)	35 (34.0)	0.334‡
Yes	29 (58.0)	68 (66.0)	
Gravidity, times			
≤ 1 time, n (%)	19 (38.0)	19 (18.4)	0.009‡
>1 time, n (%)	31 (62.0)	84 (81.6)	
Intrauterine device use, n (%)			
No	39 (78.0)	83 (80.6)	0.709‡
Yes	11 (22.0)	20 (19.4)	
Chief complaint, n (%)			
Abdominal distension	12 (24.0)	63 (61.2)	0.001§
Abdominal pain	16 (32.0)	40 (38.8)	
Menstrual alterations or post-menopausal vaginal bleeding	10 (20.0)	10 (9.7)	
Asymptomatic	16 (32.0)	13 (12.6)	
With ovarian endometriosis, n (%)			
No	32 (64.0)	102 (99.0)	< 0.001‡
Yes	18 (36.0)	1 (1.0)	
Tumor diameter, mean (SD) [cm]	11.6 (4.3)	8.4 (4.2)	< 0.001†
≤ 10 cm, n (%)	19 (38.0)	69 (67.0)	0.001‡
> 10 cm, n (%)	31 (62.0)	34 (33.0)	
Bilateral tumors, n (%)			
No	41 (82.0)	36 (35.0)	< 0.001‡
Yes	9 (18.0)	67 (65.0)	
Cystic solid tumor, n (%)			
No	12 (24.0)	46 (44.7)	0.013‡
Yes	38 (76.0)	57 (55.3)	
CA125, mean (SD), U/mL	288.5 (435.2)	1570.4 (2068.1)	< 0.001†
≤ 1000 U/mL, n (%)	46 (92.0)	59 (57.3)	< 0.001‡
> 1000 U/mL, n (%)	4 (8.0)	44 (42.7)	
The nadir of CA125 after treatment, mean (SD), U/mL	37.1 (127.5)	30.5 (113.9)	0.748†
≤ 10 U/mL, n (%)	29 (58.0)	50 (48.5)	0.272‡
> 10 U/mL, n (%)	21 (42.0)	53 (51.5)	
He4, mean (SD), pmol/L	153.0 (205.1)	689.9 (778.6)	< 0.001†
≤ 140 pmol/L, n (%)	37 (74.0)	20 (19.4)	< 0.001‡
> 140 pmol/L, n (%)	13 (26.0)	83 (80.6)	

Characteristics	OCCC (N = 50)	HGSOC (N = 103)	P
CA199, mean (SD), U/mL	51.6 (80.8)	23.3 (47.9)	0.025†
≤ 27 U/mL, n (%)	29 (58.0)	84 (81.6)	0.002‡
> 27 U/mL, n (%)	21 (42.0)	19 (18.4)	
Blood calcium, mean (SD), mmol/L	2.2 (0.2)	2.2 (0.2)	0.824†
≤ 2.52 mmol/L, n (%)	50 (100.0)	100 (97.1)	0.551§
>2.52 mmol/L, n (%)	0 (0.0)	3 (2.9)	
Lactate dehydrogenase, mean (SD), U/L	191.3 (61.4)	253.4 (166.4)	0.001†
≤ 250 U/L, n (%)	43 (86.0)	67 (65.0)	0.007‡
> 250 U/L, n (%)	7 (14.0)	36 (35.0)	
Platelet to lymphocyte ratio, mean (SD)	216.5 (104.5)	242.3 (132.2)	0.229†
≤ 200, n (%)	24 (48.0)	46 (44.7)	0.697‡
> 200, n (%)	26 (52.0)	57 (55.3)	
R0 resection, n (%)			
No	7 (14.0)	28 (27.2)	0.069‡
Yes	43 (86.0)	75 (72.8)	
FIGO stage, n (%)			
I–II	37 (74.0)	23 (22.3)	< 0.001‡
III–IV	13 (26.0)	80 (77.7)	

† — t Test; ‡ — Pearson's chi-squared test; § — Fisher exact test; HGSOC — high-grade serous ovarian carcinoma; OCCC — ovarian clear cell carcinoma; R0 — resection, complete resection; SD — standard deviation

ovarian cancer who visited our hospital (n = 697) from January 2013 to December 2017, 99 patients (14.2%) were diagnosed with OCCC, and 194 patients (27.8%) were diagnosed with HGSOC. There were 50 patients in the group of OCCC with the average age of 51.3 years old and 103 patients in the HGSOC group with the average age of 56.1 years old included in the present analysis. Patients with confirmed OCCC were younger than HGSOC (p < 0.05). Most of the OCCC showed a large unilateral cystic solid tumor. The proportion of asymptomatic patients was higher in the OCCC group than in the HGSOC group. Compared with HGSOC, OCCC patients had fewer pregnancies and more patients with ovarian endometriosis. There is a great difference between OCCC and HGSOC in terms of tumor markers. The values of CA125, HE4, and LDH in the OCCC group were lower than that in the HGSOC group, while the CA199 values were higher than that in the HGSOC group. Among the patients with CA125 < 1000, 45.7% of the patients in the OCCC group had CA199 > 27, while only 18.6% in the HGSOC group (p < 0.003). At the same time, the proportion of patients in stages I–II was 74% in the OCCC group and only 22.3% in the HGSOC group (p < 0.0001).




Figure 2. Left, overall survival in the two groups, in International Federation of Gynecology and Obstetrics (FIGO) stages I–II (A), and in FIGO stages III–IV (C). Right, progression-free survival in the two groups in FIGO stages I–II (B), and in FIGO stages III–IV (D); HGSOC — high-grade serous ovarian carcinoma; OCCC — ovarian clear cell carcinoma

Prognosis of patients in the 2 groups with different stages

During the median follow-up time of 44 months (range, 6–93 months), 15 patients (30.0%) in the OCCC group and 52 patients (50.5%) in the HGSOC group were dead ($p = 0.017$). Figure 2 depicts the OS and PFS of each group. In patients with FIGO stages I–II, the 5-year OS and PFS rates of patients with OCCC (83.8% and 89.2%, respectively) were higher than those of patients with HGSOC (67.2% and 87.0%, respectively). However, this was not statistically significant ($p = 0.786$ and $p = 0.194$, respectively). In patients with FIGO stages III–IV, the 5-year OS and PFS rates of patients with OCCC (15.4% and 0.0%, respectively) were significantly lower than those of patients with HGSOC (34.1% and 11.4%, respectively; $p < 0.001$ and $p < 0.001$, respectively). The median OS and PFS for the two groups with FIGO stages III–IV were: OCCC (OS/PFS), 20/7 months; HGSOC (OS/PFS), 53/16 months.

In the univariate survival analysis by Logistic regression, significant prognostic factors for OCCC were the lowest value of CA125, lactate dehydrogenase, bilateral tumors, R0 resection, and FIGO stage. Significant prognostic factors for HGSOC were age, menopausal status, intrauterine device use, the lowest value of CA125, R0 resection, and FIGO stage (Tab. 2). Multivariate Cox regression analyses revealed that R0 resection and FIGO stage were independent risk

factors for the prognosis of patients with OCCC. And the lowest value of CA125, R0 resection, and FIGO stage were independent risk factors for the prognosis of patients with HGSOC (Tab. 3).

DISCUSSION

Ovarian clear cell carcinoma accounts for 5% to 25% of EOC with obvious regionality. Ovarian clear cell carcinoma incidence was markedly higher in Asia than in other regions [4]. The rate of early detection of OCCC is also higher than that of HGSOC. However, although a high proportion of patients with OCCC are detected early, much literature has reported that the prognosis of patients with OCCC is similar to or worse than that of patients with HGSOC [5–9]. Compared with other histologic types of EOC, OCCC is more aggressive and less sensitive to platinum-based chemotherapy [10].

Oliver et al. [4] enrolled 544 patients with OCCC and 7054 patients with serous carcinoma. They found that patients with OCCC were younger and had a higher proportion of early stage. FIGO stages I–II accounted for more than 50% (57–81%) of patients with OCCC. Ovarian clear cell carcinoma also has some special clinical features, such as frequent presentation as a large pelvic mass, association with endometriosis, vascular thrombotic events, and hypercalcemia [2]. Although no patients with OCCC were found

Table 2. Univariate analyses of prognosis by logistic regression

Variable	OCCC			HGSOC		
	OR	95% CI	p	OR	95% CI	p
Age						
> 50 (vs ≤ 50)	0.500	0.146–1.712	0.270	2.940	1.211–7.137	0.017
Menopausal status						
Yes (vs No)	0.348	0.100–1.210	0.097	4.038	1.673–9.751	0.002
Intrauterine device use						
Yes (vs No)	2.417	0.603–9.678	0.213	0.255	0.085–0.768	0.015
The lowest value of CA125						
> 10 (vs ≤ 10)	6.875	1.766–26.765	0.005	3.775	1.669–8.534	0.001
Lactate dehydrogenase						
> 250 (vs ≤ 250)	22.667	2.411–213.102	0.006	0.971	0.432–2.182	0.942
Bilateral tumors						
Yes (vs no)	14.437	2.506–83.168	0.003	2.059	0.901–4.704	0.087
RO resection						
No (vs yes)	22.667	2.411–213.102	0.006	14.815	4.090–53.660	< 0.001
FIGO stage						
III–IV (vs I–II)	45.375	7.283–282.695	< 0.001	10.538	2.889–38.441	< 0.001

CI — confidence interval; OR — odds ratio; HGSOC — high-grade serous ovarian carcinoma; OCCC — ovarian clear cell carcinoma

Table 3. Multivariate analyses of prognosis by multivariate Cox regression

Variable	OCCC			HGSOC		
	HR	95% CI	p	HR	95% CI	p
The lowest value of CA125						
> 10 (vs ≤ 10)	2.995	0.727–12.337	0.129	2.087	1.123–3.880	0.020
RO resection						
No (vs yes)	7.884	1.689–36.811	0.009	3.693	1.940–7.030	< 0.001
FIGO stage						
III–IV (vs I–II)	4.555	1.088–19.071	0.038	3.910	1.186–12.889	0.025

CI — confidence interval; HGSOC — high-grade serous ovarian carcinoma; HR — hazard ratio; OCCC — ovarian clear cell carcinoma

to have hypercalcemia in our study, this may be related to our relatively small sample size. The lower incidence of bilaterality of OCCC compared to HGSOC observed in this study has been confirmed by other authors [11]. Strong evidence for an association between ovarian endometriosis and OCCC has been established in many studies [11–14]. Patients with ovarian endometriosis have a higher risk of developing ovarian cancer, and the risk was particularly elevated in subjects with a long-standing history of ovarian endometriosis [12]. Park et al. [14] even proposed that ovarian endometriosis may be a precancerous lesion of OCCC. In the present study, 36% of OCCC patients were complicated with ovarian endometriosis, compared with 1.0% of HGSOC patients ($p < 0.01$), which fully demonstrated the association between ovarian endometriosis and OCCC.

Compared with other types of EOC, OCCC lacks effective tumor biomarkers. CA125, which plays a vital role in other types of ovarian cancer, has less clinical significance in OCCC [15]. In recent years, some scholars have studied the clinical significance of CA199 in OCCC. Nakagawa et al. [16] reported that 54% of OCCC patients were associated with elevated CA199. Zhu et al. [17] suggested that CA199 may help to distinguish the prognosis of patients. The clinical value of CA199 in patients with OCCC is worthy of further investigation. Hypercalcemia is one of the most common paraneoplastic syndromes in malignant tumors. However, ovarian cancer with hypercalcemia is rarely reported. Japanese scholars Fujino et al. [18] proposed that OCCC is most closely related to hypercalcemia in patients with ovarian cancer. For patients with OCCC complicated with hypercalcemia, recurrence is often accompanied by an increase in serum

calcium. However, none of the 50 OCCC patients included in this study were complicated with hypercalcemia, which is inconsistent with the above reports and may be related to the insufficient sample size.

Ovarian clear cell carcinoma has been generally accepted as unfavorable when compared with other types of EOC, which has been supported by several retrospective studies [19–21]. However, Oliver et al. [4] found that patients with OCCC had a better overall prognosis compared with serous carcinoma. Oliver et al. [4] believed that this difference was related to younger age, earlier stage, and better performance status of patients with OCCC. The study also found that, after adjusting the age, stage, and performance status, the prognosis of OCCC was significantly better than serous carcinoma in stages I–II. At the same time, it was significantly worse in stages III–IV. In our study, we used HGSOC as the control group. We found no statistically significant difference in OS and PFS between the two groups in stages I–II. However, in stages III–IV patients, OCCC patients displayed worse OS and PFS.

The factors affecting the prognosis of patients with OCCC continue to be a topic of hot debate in medicine. The sample size of the clinical studies on OCCC by Nasioudis et al. [22] was relatively large. Nasioudis et al. [22] evaluated the effect of chemotherapy on prognosis in 2325 OCCC patients with stage I. They found that the survival benefit of chemotherapy on patients with OCCC in the early stage might only be apparent when the lesion was confined to the ovary. Jenison et al. [11] found that incomplete capsules had a significant adverse effect on the prognosis of patients with OCCC in stage I. Therefore, to improve the prognosis of patients with OCCC, we should try our best to avoid iatrogenic upgrading.

CONCLUSIONS

In conclusion, there are many differences in clinical features and prognosis between OCCC and HGSOC. Most of the OCCC showed a large unilateral cystic solid tumor. The detection of CA199 is more critical in patients with OCCC than in patients with HGSOC. OCCC has a high incidence of early stage. OCCC patients have a significantly worse prognosis than those with HGSOC in the advanced stage (FIGO III–IV). Clinically, we should try to maintain the integrity of the tumor envelope during surgery. R0 resection is an essential factor that can improve the prognosis of patients with both OCCC and HGSOC.

Our study was a retrospective single-center study; a larger prospective multi-center study is needed for further prospective external validation.

Article information and declarations

Ethics approval

This retrospective chart review study involving human participants was in accordance with the ethical standards of the institutional and national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards. The study was approved by the ethics committee at the Affiliated Hospital of Qingdao University in Shandong, China (Ethical approval number: QYFY WZLL 26500).

Consent to participate

All participants signed an informed consent form which was approved by the ethics committee at the Affiliated Hospital of Qingdao University in Shandong, China. (Ethical approval number: QYFY WZLL 26500)

Availability of data and materials

The data during the current study are available from the corresponding author on reasonable request. An supplemental file shows the data in more detail (see Supplemental file 1).

Competing interests

The authors declare that there are no competing interests in this study.

Funding

No funding was received to assist with the preparation of this manuscript.

Acknowledgements

The authors would like to show their sincere gratitude to all the patients and their families involved in this study.

REFERENCES

1. Lu Z, Chen J. Introduction of WHO classification of tumours of female reproductive organs, fourth edition. *Zhonghua Bing Li Xue Za Zhi*, (Chinese). 2014; 43(10): 649–650, indexed in Pubmed: [25567588](#).
2. Behbakht K, Randall TC, Benjamin I, et al. Clinical characteristics of clear cell carcinoma of the ovary. *Gynecol Oncol*. 1998; 70(2): 255–258, doi: [10.1006/gyno.1998.5071](#), indexed in Pubmed: [9740700](#).
3. Vergote I, Rustin GJ, Eisenhauer EA, et al. Re: new guidelines to evaluate the response to treatment in solid tumors [ovarian cancer]. *Gynecologic Cancer Intergroup. J Natl Cancer Inst*. 2000; 92(18): 1534–1535, doi: [10.1093/jnci/92.18.1534](#), indexed in Pubmed: [10995813](#).
4. Oliver KE, Brady WE, Birrer M, et al. An evaluation of progression free survival and overall survival of ovarian cancer patients with clear cell carcinoma versus serous carcinoma treated with platinum therapy: An NRG Oncology/Gynecologic Oncology Group experience. *Gynecol Oncol*. 2017; 147(2): 243–249, doi: [10.1016/j.ygyno.2017.08.004](#), indexed in Pubmed: [28807367](#).
5. Chan JK, Teoh D, Hu JM, et al. Do clear cell ovarian carcinomas have poorer prognosis compared to other epithelial cell types? A study of

- 1411 clear cell ovarian cancers. *Gynecol Oncol.* 2008; 109(3): 370–376, doi: [10.1016/j.ygyno.2008.02.006](https://doi.org/10.1016/j.ygyno.2008.02.006), indexed in Pubmed: [18395777](https://pubmed.ncbi.nlm.nih.gov/18395777/).
6. Crozier MA, Copeland LJ, Silva EG, et al. Clear cell carcinoma of the ovary: a study of 59 cases. *Gynecol Oncol.* 1989; 35(2): 199–203, doi: [10.1016/0090-8258\(89\)90043-7](https://doi.org/10.1016/0090-8258(89)90043-7), indexed in Pubmed: [2807010](https://pubmed.ncbi.nlm.nih.gov/2807010/).
 7. Dembo AJ, Davy M, Stenwig AE, et al. Prognostic factors in patients with stage I epithelial ovarian cancer. *Obstetrics & Gynecology.* 1990; 72(2): 263–273, indexed in Pubmed: [2300355](https://pubmed.ncbi.nlm.nih.gov/2300355/).
 8. Kennedy AW, Markman M, Biscotti CV, et al. Survival probability in ovarian clear cell adenocarcinoma. *Gynecol Oncol.* 1999; 74(1): 108–114, doi: [10.1006/gyno.1999.5445](https://doi.org/10.1006/gyno.1999.5445), indexed in Pubmed: [10385560](https://pubmed.ncbi.nlm.nih.gov/10385560/).
 9. Leitao MM, Boyd J, Hummer A, et al. Clinicopathologic analysis of early-stage sporadic ovarian carcinoma. *Am J Surg Pathol.* 2004; 28(2): 147–159, doi: [10.1097/00000478-200402000-00001](https://doi.org/10.1097/00000478-200402000-00001), indexed in Pubmed: [15043303](https://pubmed.ncbi.nlm.nih.gov/15043303/).
 10. Pectasides D, Fountzilas G, Aravantinos G, et al. Advanced stage clear-cell epithelial ovarian cancer: the Hellenic Cooperative Oncology Group experience. *Gynecol Oncol.* 2006; 102(2): 285–291, doi: [10.1016/j.ygyno.2005.12.038](https://doi.org/10.1016/j.ygyno.2005.12.038), indexed in Pubmed: [16516283](https://pubmed.ncbi.nlm.nih.gov/16516283/).
 11. Jenison EL, Montag AG, Griffiths CT, et al. Clear cell adenocarcinoma of the ovary: a clinical analysis and comparison with serous carcinoma. *Gynecol Oncol.* 1989; 32(1): 65–71, doi: [10.1016/0090-8258\(89\)90852-4](https://doi.org/10.1016/0090-8258(89)90852-4), indexed in Pubmed: [2642454](https://pubmed.ncbi.nlm.nih.gov/2642454/).
 12. Brinton LA, Gridley G, Persson I, et al. Cancer risk after a hospital discharge diagnosis of endometriosis. *Am J Obstet Gynecol.* 1997; 176(3): 572–579, doi: [10.1016/s0002-9378\(97\)70550-7](https://doi.org/10.1016/s0002-9378(97)70550-7), indexed in Pubmed: [9077609](https://pubmed.ncbi.nlm.nih.gov/9077609/).
 13. King CM, Barbara C, Prentice A, et al. Models of endometriosis and their utility in studying progression to ovarian clear cell carcinoma. *J Pathol.* 2016; 238(2): 185–196, doi: [10.1002/path.4657](https://doi.org/10.1002/path.4657), indexed in Pubmed: [26456077](https://pubmed.ncbi.nlm.nih.gov/26456077/).
 14. Park JY, Kim DY, Suh DS, et al. Significance of ovarian endometriosis on the prognosis of ovarian clear cell carcinoma. *Int J Gynecol Cancer.* 2018; 28(1): 11–18, doi: [10.1097/IGC.0000000000001136](https://doi.org/10.1097/IGC.0000000000001136), indexed in Pubmed: [28930811](https://pubmed.ncbi.nlm.nih.gov/28930811/).
 15. Kim HS, Choi HY, Lee M, et al. Systemic inflammatory response markers and CA-125 levels in ovarian clear cell carcinoma: a two center cohort study. *Cancer Res Treat.* 2016; 48(1): 250–258, doi: [10.4143/crt.2014.324](https://doi.org/10.4143/crt.2014.324), indexed in Pubmed: [25761476](https://pubmed.ncbi.nlm.nih.gov/25761476/).
 16. Nakagawa N, Koda H, Nitta N, et al. Reactivity of CA19-9 and CA125 in histological subtypes of epithelial ovarian tumors and ovarian endometriosis. *Acta Med Okayama.* 2015; 69(4): 227–235, doi: [10.18926/AMO/53559](https://doi.org/10.18926/AMO/53559), indexed in Pubmed: [26289914](https://pubmed.ncbi.nlm.nih.gov/26289914/).
 17. Zhu J, Jiang L, Wen H, et al. Prognostic value of serum CA19-9 and perioperative CA-125 levels in ovarian clear cell carcinoma. *Int J Gynecol Cancer.* 2018; 28(6): 1108–1116, doi: [10.1097/IGC.0000000000001293](https://doi.org/10.1097/IGC.0000000000001293), indexed in Pubmed: [29781825](https://pubmed.ncbi.nlm.nih.gov/29781825/).
 18. Fujino T, Watanabe T, Yamaguchi K, et al. The development of hypercalcemia in a patient with an ovarian tumor producing parathyroid hormone-related protein. *Cancer.* 2015; 70(12): 2845–2850, doi: [10.1002/1097-0142\(19921215\)70:12<2845::aid-cncr2820701221>3.0.co;2-f](https://doi.org/10.1002/1097-0142(19921215)70:12<2845::aid-cncr2820701221>3.0.co;2-f).
 19. Goff BA, Sainz de la Cuesta R, Muntz HG, et al. Clear cell carcinoma of the ovary: a distinct histologic type with poor prognosis and resistance to platinum-based chemotherapy in stage III disease. *Gynecol Oncol.* 1996; 60(3): 412–417, doi: [10.1006/gyno.1996.0065](https://doi.org/10.1006/gyno.1996.0065), indexed in Pubmed: [8774649](https://pubmed.ncbi.nlm.nih.gov/8774649/).
 20. Winter WE, Maxwell GL, Tian C, et al. Prognostic factors for stage III epithelial ovarian cancer: a Gynecologic Oncology Group Study. *J Clin Oncol.* 2007; 25(24): 3621–3627, doi: [10.1200/JCO.2006.10.2517](https://doi.org/10.1200/JCO.2006.10.2517), indexed in Pubmed: [17704411](https://pubmed.ncbi.nlm.nih.gov/17704411/).
 21. Duska LR, Garrett L, Henretta M, et al. When 'never-events' occur despite adherence to clinical guidelines: the case of venous thromboembolism in clear cell cancer of the ovary compared with other epithelial histologic subtypes. *Gynecol Oncol.* 2010; 116(3): 374–377, doi: [10.1016/j.ygyno.2009.10.069](https://doi.org/10.1016/j.ygyno.2009.10.069), indexed in Pubmed: [19922988](https://pubmed.ncbi.nlm.nih.gov/19922988/).
 22. Nasioudis D, Mastroyannis SA, Albright BB, et al. Adjuvant chemotherapy for stage I ovarian clear cell carcinoma: Patterns of use and outcomes. *Gynecol Oncol.* 2018; 150(1): 14–18, doi: [10.1016/j.ygyno.2018.04.567](https://doi.org/10.1016/j.ygyno.2018.04.567), indexed in Pubmed: [29751993](https://pubmed.ncbi.nlm.nih.gov/29751993/).