

Figure S1. In-hospital mortality in patients managed invasively by age groups




Figure S2. In-hospital mortality in patients managed non-invasively by age groups




Figure S3. Proportions of patients suffering from recurrent myocardial infarction within one year following discharge by age groups




Figure S4. Proportions of patients suffering from stroke within one year following discharge by age groups




Figure S5. Proportions of patients hospitalized at least once for cardiovascular reasons within one year following discharge by age groups




Table S1. ICD-10 codes used to define the index hospitalization for acute myocardial infarction and the end-points

Diagnosis	ICD-10 codes
Acute myocardial infarction	I21; I21.0; I21.1; I21.2; I21.3; I21.4; I21.9; I22; I22.0; I22.1; I22.8; I22.9
Stroke	I60; I60.0; I60.1; I60.2; I60.3; I60.4; I60.5; I60.6; I60.7; I60.8; I60.9; I61; I61.0; I61.1; I61.2; I61.3; I61.4; I61.5; I61.6; I61.8; I61.9; I62; I62.0; I62.1; I62.9; I63; I63.0; I63.1; I63.2; I63.3; I63.4; I63.5; I63.6; I63.8; I63.9; I64
Hospitalizations for cardiovascular reasons	I00-I95; G46; J81; K55; Q20-Q28; R00, R01; R03; R07.2; R09.2; R57.0