



Biochemical and clinical evaluation of endothelial injury after distal or traditional transradial access in percutaneous interventions

Authors: Karol Momot, Maciej Zarębiński, Krzysztof Flis, Katarzyna Czarzasta, Liana Puchalska,
Małgorzata Wojciechowska

Article type: Original article

Received: December 14, 2021

Accepted: April 20, 2022

Early publication date: April 21, 2022

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

Biochemical and clinical evaluation of endothelial injury after distal or traditional transradial access in percutaneous interventions

Short title: Local endothelial injury after percutaneous interventions

Karol Momot¹, Maciej Zarębiński², Krzysztof Flis², Katarzyna Czarzasta¹, Liana Puchalska¹, Małgorzata Wojciechowska¹

¹Laboratory of Center for Preclinical Research, Department of Experimental and Clinical Physiology, Medical University of Warsaw, Warszawa, Poland

²Department of Invasive Cardiology, Lazarski University, Independent Public Specialist Western Hospital John Paul II, Grodzisk Mazowiecki, Poland

Correspondence to:

Maciej Zarębiński, MD, PhD,

Department of Invasive Cardiology,

Lazarski University,

Independent Public Specialist Western Hospital John Paul II,

Daleka 11, 05–825 Grodzisk Mazowiecki, Poland,

phone: +48 22 755 92 90,

e-mail: maciej@zarebinski.pl

Conflict of interest: None declared

WHAT'S NEW?

This study provides a wider perspective of the distal transradial access in percutaneous interventions. This report is the first one which compares transradial approaches in terms of endothelial injury. We list advantages and disadvantages of distal and traditional transradial access. We confirm that new distal transradial access should be widely used in invasive cardiology and radiology.

ABSTRACT

Background: Distal transradial access (dTRA) has been proposed as an alternative to traditional transradial access (TRA) in cardiac catheterization.

Aims: The aim of the study was to compare these two transradial approaches: TRA and dTRA in terms of clinical and biochemical aspects.

Methods: Two hundred patients qualified for elective coronary procedure were included. The patients were assigned to one of the groups depending on the vascular access. The groups were compared in terms of perceived pain using Visual Analogue Scale (VAS), time of gaining the access, need for conversion and local complications. Additionally, in forty patients circulating endothelial injury markers: Endothelin-1 (ET-1), Interleukin-8 (IL-8) and Soluble Vascular Cell Adhesion Molecule-1 (sVCAM-1) were assessed.

Results: Successful cannulation was obtained in 84 (100%) in the TRA group and in 98 (84%) subjects in the dTRA ($P < 0.001$). dTRA was associated with higher level of pain perceived at the time of gaining vascular approach than TRA; median VAS score (IQR): 4 (2-5) vs. 2 (2-4) ($P = 0.04$). The mean time (standard deviation [SD]) needed to cannulate the artery in dTRA was longer than in TRA: 81 (8) seconds vs. 50 (4) seconds ($P = 0.04$). ET-1 concentration was (SD): 2.08 (0.19) [dTRA] vs. 2.00 (0.29) [TRA] pg/ml ($P = 0.83$); sVCAM-1: 12.71 (3.97) vs. 12.86 (4.29) ng/ml ($P = 0.98$); IL-8: 8.81 (0.42) vs. 9.15 (0.52) ng/ml ($P = 0.62$). Number of complications after procedures did not differ between these two approaches.

Conclusions: Cannulation of dTRA is associated with lower success rate and higher pain perceived. dTRA is not inferior to TRA when safety issues and vascular injury are considered.

Key words: percutaneous interventions, coronarography, radial access

INTRODUCTION

Vast increase in number of percutaneous diagnostic and therapeutic procedures performed within the last two decades lead to the need of arterial access associated with decreased complication rates and shorter postoperative care. Advantages of radial artery access over femoral has been well proven. The radial cannulation, which is predominantly used in coronary interventions, may be complicated by occlusion of the artery. The occlusion rate increases with repeating of interventions. The distal transradial access (dTRA) (Figure 1) via the anatomical snuffbox

decreases this complication rate even more and it has been proposed as an alternative to traditional transradial access (TRA) (Figure 2) in percutaneous interventions. The anatomical snuffbox is located on the radial side of the wrist, it is bounded by the tendon of the extensor pollicis longus posteriorly and of the tendons of the extensor pollicis brevis and abductor pollicis longus anteriorly. The radial artery crosses the floor that is formed by the scaphoid and the trapezium bones [1].

First studies pointed out that dTRA was associated with increased rate of cannulation failure, prolonged duration of cannulation, increased number of attempts and number of skin punctures compared to the TRA [2]. This was mainly due to the smaller diameter of the vessel, tortuosity of the radial artery in the area of the anatomical snuffbox and the lack of experience of operators. There is no data comparing the two accesses in terms of endothelial damage. During percutaneous intervention and hemostatic compression mechanical stress of the cannulated vessels occurs. It is caused by the needle puncturing the artery wall, the sheath inserted in the lumen of the artery and external pressure of the dressing. Studies show that after exposure to stress factors, endothelium releases numerous substances like cytokines which can be assessed in blood plasma [3–9]. We believe that these substances are also released during percutaneous interventions.

In this study, concentrations of markers of endothelial injury were measured. This is a novel perspective, as it is the first comparison of the analyzed percutaneous approaches based on biochemical assessment. The markers: Endothelin-1 (ET-1), Interleukin-8 (IL-8) and Soluble Vascular Cell Adhesion Molecule-1 (sVCAM-1) were chosen based on relevant literature.

METHODS

Study patients

Two hundred adult patients, scheduled for the elective coronary angiography or angioplasty, were recruited. Procedures were performed between November 2020 and April 2021. Participants signed written informed consent forms. The study was approved by the Local Ethics Committee (KB/167/2020). Patients with estimated glomerular filtration rate (eGFR) <30 ml/min/1.73 m², dialyzed, with coronary artery bypass grafting (CABG) and/or diagnosed with active cancer were not included in this study. Based on the research of others, like Koutouzis et al. and Wretowski et al. [2, 10] and our experience with the lack of distal pulse and the need for conversion from distal

to traditional access, all qualified patients were allocated to two approaches (dTRA and TRA) by block randomization at ratio 3:2, receiving group sizes: 120 and 80, respectively. Four subjects out of 120 initially assigned to the dTRA approach did not present palpable pulse in an anatomical snuffbox before the procedure and they were reassigned to the TRA access without any attempts to cannulate dTRA. Therefore, operators attempted to obtain 116 distal approaches and 84 traditional approaches. After failed attempt to cannulate dTRA, these subjects were converted to TRA and were included in a third group named conversion ($n = 18$). Thus, final dTRA group included 98 subjects. Demographic data of the patients are presented in [Table 1](#). In forty random patients (20 from dTRA group and 20 from TRA group) after the dressing removal, blood from cephalic vein was collected and plasma concentrations of ET-1, IL-8, sVCAM-1 were determined using the enzyme-linked immunoassays (ELISA).

Procedure

The procedures of coronary interventions without ultrasound guidance were performed by EAPCI (European Association of Percutaneous Cardiovascular Interventions) certified operators, using radial access in more than 95% of routine procedures. Sheath size used for all the procedures was 6 F. Time needed to gain vascular access was assessed. The amount of injected contrast and the total radial dose were recorded. After the procedure a pressure dressing was applied to the puncture site. The dressing was removed after 120 minutes. The puncture site was assessed for the presence of hematoma and pulse. After the removal of the dressing, 10 ml of blood from the cephalic vein was collected to tubes with EDTA-K2 anticoagulant and then centrifuged. Samples with obtained plasma were immediately frozen and stored at -20°C until the moment of biochemical assessment. Plasma concentrations of markers were analyzed using ELISA: IL-8 Human ELISA Kit (KHC0081; Thermo Fisher Scientific, Inc., Waltham, MA, US), Human sVCAM-1/CD106 ELISA Kit (MBS2505831; MyBioSource, San Diego, CA, US) and Endothelin-1 Quantikine ELISA Kit (DET100; R&D Systems, Inc., Minneapolis, MN, US). Each ELISA test was carried out in accordance with the instructions provided by the manufacturer.

Statistical analysis

Quantitative variables are presented as medians (interquartile range [IQR]) or means (standard deviation [SD]). The ANOVA test (normal distribution) and the Kruskal-Wallis H test (non-

normal distribution) were performed in the comparison of numerical variables between the three groups. Appropriate post-hoc tests were then performed (Dunn's and Tukey's tests, respectively). Student's t test (normal distribution) and Mann–Whitney U test (non-normal distribution) were used to perform inter-group comparisons. Equality of variances was assessed by Levene's test. Categorical variables are expressed as numbers and percentages and compared using χ^2 test. Statistical data were considered significant with a *P*-value <0.05. All statistical analyses were performed using Statistica 13 software.

RESULTS

Success rate of obtaining a vascular approach in the dTRA group was 84% and 100% in the TRA group (*P* <0.001). In eighteen dTRA subjects (16%) operators failed to gain the vascular access. The approach was changed to TRA, and then successfully gained. These eighteen subjects were included in the third group named “conversion”. Data on the procedure, the mean time required to gain the access; local complications; pain when gaining the vascular approach and during the maintenance of pressure dressing are presented in **Table 2**. Results of biochemical evaluation are presented in **Figure 3**. Regardless of the approach, concentrations of endothelial markers were not correlated with smoking, diabetes, hypertension, kidney disease and coronary disease. Subjects treated with statins had lower ET-1 concentration (SD) than subjects without statin therapy, irrespective of the access: 1.63 (0.24) vs. 2.33 (0.21) pg/ml (*P* = 0.04). Subjects with obesity had higher levels of IL-8 than subjects without obesity, regardless of the approach (*P* = 0.04) (**Figure 4**).

DISCUSSION

Markers of endothelial injury

At the moment of publication, there are several studies which compare these two approaches. However, this research is the first one which provides a closer look at them in terms of endothelial injury. The aim was to evaluate selected markers of vascular injury, dysfunction and inflammation between patients after distal transradial access and traditional transradial access. It is assumed that during percutaneous interventions endothelial injury, inflammation and dysfunction is caused by vascular sheath and catheter insertion and hemostatic compression. These factors are closely linked to mechanical stretch, shear stress and external pressure, which are factors stimulating release of

endothelial injury markers in vitro. The choice of these markers was based on the literature found via PubMed, ScienceDirect, Scopus or Google Scholar.

Endothelin-1

ET-1 is produced mainly by vascular ECs (endothelial cells) and it is acknowledged as the most common ET in humans. This particle is frequently assessed in diagnostics of endothelial dysfunction, injury or inflammation. At the normal state, ET-1 is mainly secreted abluminally towards the vascular smooth muscles and its levels in blood are fairly low. However, in the case of endothelium stimulus, ET-1 is released into the blood from ECs [6, 11]. Mechanical strain of vessel damages vascular wall and stimulates secretion of ET-1 from ECs [12, 13]. Previous studies also indicate shear stress as a factor promoting ET-1 production [7, 8].

Soluble Vascular Cell Adhesion Molecule-1

sVCAM-1 is a circulating particle derived from damaged or activated ECs [3]. Damaged endothelial glycocalyx of vessel walls leads to the increase of sVCAM-1 levels [14]. Shear stress is also the factor that stimulates sVCAM-1 from ECs (9). Elevation of blood pressure activates the expression of adhesion molecules (15). Prolonged mechanical wall stretch promotes VCAM-1 gene expression in ECs [16]. sVCAM-1 plays an important role in accelerating atherosclerosis by facilitating the attachment of inflammatory cells to the vascular endothelial wall and promoting their subsequent migration through the endothelium [17].

Interleukin-8

IL-8 is a proinflammatory chemokine produced by ECs. IL-8 is stored inside ECs and ET-1 promotes releasing this chemokine in vitro [18]. Elevated levels of IL-8 have been found in the area of injured endothelium [5]. The IL-8 concentration starts to increase 1 hour after the exposure to the stress factor, and it is caused by IL-8 gene expression [19]. External mechanical pressure on the ECs significantly raises IL-8 secretion from these cells in vitro [4]. IL-8 levels are increased as a result of many inflammatory conditions, so careful exclusion criteria of patients is required. Platelet-derived microparticles (PMPs), which are produced in case of shear stress, induce IL-8 secretion by ECs [20]. IL-8 is a proinflammatory cytokine with atherogenic effects, it accelerates

the movement of neutrophils and T lymphocytes under the endothelium and promotes monocyte adhesion to the vascular wall.

General

There were no statistically significant differences between the dTRA and TRA groups in the number of: cases of hematoma, lack of the distal pulse after interventions, mean amount of contrast used and radial dose. The findings were similar to the results in other studies [2, 10, 21]. Researchers point out that the cannulation time was longer in the dTRA group, which was also in line with our observations [21]. In our research, the level of pain at the time of gaining vascular approach was significantly higher in dTRA group and conversion. The longer time of cannulation and more severe pain during the procedure can be probably explained by less experience of operators in using dTRA. Additionally, in some subjects inexperienced operators had to make conversions when using dTRA. Probably anatomical characteristics of the radial artery in the snuffbox (tortuosity and small diameter) makes the distal approach more complicated and requires more experience from operators. In accordance with the results of other studies, operators should practice in gaining distal approach in order to obtain the same successful cannulation rate, level of pain and time needed to gain the access as in TRA [22, 23]. The advantage of dTRA postulated in other studies is shorter hemostatic compression after the procedure, but in the presented report it was identical in all patients as we wanted to provide the same conditions for biochemical and pain assessment [21, 24]. dTRA offers two more forearm approaches to evaluate and this may reduce the need of femoral artery cannulation. If the radial artery occlusion has occurred during TRA, dTRA provides a possibility to recannulate the occluded radial artery [24]. dTRA offers the option to have the patient's left hand close to the right groin, which is more comfortable for the patient and the operator. dTRA is also beneficial for right-handed patients whose dominant upper limb is without immobilization during the hemostatic compression [1]. Radial artery gives branches before entering the anatomical snuffbox, therefore occlusion after dTRA is related to a smaller area of ischemia than after TRA. Since standard sheath size of 6F allows most of coronary interventions, it is probable that dTRA may serve as a good choice also for complicated high-risk procedures. The safety cannulation with larger sheath has not been tested in our study but with a use of thin-walled sheaths it seems quite possible to use advanced intravascular techniques which requires larger lumen. Januszko et al. showed, that TRA, as opposed to femoral access is related

to higher risk of coronary artery perforation in patients treated with rotational atherectomy [25]. As this complication may also refer to the use of dTRA, future studies should be conducted.

There were no differences between dTRA and TRA in the plasma markers of endothelial injury. This means that in both groups the endothelial damage was similar and that in terms of biochemical assessment, dTRA is at least as safe as TRA. Regardless of the approach, elevated IL-8 levels in obese patients suppose that obesity may be connected with a greater damage of endothelium, but it cannot be excluded that it is constantly elevated in obese subjects, which would be in accordance with other studies [20]. Furthermore, patients treated with statins have significantly lower levels of ET-1 than patients without this treatment, which confirms that statin therapy reduces vascular inflammation [26].

The main limitations of this study were the lack of biochemical evaluation before percutaneous intervention and a small sample size.

CONCLUSIONS

There were no differences between dTRA and TRA in the quotative markers of local endothelial injury. dTRA was more painful for the patient during the cannulation, but the difference should be diminished as the operators gain experience. Consequently, the choice of dTRA is as good as that of TRA. Considering dTRA advantages listed in the discussion, it should be widely used in percutaneous interventions in invasive cardiology, neurology and radiology.

Funding

This study was financed from the student research grant of the Medical University of Warsaw to [KM] (MG/M/13/13/20).

REFERENCES

1. Sgueglia GA, Di Giorgio A, Gasparone A, et al. Anatomic basis and physiological rationale of distal radial artery access for percutaneous coronary and endovascular procedures. *JACC Cardiovasc Interv.* 2018; 11(20): 2113–2119, doi: [10.1016/j.jcin.2018.04.045](https://doi.org/10.1016/j.jcin.2018.04.045), indexed in Pubmed: [30336816](https://pubmed.ncbi.nlm.nih.gov/30336816/).

2. Koutouzis M, Kontopodis E, Tassopoulos A, et al. Distal versus traditional radial approach for coronary angiography. *Cardiovasc Revasc Med*. 2019; 20(8): 678–680, doi: [10.1016/j.carrev.2018.09.018](https://doi.org/10.1016/j.carrev.2018.09.018), indexed in Pubmed: [30314833](https://pubmed.ncbi.nlm.nih.gov/30314833/).
3. Kjaergaard AG, Dige A, Krog J, et al. Soluble adhesion molecules correlate with surface expression in an in vitro model of endothelial activation. *Basic Clin Pharmacol Toxicol*. 2013; 113(4): 273–279, doi: [10.1111/bcpt.12091](https://doi.org/10.1111/bcpt.12091), indexed in Pubmed: [23724832](https://pubmed.ncbi.nlm.nih.gov/23724832/).
4. Weiss D, Avraham S, Guttlieb R, et al. Mechanical compression effects on the secretion of vWF and IL-8 by cultured human vein endothelium. *PLoS One*. 2017; 12(1): e0169752, doi: [10.1371/journal.pone.0169752](https://doi.org/10.1371/journal.pone.0169752), indexed in Pubmed: [28081186](https://pubmed.ncbi.nlm.nih.gov/28081186/).
5. von der Thüsen JH, Kuiper J, van Berkel TJC, et al. Interleukins in atherosclerosis: molecular pathways and therapeutic potential. *Pharmacol Rev*. 2003; 55(1): 133–166, doi: [10.1124/pr.55.1.5](https://doi.org/10.1124/pr.55.1.5), indexed in Pubmed: [12615956](https://pubmed.ncbi.nlm.nih.gov/12615956/).
6. Dhaun N, Webb DJ. Endothelins in cardiovascular biology and therapeutics. *Nat Rev Cardiol*. 2019; 16(8): 491–502, doi: [10.1038/s41569-019-0176-3](https://doi.org/10.1038/s41569-019-0176-3), indexed in Pubmed: [30867577](https://pubmed.ncbi.nlm.nih.gov/30867577/).
7. Marasciulo FL, Montagnani M, Potenza MA. Endothelin-1: the yin and yang on vascular function. *Curr Med Chem*. 2006; 13(14): 1655–1665, doi: [10.2174/092986706777441968](https://doi.org/10.2174/092986706777441968), indexed in Pubmed: [16787211](https://pubmed.ncbi.nlm.nih.gov/16787211/).
8. Davenport AP, Hyndman KA, Dhaun N, et al. Endothelin. *Pharmacol Rev*. 2016; 68(2): 357–418, doi: [10.1124/pr.115.011833](https://doi.org/10.1124/pr.115.011833), indexed in Pubmed: [26956245](https://pubmed.ncbi.nlm.nih.gov/26956245/).
9. Schmidt-Lucke C, Reinhold D, Ansorge S, et al. Changes of plasma concentrations of soluble vascular cell adhesion molecule-1 and vascular endothelial growth factor after increased perfusion of lower extremities in humans. *Endothelium*. 2003; 10(3): 159–165, doi: [10.1080/10623320390233472](https://doi.org/10.1080/10623320390233472), indexed in Pubmed: [13129819](https://pubmed.ncbi.nlm.nih.gov/13129819/).
10. Wretowski D, Krakowian M, Łabyk A, et al. Very distal transradial approach (VITRO) for coronary interventions. *Postepy Kardiol Interwencyjnej*. 2019; 15(1): 42–45, doi: [10.5114/aic.2019.83771](https://doi.org/10.5114/aic.2019.83771), indexed in Pubmed: [31043983](https://pubmed.ncbi.nlm.nih.gov/31043983/).
11. Levin ER. Endothelins. *N Engl J Med*. 1995; 333(6): 356–363, doi: [10.1056/NEJM199508103330607](https://doi.org/10.1056/NEJM199508103330607), indexed in Pubmed: [7609754](https://pubmed.ncbi.nlm.nih.gov/7609754/).

12. Wang DL, Tang CC, Wung BS, et al. Cyclical strain increases endothelin-1 secretion and gene expression in human endothelial cells. *Biochem Biophys Res Commun.* 1993; 195(2): 1050–1056, doi: [10.1006/bbrc.1993.2150](https://doi.org/10.1006/bbrc.1993.2150), indexed in Pubmed: [8373384](https://pubmed.ncbi.nlm.nih.gov/8373384/).
13. Russell FD, Skepper JN, Davenport AP. Human endothelial cell storage granules: a novel intracellular site for isoforms of the endothelin-converting enzyme. *Circ Res.* 1998; 83(3): 314–321, doi: [10.1161/01.res.83.3.314](https://doi.org/10.1161/01.res.83.3.314), indexed in Pubmed: [9710124](https://pubmed.ncbi.nlm.nih.gov/9710124/).
14. Padberg JS, Wiesinger A, di Marco GS, et al. Damage of the endothelial glycocalyx in chronic kidney disease. *Atherosclerosis.* 2014; 234(2): 335–343, doi: [10.1016/j.atherosclerosis.2014.03.016](https://doi.org/10.1016/j.atherosclerosis.2014.03.016), indexed in Pubmed: [24727235](https://pubmed.ncbi.nlm.nih.gov/24727235/).
15. Preston RA, Ledford M, Materson BJ, et al. Effects of severe, uncontrolled hypertension on endothelial activation: soluble vascular cell adhesion molecule-1, soluble intercellular adhesion molecule-1 and von Willebrand factor. *J Hypertens.* 2002; 20(5): 871–877, doi: [10.1097/00004872-200205000-00021](https://doi.org/10.1097/00004872-200205000-00021), indexed in Pubmed: [12011647](https://pubmed.ncbi.nlm.nih.gov/12011647/).
16. Wójtowicz A, Babu SS, Li Li, et al. Zyxin mediation of stretch-induced gene expression in human endothelial cells. *Circ Res.* 2010; 107(7): 898–902, doi: [10.1161/CIRCRESAHA.110.227850](https://doi.org/10.1161/CIRCRESAHA.110.227850), indexed in Pubmed: [20705921](https://pubmed.ncbi.nlm.nih.gov/20705921/).
17. Krieglstein C, Granger DN. Adhesion molecules and their role in vascular disease. *Am J Hypertens.* 2001; 14(11): S44–S54, doi: [10.1016/s0895-7061\(01\)02069-6](https://doi.org/10.1016/s0895-7061(01)02069-6), indexed in Pubmed: [11411765](https://pubmed.ncbi.nlm.nih.gov/11411765/).
18. Øynebråten I, Bakke O, Brandtzaeg P, et al. Rapid chemokine secretion from endothelial cells originates from 2 distinct compartments. *Blood.* 2004; 104(2): 314–320, doi: [10.1182/blood-2003-08-2891](https://doi.org/10.1182/blood-2003-08-2891), indexed in Pubmed: [15044249](https://pubmed.ncbi.nlm.nih.gov/15044249/).
19. Tang R, Zhou J, Cheng M, et al. Time-dependent increase of interleukin-8 production in endothelial cells exposed to fluid shear stress [article in Chinese]. *Sheng Wu Yi Xue Gong Cheng Xue Za Zhi.* 2004; 21(1): 34–37, indexed in Pubmed: [15022458](https://pubmed.ncbi.nlm.nih.gov/15022458/).
20. Kim CS, Park HS, Kawada T, et al. Circulating levels of MCP-1 and IL-8 are elevated in human obese subjects and associated with obesity-related parameters. *Int J Obes (Lond).* 2006; 30(9): 1347–1355, doi: [10.1038/sj.ijo.0803259](https://doi.org/10.1038/sj.ijo.0803259), indexed in Pubmed: [16534530](https://pubmed.ncbi.nlm.nih.gov/16534530/).
21. Aoi S, Htun WW, Freeo S, et al. Distal transradial artery access in the anatomical snuffbox for coronary angiography as an alternative access site for faster hemostasis. *Catheter*

- Cardiovasc Interv. 2019; 94(5): 651–657, doi: [10.1002/ccd.28155](https://doi.org/10.1002/ccd.28155), indexed in Pubmed: [30801939](https://pubmed.ncbi.nlm.nih.gov/30801939/).
22. Oliveira MD, Caixeta A. Distal transradial access (dTRA) for coronary angiography and interventions: A quality improvement step forward? *Case Medical Research*. 2020; 32(9): E238–E239, doi: [10.31525/ct1-nct03948165](https://doi.org/10.31525/ct1-nct03948165), indexed in Pubmed: [32865510](https://pubmed.ncbi.nlm.nih.gov/32865510/).
 23. Cai G, Huang H, Li F, et al. Distal transradial access: a review of the feasibility and safety in cardiovascular angiography and intervention. *BMC Cardiovasc Disord*. 2020; 20(1): 356, doi: [10.1186/s12872-020-01625-8](https://doi.org/10.1186/s12872-020-01625-8), indexed in Pubmed: [32758150](https://pubmed.ncbi.nlm.nih.gov/32758150/).
 24. Davies RE, Gilchrist IC. Dorsal (distal) transradial access for coronary angiography and intervention. *Interv Cardiol Clin*. 2019; 8(2): 111–119, doi: [10.1016/j.iccl.2018.11.002](https://doi.org/10.1016/j.iccl.2018.11.002), indexed in Pubmed: [30832936](https://pubmed.ncbi.nlm.nih.gov/30832936/).
 25. Januszek R, Siudak Z, Malinowski KP, et al. Radial versus femoral access in patients treated with percutaneous coronary intervention and rotational atherectomy. *Kardiol Pol*. 2020; 78(6): 529–536, doi: [10.33963/KP.15286](https://doi.org/10.33963/KP.15286), indexed in Pubmed: [32267133](https://pubmed.ncbi.nlm.nih.gov/32267133/).
 26. Liberale L, Carbone F, Montecucco F, et al. Statins reduce vascular inflammation in atherogenesis: A review of underlying molecular mechanisms. *Int J Biochem Cell Biol*. 2020; 122: 105735, doi: [10.1016/j.biocel.2020.105735](https://doi.org/10.1016/j.biocel.2020.105735), indexed in Pubmed: [32126319](https://pubmed.ncbi.nlm.nih.gov/32126319/).

Table 1. Demographic data

Characteristics	TRA group (n = 84)	dTRA group (n = 98)	Conversion group (n = 18)	P-value
Age, years, mean (SD)	67 (10)	65 (10)	63.3 (9)	0.20
Male gender, n (%)	53 (63)	63 (64)	8 (44)	0.31
BMI, kg/m ² , mean (SD)	29.4 (5.7)	29.0 (5.2)	28.6 (6.4)	0.83
Obesity, n (%)	29 (35)	30 (31)	5 (28)	0.65
Current smoking, n (%)	40 (48)	35 (36)	6 (33)	0.21
Lipid disorders, n (%)	38 (45)	44 (45)	6 (33)	0.60
Diabetes or prediabetes, n (%)	30 (36)	27 (28)	4 (22)	0.35
Hypertension, n (%)	58 (69)	73 (74)	12 (66)	0.54
CKD, n (%)	4 (5)	9 (9)	1 (6)	0.36
Medications				
ASA, n (%)	40 (48)	62 (63)	10 (56)	0.13
ADP/P2Y inhibitors, n (%)	13 (15)	23 (23)	3 (17)	0.38
NOAC, n (%)	11 (13)	12 (12)	2 (11)	0.96
Statins, n (%)	38 (45)	44 (45)	6 (33)	0.60

Abbreviations: ASA, acetylsalicylic acid; BMI, body mass index; CKD, chronic kidney disease; NOAC, novel oral anticoagulants

Table 2. Characteristics of procedures

Characteristics	TRA group (n = 84)	dTRA group (n = 98)	Conversion group (n = 18)	P-value
Time needed to gain vascular access, seconds, mean (SD)	50 (4) ^a	81 (8)	277 (51) ^{b, c}	<0.001
Hematoma after procedure, n (%)	5 (6)	12 (12)	4 (22)	0.09
Radial artery occlusion after procedure, n (%)	2 (2)	3 (3)	1 (6)	0.78
VAS 1 score, median (IQR)	2 (2–4)	4 (2–5)	4 (2–5)	0.04
VAS 2 score, median (IQR)	2 (1–4)	2 (1–4)	2 (2–4)	0.57
Revascularization with stent implantation, n (%)	34 (40)	40 (41)	5 (28)	0.57
Radial dose during procedure, mGy, mean (SD)	958 (115)	888 (79)	630 (161)	0.37
Amount of contrast during procedure, ml, mean (SD)	110 (6.6)	117 (6.8)	101 (12.4)	0.50

^aTRA vs. dTRA, $P = 0.04$. ^bdTRA vs. Conversion, $P < 0.001$. ^cTRA vs. Conversion, $P < 0.001$

Abbreviations: TRA, traditional transradial access VAS 1, pain at the time of gaining vascular approach; VAS 2, pain during the maintenance of pressure dressing



Figure 1. Distal transradial access



Figure 2. Traditional transradial access




Figure 3. Biochemical evaluation of endothelial markers depending on the approach, 15 minutes after removal of hemostatic compression




Figure 4. Interleukin-8 concentration regardless the approach, 15 minutes after removal of hemostatic compression