



Clinical significance of plasma candidate biomarkers of Alzheimer's Disease

Piotr Lewczuk^{1,2}, Marta Łukaszewicz-Zajac³, Johannes Kornhuber¹, Barbara Mroczo^{2,3}

¹Department of Psychiatry and Psychotherapy, Universitätsklinikum Erlangen, and Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, Germany

²Department of Neurodegeneration Diagnostics, Medical University of Białystok, Białystok, Poland

³Department of Biochemical Diagnostics, Medical University of Białystok, Białystok, Poland

ABSTRACT

The number of patients with Alzheimer's Disease (AD) has increased rapidly in recent decades. AD is a complex progressive neurodegenerative disease affecting c.14 million patients in Europe and the United States. The hallmarks of this disease are neurotic plaques composed of the amyloid- β (A β) peptide and neurofibrillary tangles formed of hyperphosphorylated tau protein (pTau). To date, four CSF biomarkers: amyloid beta 42 (A β 42), A β 42/40 ratio, Tau protein, and Tau phosphorylated at threonine 181 (pTau181) have been validated as core neurochemical AD biomarkers. Imaging biomarkers are valuable for AD diagnosis, although they suffer from limitations in their cost and accessibility, while CSF biomarkers require lumbar puncture. Thus, there is an urgent need for alternative, less invasive and more cost-effective biomarkers capable of diagnosing and monitoring AD progression in a clinical context, as well as expediting the development of new therapeutic strategies. This review assesses the potential clinical significance of plasma candidate biomarkers in AD diagnosis. We conclude that these proteins might hold great promise in identifying the pathological features of AD. However, the future implementation process, and validation of the assays' accuracy using predefined cut-offs across more diverse patient populations, are crucial in establishing their utility in daily practice.

Keywords: Alzheimer's Disease, amyloid beta, tau, plasma biomarkers

Introduction

Alzheimer's Disease (AD) is a progressive neurodegenerative disorder that leads to a gradual decline in cognitive abilities and behavioural changes. Currently, more than 55 million people have dementia worldwide, and every year there are nearly 10 million new cases. The cumulative expenses linked to healthcare, long-term care, and hospice care for AD and other forms of dementia was estimated in 2019 to cost economies globally 1.3 trillion US dollars [1]. Furthermore, given that AD occurrence escalates with age, the prevalence of the disease is inexorably on the rise as the population ages. An estimated 44 million individuals worldwide currently live

with dementia, a figure projected to triple by 2050 owing to an ageing population. The most significant increase in dementia prevalence is anticipated in low and middle-income countries [1–3]. The onset of AD is associated with pathological changes that begin in the medial temporal lobe and then spread to areas of the neocortex, and these changes start decades before any clinical symptoms appear [3, 4].

AD progresses through three stages: a pre-symptomatic phase, a prodromal phase marked by cognitive symptoms, and finally a symptomatic phase leading to dementia [5]. Additionally, mild cognitive impairment (MCI) is often a precursor to cognitive dysfunction in AD dementia. Each year, c.10–20% of individuals with MCI will develop AD [6].

Address for correspondence: Prof. Piotr Lewczuk, Department of Psychiatry and Psychotherapy, Lab for Clinical Neurochemistry and Neurochemical Dementia Diagnostics, Schwabachanlage 6, 91054 Erlangen, Germany; e-mail: Piotr.Lewczuk@uk-erlangen.de

Submitted: 13.05.2024 Accepted: 21.06.2024 Early publication date: 02.08.2024

This article is available in open access under Creative Commons Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

The clinical presentation of AD typically follows a predominantly symptom-free preclinical phase, making early diagnosis especially challenging [5–8]. Therefore, early detection of AD is crucial for effective treatment.

The pathophysiology of AD is centred on the buildup of amyloid beta ($A\beta$) plaques and neurofibrillary tangles formed by Tau fibrils, alongside glial activation, neuronal and synaptic degeneration, and neuroinflammation. The presence of extracellular senile plaques made up of $A\beta$ peptides and intracellular neurofibrillary tangles that contain a hyperphosphorylated form of Tau (pTau) proteins are two of the most frequent neuropathological characteristics observed in the brains of patients with AD. It is widely accepted that the most reliable biomarkers of AD are these two categories of molecules [5–8].

Cerebrospinal fluid (CSF) biomarkers

$A\beta$ peptides

Amyloid plaques consist of peptides formed from the enzymatic breakdown of β -amyloid precursor protein (APP). Consequently, the generation of $A\beta$ occurs through the successive cleavage of APP by β -site amyloid precursor protein cleaving enzyme 1 (BACE1) and γ -secretase, resulting in the liberation of multiple forms of $A\beta$ peptides, the most abundant being 40 amino acids in length ($A\beta_{40}$) [9]. Furthermore, α -secretase may also process APP, leading to the release of soluble APP α via a pathway that does not produce amyloid. Moreover, the cleavage site of γ -secretase within the transmembrane domain of APP lacks precision, leading to the production of $A\beta$ peptides varying in length [9, 10]. AD-associated $A\beta$ is present in the central nervous system (CNS) in many different isoforms, having both N- and C-terminal variants. The most hydrophobic and longest $A\beta$ consists of 42 amino acids ($A\beta_{42}$) and is the major constituent of senile plaques in AD brains. $A\beta_{42}$ is also present in cerebrospinal fluid (CSF), but at a lower concentration compared to the shorter and more hydrophilic $A\beta_{40}$ and $A\beta_{38}$ isoforms [10].

Some studies have shown that the levels of $A\beta_{42}$ in CSF are inversely related to the amount of plaque in the brain. This has been observed both through *in vivo* imaging with positron emission tomography (PET) and in post mortems [11]. The reduction in CSF $A\beta_{42}$ levels among AD patients has been confirmed by numerous studies, consistently demonstrating an average fold change of 0.56 for CSF $A\beta_{42}$ compared to cognitively intact older individuals [12]. It has been proved that the measurement of CSF $A\beta_{42}$ is characterised by sensitivity of 78% and specificity of 81% in distinguishing AD patients from elderly controls [13]; similar results have been presented by other authors [14]. Moreover, it has been estimated that CSF $A\beta_{42}$ measurements aid in the accurate classification of 87% of individuals when distinguishing between non-Alzheimer's (non-AD) dementia patients and non-demented individuals [15]. Furthermore, it is suggested that a decrease in CSF

$A\beta_{42}$ serves as an early indicator of clinically 'silent' brain amyloidosis [16].

Several investigators have shown improved diagnostic accuracy of CSF $A\beta_{42}/A\beta_{40}$ ratio compared to $A\beta_{42}$ alone [17]. The usefulness of $A\beta_{42}/A\beta_{40}$ values across the spectrum of AD can be grouped into three main categories. The first category includes diagnostic studies for AD, which use clinical diagnosis as a reference (case control design) and compare their results to amyloid PET as an indicator of AD pathology. The second category focuses on differential diagnosis between AD and other types of neurodegenerative disorders. The third category comprises prognostic studies that test the ability of the $A\beta_{42}/A\beta_{40}$ ratio to predict progression from pre-clinical to dementia stages of the disorder.

CSF biochemical markers such as $A\beta_{42}$, total Tau (tTau) and Tau phosphorylated at threonine 181 (pTau181) have proven diagnostic accuracy for MCI and dementia due to AD [7, 17–18]. It is important to be able to distinguish between AD and other types of dementia. Previously, we compared the accuracy of CSF $A\beta_{42}$ and $A\beta_{40}$ as well as tTau concentrations in discriminating between patients with AD, non-AD, and control subjects using the enzyme-linked immunosorbent assay (ELISA) method [17].

The concentrations of CSF $A\beta_{42}$ have been shown to be significantly lower, while CSF tTau levels are significantly higher, in AD patients compared to controls and non-AD [17]. CSF levels of $A\beta_{40}$ did not differ significantly among the analysed groups. Moreover, receiver operating characteristic (ROC) analysis was assessed to define cut-off values for maximised sensitivity and specificity. For all analysed groups, the $A\beta_{42}/A\beta_{40}$ ratio classified more patients correctly than did the concentration of $A\beta_{42}$ alone (AD vs controls, 94% and 86.7%; AD vs non-AD, 90% and 85%; and AD vs non-AD and controls, 90.8 and 87%) [17]. In addition, the percentage of correctly classified patients was further improved when the $A\beta_{42}/A\beta_{40}$ ratio was combined with the analysis of the CSF tTau levels [17].

CSF levels of $A\beta$ peptides ending at the amino acid position of 42 are widely accepted biomarkers of AD. However, a neurochemical dementia diagnostics (NDD) interpretation of subjects with constitutively high or low CSF levels of total $A\beta$ peptides, could lead to erroneous conclusions, because these biomarkers seem to correlate better with the total $A\beta$ load than with the pathological status.

In our previous study, we reported significantly increased CSF concentrations of phosphorylated Tau (pTau181) and tTau in subjects with high CSF $A\beta_{40}$ concentrations and decreased $A\beta_{42}/A\beta_{40}$ ratio compared to those with a low CSF $A\beta_{40}$ and a normal $A\beta_{42}/A\beta_{40}$ ratio [18]. Furthermore, we found a significantly decreased $A\beta_{42}/A\beta_{40}$ ratio in the group of subjects with apolipoprotein E epsilon 4 (APOE ϵ 4) allele compared with the group of subjects without this allele. Surprisingly, patients with low $A\beta_{40}$ and decreased $A\beta_{42}/A\beta_{40}$ ratio were characterised by decreased pTau181 and unaltered tTau compared

to subjects with high A β 40 and a A β 42/A β 40 ratio within the normal range.

We suggest that the A β concentration ratio should replace the 'raw' concentrations of corresponding A β peptides to improve reliability of the neurochemical dementia diagnosis [18]. In addition, based on the body of evidence, we suggest that the CSF A β 42/40 ratio, rather than the absolute value of CSF A β 42, should be used when analysing CSF AD biomarkers to improve the percentage of appropriately diagnosed patients [7]. Furthermore, it is worth emphasising that the empirical findings of a better diagnostic performance of A β 42/40 ratio, compared to A β 42 alone, is grounded in fundamental principles of mathematical stochastics and theory of distributions, as was derived in our previous study [19]. With this derivation, it was proven that under particular conditions, a quotient of two variables (in our case, A β 42/40) has always more compacted and less dispersed distribution than has the numerator (A β 42), which leads to less overlapping of the distributions from two populations e.g. subjects with and without disease [19].

Research involving AD patients and cognitively normal individuals highlighted a strong agreement between CSF A β 42 levels and amyloid- β PET imaging [11, 20]. The concentration of CSF A β 42 decreases before the detection of amyloid- β using PET imaging, implying that A β 42 may serve as a more sensitive marker of AD during its very early stages, while A β PET could provide better at grading the severity of early AD [11, 20]. Palmqvist et al. [21] found that both amyloid PET and CSF biomarkers accurately identify early AD in patients with MCI-AD, with no enhanced accuracy when combining CSF and PET amyloid measures compared to using either CSF A β 42 or tTau alone [21]. Therefore, the decision about using CSF or A β PET biomarkers for early AD identification may currently depend on PET scanner availability, physician/patient preference, and associated costs [22, 23].

Research has been conducted to investigate the potential of various A β peptides as biomarkers for AD. Peptides longer than 42 or shorter than 40 amino acid residues have been assessed for their diagnostic accuracy. For instance, the A β 43 isomer is decreased in AD and has a similar diagnostic accuracy to that of CSF A β 42. Clinical investigations have revealed that there is no significant difference between CSF A β 38 levels in AD patients and control subjects. However, there is a correlation between CSF A β 38 and PET A β [24]. The authors concluded that the CSF A β 42/A β 38 and CSF A β 42/A β 40 ratios are better indicators than CSF A β 42 alone in detecting brain amyloid deposition in prodromal AD and in distinguishing AD dementia from non-AD dementias [25]. Additionally, the CSF A β 42/A β 38 ratio may aid in differentiating between AD and other non-AD dementias [21] e.g. dementia with Lewy bodies (DLB) [26]. It seems that an initial stage in the development of AD is the oligomerisation of A β monomers, especially those ending at the C-terminal 42. Therefore, A β oligomers may have a diagnostic role in AD [28–31], something we will look at in the next section.

Tau protein and its phosphorylated forms

Tau proteins are a type of microtubule-associated molecule that can be found in both neuronal and non-neuronal cells. There are six different isoforms of this protein, varying in length from 352 to 441 amino acid residues [32]. The primary function of Tau proteins is to promote stability and growth of neuronal microtubules. tTau protein concentration is a non-specific marker of neuronal destruction in neurodegenerative diseases. Various studies have demonstrated that tTau concentrations are elevated in the CSF of AD patients [33].

The pTau molecules play a crucial role in the regulation of Tau-microtubule interactions. It is thought that hyperphosphorylated Tau or oligomeric Tau may contribute to synaptic degeneration, while granular Tau oligomers could be responsible for neuronal loss. There is a suggestion that adding the measurement of soluble oligomers of Tau protein (TauOs) to the panel of CSF biomarkers might enhance the diagnosis of AD. It is believed that the toxicity of TauOs could be a potential factor in the pathogenesis of this disease, acting in the initial stages and seeding Tau pathology within the brains of individuals with AD. It has been identified that there are multiple phosphorylated tau residues in the mid-domain, including pTau181, threonine 231 (pTau231), serine 235 (pTau235), serine 199 (pTau199) as well as for the C-terminal residues serine 396 and 404 [34]. Studies have shown that CSF levels of pTau181, pTau199, and pTau231 are effective in distinguishing AD from other neurodegenerative disorders and non-demented controls [35]. In AD patients, CSF concentrations of pTau181 were significantly higher, especially in those with neurochemically supported clinical diagnoses with decreased A β 42 in the CSF [15]. According to a study by Parnetti et al. [36], pTau181 could potentially serve as a biomarker to differentiate between AD and DLB. Furthermore, patients with MCI who progressed to AD during the study showed elevated levels of Tau protein phosphorylated at threonine 231 and serine 235 positions [37]. It has been assessed that levels of CSF pTau217 and pTau231 show pathological changes earlier than CSF pTau181. Thus, studies have reported that CSF pTau 217 correlates more strongly with amyloid PET and tau PET positivity than other forms of tau [38,39]. Buchhave et al. [40] examined the predictive capability of CSF pTau in forecasting the development of AD within 10 years in individuals with MCI. They compared CSF biomarkers between those who converted to AD early and late during follow-up. Initially, patients who progressed to AD had notably higher pTau levels compared to nonconverters. Additionally, early converters exhibited significantly elevated pTau levels compared to late converters. Some authors claim that so-called 'Tau/A β ' quotients have the potential to upgrade diagnostic performance or, to put it more precisely, improve its interpretation. Such approaches must be taken with caution.

Firstly, in contrast to A β 42/40 ratio, which is a measure for normalisation of A β 42 for the total CSF amount of A β (of which A β 40 is the most abundant isoform), Tau/A β attempts




Figure 1. Core and novel cerebrospinal fluid (CSF) biomarkers of Alzheimer's Disease (AD) [17–19, 34–58]

to normalise a biomarker of one pathophysiological process (amyloidosis) for a biomarker of another process (neurodegeneration). This is methodologically questionable. Secondly, any quotient, by its mathematical definition, explicitly assumes that the relation of the two quantities (here, a biomarker of amyloidosis and a biomarker of neurodegeneration) is linear, which is simply incorrect [41]. An example will clarify this.

Let us imagine a patient with an early stage of disease (and note that such patients are the ultimate target population for early AD diagnostics) whose $A\beta_{42}$ just starts declining but who has constitutively low concentrations of Tau and/or other pTau. Such a patient will have a normal $A\beta$ /Tau ratio until a much later stage, when Tau starts increasing, and will be diagnosed falsely negatively. In an even worse scenario, consider a patient with physiologically low $A\beta_{42}$ (due to a low total $A\beta$ amount) who has Tau larger than the median, albeit within the normal range. Such a patient will have a decreased $A\beta$ /Tau quotient, and will be diagnosed falsely positively.

Potential further CSF biomarkers for AD

Potential CSF biomarkers have been categorised according to the underlying pathophysiological mechanisms of AD, encompassing various domains such as $A\beta$ metabolism ($A\beta_{38}$, BACE1), inflammation and glial activation (triggering receptor expressed on myeloid cells 2 – TREM2 and its soluble variant – sTREM2, chitinase-3-like protein 1 – YKL-40, vascular dysregulation (heart-type fatty acid-binding protein – hFABP), α -Synuclein pathology (α -Syn), synaptic dysfunction (neurogranin, synaptosome-associated protein 25 – SNAP-25 and synaptotagmin-1 – SYT-1), TDP-43 pathology (TAR DNA binding protein 43 – TDP-43), and other neuronal proteins (visinin-like protein 1 – VILIP-1 and neurofilament light – NfL) [42–58]. While alterations in $A\beta$ metabolism are recognised as

the earliest detectable events in AD, interventions based on the $A\beta$ hypothesis have to date yielded disappointing results [59,60]. This underlines the need for a broader exploration of alternative hypotheses, with particular interest in those related to Tau [61]. This is reinforced by the observation that cognitive symptoms in AD correlate more directly with biomarkers of neurodegeneration than with $A\beta$ deposition. Following this rationale, a novel assay designed to specifically measure non-phosphorylated forms of Tau molecules (Non-pTau) has been developed [62]. Intriguingly, it significantly improved the accuracy of patient classification (99%) compared to routinely used assays: Tau (90%), pTau181 (62%) [63]. Additionally, the Non-pTau assay is extensively employed in the differential diagnosis of other dementias, particularly those with substantial Tau pathology, although definitive conclusions have yet to be drawn [62–63]. Well-established, and candidates for novel, CSF biomarkers of AD are set out in Figure 1.

Compared to non-soluble forms in $A\beta$ plaques, soluble $A\beta$ oligomers ($A\beta$ Os) are more toxic [28]. Clinical investigations have demonstrated a notable increase in CSF $A\beta$ Os and/or the $A\beta$ Os/ $A\beta_{42}$ ratio in AD patients compared to age-matched controls, along with an inverse correlation between $A\beta$ O levels and mini-mental state examination (MMSE) score, suggesting the significance of oligomers as a diagnostic marker for AD [29]. Authors have concluded that $A\beta$ Os could serve as a test to differentiate between AD and MCI patients and cognitively normal controls, suggesting that elevated $A\beta$ O levels may predict progression from MCI to AD [30] and might serve as a potential biomarker for AD diagnosis, with diagnostic sensitivity and specificity of more than 95% and 90% respectively [31].

Many papers have indicated that total α Syn levels tend to be higher in AD patients than in controls [64–67]. The increased levels of CSF α Syn in AD and the moderate correlation

between α Syn and Tau/pTau181 support the findings of previous research suggesting that increased CSF α Syn in AD is due to general neurodegeneration, rather than any process specific to AD. As the primary source of α Syn in the brain is presynaptic neuronal terminals, it seems reasonable to suggest that degenerating neurons release α Syn molecules, which can then diffuse into the CSF at a higher rate. Further studies are needed to determine the relationship between biomarkers and clinical presentation, such as cognitive measures, as well as the impact of patient variables such as sex, APOE ϵ 4 status, and comorbidities [66, 67]. This is currently being looked at for several other candidates, such as selected metalloproteinases (MMPs) and their tissue inhibitors [56]. For instance, compared to elderly individuals without cognitive deficits, AD patients have significantly lower CSF concentrations of matrix metalloproteinase 9 (MMP-9), and significantly higher levels of matrix metalloproteinase 3 (MMP-3). These biomolecules might contribute to the pathophysiology and diagnosis of AD, indicating the need for further studies involving larger patient cohorts to determine their potential diagnostic value [56].

The Erlangen Score

The combination of reduced CSF A β 42 levels and/or A β 42/40 ratio, along with elevated levels of Tau and/or pTau, as discussed previously, signifies the dual pathological processes of AD i.e. amyloidosis and neurodegeneration. Despite the high accuracy of these CSF biomarkers for diagnosis, their global acceptance has been hampered by challenges related to result comparability across different laboratories (or even within the same lab) using different analytical methods. Efforts have been made to standardise procedures for sample collection, measurement protocols, and assay calibrators to address this issue, but widespread adoption of these new approaches will require time [68–71]. Furthermore, with the increasing use of AD CSF biomarkers in routine clinical settings, interpreting the results requires expertise and caution. The challenge lies in deciphering the frequently heterogeneous information these biomarkers provide, which might not always fit neatly into clear normal/abnormal categories. Various approaches have been proposed to standardise the diagnostic interpretation of CSF biomarker profiles. The ES interpretation algorithm was introduced in 2009 [71], and this has been followed by other methods such as logistic regression models, classification scales like the Paris-Lille-Montpellier (PLM) scale [72], or a descriptive nominal-scale A/T/N system [73].

Currently, the Erlangen Score provides a system based on CSF biomarkers, but its flexibility allows its easy adaptation to other biomarkers e.g. those derived from the blood or neuroimaging [71]. The sole prerequisite is that there is at least one biomarker of amyloidosis and at least one biomarker of neurodegeneration, and that the biomarkers are validated in terms of a centre-specific reference range. A score of 0 is given

if all biomarkers are normal, which equates to ‘no neurochemical evidence for AD’. If there are border zone alterations in one biomarker group (either A β or Tau/pTau but not both), the score is 1, and this is referred to as ‘neurochemically improbable AD’. A score of 2 is given if there are evident alterations in either A β metabolism (decreased A β 42 concentration or A β 42/A β 40 ratio) or tau metabolism (increased concentrations of Tau and/or pTau181), but not in both, or if there are border zone alterations in the CSF biomarkers of both groups. When there are noticeable changes in either A β or Tau biomarkers group along with border zone changes in the other group, a score of 3 points is assigned. These cases, when ES = 2 or 3, are referred to as ‘neurochemically possible AD’ (see Table 1). If there are noticeable changes in both A β and Tau groups, it results in 4 points, known as ‘neurochemically probable AD’. In cases where there is isolated, very high concentration of tTau, it is interpreted as suspected rapidly progressing neurodegeneration with unlikely AD. However, if this same concentration of Tau is accompanied by pathological A β 42 concentrations/ratio, then the interpretation would shift to possible or even probable AD depending on the normality of pTau. The ES pattern can be presented graphically for clinicians to review.

This concept offers distinct advantages over alternative methods. It facilitates the classification of CSF results into five classes on a graded scale (0-4), indicating increasing deviations in AD CSF biomarkers. Additionally, it introduces the novel concept of border zone results, enhancing the interpretation of CSF AD biomarkers. ES is straightforward and does not require computer-based support in routine laboratory operations, although it can be easily adapted for automated systems in high-throughput labs. In contrast to the A/T/N classification, ES categorises subjects on a graded scale, facilitating semi-quantitative correlation of CSF findings with other metrics such as progression hazards, odds ratios, or time to progression from MCI to dementia. Moreover, as an ordinal-scale system, ES accommodates border-zone laboratory results, seamlessly integrating them into the interpretation algorithm. Furthermore, compared to the PLM approach, which focuses on the number of abnormal CSF biomarkers [72], ES is more adaptable, allowing for the inclusion of additional potential biomarkers reflecting amyloid pathology or neurodegeneration without the need to adjust the categories. Regardless of the number of biomarkers considered, ES consistently categorises CSF patterns into five ordinal categories. ES has undergone thorough validation using cohorts from various expert centres and a wide range of predefined endpoints. Initially, ES demonstrated accurate classification of individuals with non-demented/mild cognitive impairment who were at higher risk of dementia development in two separate, large-scale, multicentre studies (the German Competence Network Dementias and the US-ADNI). Notably, these studies used distinct sample handling protocols, laboratory analytical platforms, and centre-specific reference ranges. Nonetheless, ES exhibited consistent performance across both [71, 74].

Table 1. Interpretation of Erlangen Score [71]

Score	Interpretation of amyloid β results	Interpretation of Tau/ pTau results	Interpretation of Erlangen Score
0	A β 42 — normal A β 42/A β 40 ratio — normal	Tau — normal pTau — normal	No neurochemical evidence for AD
1	Levels in border zone alterations in one biomarker group (either A β or Tau/pTau, but not both)		Neurochemically improbable AD
2	Evident alterations in either A β metabolism (decreased A β 42 concentration or A β 42/A β 40 ratio) or tau metabolism (increased concentrations of Tau and/or pTau181) but not both, or if there are border zone alterations in CSF biomarkers of both groups		Neurochemically possible AD
3	Notable changes in either A β or Tau biomarkers group along with border zone changes in other group		Neurochemically possible AD
4	Notable changes in both A β and Tau groups		Neurochemically probable AD

Note that this does not cover cases with extremely large concentrations of Tau, which may point at rapidly progressing neurodegenerative conditions like Creutzfeldt-Jakob Disease rather than at AD, and which need special consideration

A separate investigation revealed that individuals diagnosed with MCI and categorised as ‘neurochemically probable AD’ had an 8-12 times greater risk of dementia onset compared to those classified as ‘neurochemically improbable AD’, even after accounting for age, gender, MMSE score, and APOE ϵ 4 genotype. Importantly, these hazard ratios remained consistent regardless of time. Conversely, the ES fully accounted for the risks associated with demographic, cognitive, and genetic factors [75]. When examining neuropsychological and neuroimaging outcomes, it has been observed that a greater ES correlates with accelerated disease progression in individuals with MCI. Those with higher ES scores exhibit swifter declines in both whole brain and hippocampal volumes, as well as more rapid decreases in MMSE scores [76].

Ultimately, the ES algorithm has facilitated accurate prediction of post-mortem neuropathological outcomes based on *in vivo* CSF results of three core AD biomarkers. The likelihood of having AD pathology post mortem, as opposed to non-AD pathologies such as DLB, vascular dementia (VaD), or frontotemporal dementia (FTLD), increases nearly linearly with higher ES ordered categories. Interestingly, less than 3% of neuropathologically confirmed AD patients (3/106) were classified as ‘probable AD’ (ES = 0 or 1) [77].

It has been proved that neuropathological assessment of the amount and distribution of plaques and neurofibrillary tangles is the most crucial criterion for the diagnosis of AD [78, 79]. In daily practice, PET and the analysis of CSF biomarkers have been widely used to detect and monitor AD-related amyloid and tau pathologies [80]. However, recent investigators have focused on developing blood tests for AD because blood sample collection is less invasive and more cost- and time-effective.

Therefore, the aim of the next section is to critically evaluate the potential significance of plasma candidate biomarkers, such as plasma amyloid beta (A β), plasma pTau, inflammatory plasma candidate biomarkers including YKL-40, glial fibrillary acidic protein (GFAP), monocyte chemoattractant protein-1 (MCP-1) and eotaxin-1, as well as other plasma candidate biomarkers of AD (e.g. NfL) in the diagnosis of AD.

Many investigators have confirmed that plasma candidate biomarkers that would be useful in early diagnosis, stratification, prediction of disease course, or monitoring response to therapy in AD are sorely needed. However, technical challenges with the measurement of these molecules in the blood, and a lack of validation and cutoff values, have limited their use in daily practice. Recent technological advances have improved assay sensitivity, delivering ultrasensitive assays capable of measuring specific plasma biomarkers. The potential significance of plasma candidate biomarkers of AD is set out in Table 2.

Plasma candidate biomarkers for AD

Plasma amyloid beta (A β)

First of all, it needs to be emphasised that currently (June 2024) there is no validated ‘plasma biomarker’ that could be used for an individual-level diagnostic. Neurochemical assessment of AD patients relies on identifying pathology through A β aggregates via brain scans or CSF analysis, typically performed at specialised medical centres. Consequently, the quest for blood biomarkers accurately reflecting abnormal A β accumulation is paramount for enhancing AD diagnosis [81–83]. The recent focus has been on blood-based biomarkers facilitating early AD detection via cost-effective, minimally invasive methods. However, plasma A β concentrations are notably lower (50–100 times) than in CSF [84–88]. Due to insufficient precision and conflicting outcomes, distinguishing between an AD and a non-AD individual poses significant challenges [89]. Consequently, research is increasingly focusing on correlating plasma AD biomarkers with PET imaging, core CSF biomarkers, and cognitive staging.

In recent years, research into mass spectrometry (MS) has witnessed significant advances in sensitivity and precision, enabling the detection of protein concentrations at femtomolar levels with a coefficient of variation (CV) of below 4% [84]. Consequently, numerous investigators have used immunoprecipitation-mass spectrometry (IP-MS) to assess plasma A β levels, aiming to differentiate between individuals with

Table 2. Potential significance of plasma candidate biomarkers of AD

Pathophysiological mechanism	Plasma biomarker	Alterations in AD	References
A β metabolism	A β 42	– significantly lower in amyloid PET-positive individuals than in PET negative cases	[26, 87, 94–95]
	A β 42/40 ratio	– significantly lower in MCI and AD cases compared to cognitively healthy controls – best biomarker for discriminating A β -positive vs. A β -negative individuals	
Intracellular neurofibrillary tangles formed by hyperphosphorylated form of Tau (pTau) proteins	pTau181	– higher in AD patients compared to CU and strongly correlated with both A β and Tau PET – strong correlation with CSF pTau181 – differentiated AD dementia from non-AD neurodegenerative diseases with an accuracy similar to that of Tau PET and CSF ptau181	[107–111]
	pTau217	– differentiated clinically diagnosed AD dementia from non-AD neurodegenerative disorders, with accuracy comparable to CSF pTau and tau PET – significantly better performance than all other plasma pTau biomarkers when detecting abnormal A β status or progression to Alzheimer's dementia – more useful biomarker than tau PET in earliest stages of AD	[78, 114, 116]
	pTau231	– distinguished AD patients from patients with non-AD neurodegenerative disorders as well as from A β -negative MCI patients	[94, 118]
Inflammation/microglial activation	chitinase-3-like protein 1 (YKL-40)	– significantly different levels between preclinical AD based on pTau181/A β 1-42 ratio and healthy elderly individuals	[127]
	glial fibrillary acidic protein (GFAP)	– significantly elevated in patients with preclinical AD compared to healthy elderly – significantly higher in A β -positive group compared to A β -negative group – correlated with increased risk of progression to dementia and steeper cognitive decline (valuable prognostic biomarker predicting incident dementia)	[127, 130, 132]
Neuronal proteins	Neurofilament light (NfL)	– increased levels in MCI and AD A β -positive individuals compared to cognitively unimpaired and AD A β -negative group – strong correlation with CSF levels – significant increase in AD cases, comparable to core AD CSF biomarkers – elevated levels in MCI cases with positive amyloid PET scans – screening test at first clinical evaluation of patients with cognitive disturbance – novel biomarker for early neurodegeneration in AD	[122, 123, 145] [16, 135, 139, 143, 144, 146]

amyloidosis and those without, with reference to amyloid PET and CSF A β measurements [84, 86]. The IP-MS approach employs selected reaction monitoring (SRM) for plasma A β 42 and A β 40 quantification, incorporating stable isotope-labelled A β peptides into samples prior to analysis (simultaneous assessment), and using octyl glucopyranoside detergent to disrupt A β -protein complexes in plasma [90]. Ovod et al. employed an IP-MS technique involving LysN proteolytic digestion of A β peptides prior to analysis. They found that plasma A β 42 levels and the A β 42/40 ratio were notably lower in individuals with positive amyloid PET scans compared to those with negative scans [87]. Furthermore, the plasma A β 42/40 ratio exhibited a 14% decrease in the amyloid PET-positive group compared to amyloid PET-negative participants, demonstrating a high ROC value of 0.89 [87]. Several clinical studies have suggested that currently available plasma A β 42/40 tests using IP-MS technologies could serve as valuable screening tools, potentially reducing the necessity for amyloid PET scans in c.49–64% of patients [91–93]. Additionally, researchers have also observed a correlation between the plasma levels of A β 42/40 [91, 94–97] and cerebral amyloidosis detected by PET imaging, with the area under the ROC curve (AUC) ranging from 0.7–0.8 or

even higher in studies incorporating age and APOE genotype into the analysis [94].

In the extensive Swedish BioFINDER study cohort, significant correlations were observed between plasma A β 42 levels, the A β 42/40 ratio, and corresponding CSF biomarkers, as well as cortical 18 F-florbetapir amyloid PET retention. These findings were achieved using an ultra-sensitive immunoassay platform (SIMOA, single molecule array for protein detection), which employs a single-molecule array based on immunocapture of protein biomarkers on magnetic beads [98]. Furthermore, markedly lower plasma A β 42/40 ratios were detected in individuals with both MCI and AD compared to cognitively healthy controls [98]. Smirnov et al. similarly reported the relationship between plasma AD biomarkers and post mortem brain findings, employing the SIMOA platform to validate longitudinal plasma candidate biomarkers against PM-confirmed diagnoses, and they investigated specific associations with AD. Notably, subjects with significant amyloid AD pathology were shown to exhibit lower plasma A β 42/40 levels, with a diagnostic accuracy of AUC = 0.601 [94].

Keshavan et al. conducted a comparative analysis of various blood-based methodologies, including liquid

chromatography mass spectrometry (LC-MS) and SIMOA assays, to ascertain plasma A β levels and their ability to detect cortical 18 F-florbetapir amyloid PET positivity in individuals without dementia [95]. Thus, they assessed the potential utility of blood biomarkers as predictors of amyloid PET status using logistic regression models [95]. The authors found that while the AUC for amyloid PET status using a base model incorporating age, sex, and APOE ϵ 4 status was 0.695, the two most effective SIMOA plasma candidate biomarkers were A β 42/40 (AUC = 0.620) and phospho-tau181 (AUC = 0.707), neither of which surpassed the base model. In contrast, MS plasma measures such as A β 42/40 (AUC = 0.817) and A β composite (AUC = 0.820) performed significantly better. Additionally, MS measures of A β 42/40 exhibited 86.6% sensitivity and 71.9% specificity in detecting amyloid PET positivity.

These findings suggest that the LC-MS platform is superior to SIMOA assay for plasma A β and pTau181 analysis when screening a large population of dementia-free individuals for concurrent PET amyloid status. Furthermore, the authors propose that plasma screening could reduce the need for amyloid PET scans in identifying A β -positive individuals for clinical trial recruitment or anti-amyloid therapy administration, which may be feasible even in preclinical cohorts [95].

Palmqvist et al. used fully automated plasma assays, specifically Elecsys prototype immunoassays, to evaluate plasma A β 42/A β 40 levels across cohorts comprising cognitively unimpaired (CU), MCI, and AD individuals [81]. The authors found that the most effective biomarker for distinguishing A β -positive from A β -negative individuals was plasma A β 42/A β 40 ratio, with AUC values ranging from 0.83 to 0.87. Additionally, the combined measurement of plasma A β 42/A β 40, pTau181, and ApoE4 significantly enhanced the AUCs to 0.90–0.93 [81]. These findings suggest that a fully automated instrument capable of evaluating a combination of three biomarkers accurately identified A β positivity in two independent cohorts, effectively predicting the future development of AD dementia [81]. The authors proposed that blood-based biomarkers have the potential to enhance AD diagnosis, facilitate recruitment for AD trials, and monitor anti-A β therapies due to their high precision combined with high accuracy. Nonetheless, they emphasise that in the clinical implementation process, validation of the assays' accuracy and robustness using predefined cut-offs across more diverse patient populations is crucial to further establish the utility of plasma candidate biomarkers in daily practice [81].

The findings regarding plasma A β 42 as a biomarker reflecting brain amyloid pathology present conflicting outcomes, attributable to the overlap in plasma A β 42 and A β 40 levels between AD patients and non-AD controls as assessed by various techniques [12]. This discrepancy may stem from contributions by peripheral tissues to plasma A β , leading to a lack of correlation between plasma and CSF A β concentrations [99]. Furthermore, potential interference could be mitigated through analytical enhancements like standard

immunoassays [100]. Nonetheless, the reported findings are promising, underlining the importance of further investigations into plasma A β as a screening tool for brain amyloidosis and AD, particularly on larger clinical cohorts to compare different analytical platforms for measurement. The authors conclude that the high accuracies observed for A β pathology and future AD dementia using fully automated instruments offer promising prospects for integrating plasma candidate biomarkers into both clinical trials and clinical practice [84].

Plasma Phospho-Tau

CSF pTau and tau PET are established biomarkers indicative of AD-related tau pathology, showing distinct alteration patterns throughout AD progression. CSF pTau levels reflect changes in tau metabolism within the brain, with elevated concentrations observed across all stages of AD, particularly increasing in the earliest stages, whereas tau PET remains unaltered during the asymptomatic phase [101-103]. Conversely, tau PET tracers bind to insoluble paired helical tau filaments in neurofibrillary tangles, exhibiting abnormal levels primarily in symptomatic AD and correlating with brain atrophy and cognitive decline [104-105]. These findings suggest that fluid-based assessment of pTau may offer greater sensitivity than tau PET in detecting early AD stages.

Furthermore, increased plasma tau levels in AD patients have been identified using MRI and SIMOA techniques in extensive studies conducted on the ADNI and BioFINDER cohorts. The authors uncovered significant correlations between plasma tau levels and future cognitive decline, as well as increased atrophy measured by magnetic resonance imaging (MRI) and hypometabolism measured by fluorodeoxyglucose PET (FDG-PET) during a longitudinal follow-up [106]. A growing body of research suggests that blood pTau holds promise as both a diagnostic and a prognostic biomarker for AD. While CSF pTau181 has been widely validated and accepted as a core AD biomarker, recent studies have highlighted the potential of elevated plasma pTau181 in distinguishing AD patients with dementia from CU individuals [107-108]. Using the SIMOA technique, Mielke et al. demonstrated that plasma tTau and pTau181 levels were significantly higher in AD dementia patients compared to CU individuals, showing strong correlations with both A β and tau PET [107]. They proposed that plasma pTau181 might offer greater sensitivity and specificity as a predictor of elevated brain A β than tTau, serving as a valuable biomarker for AD pathophysiology and a non-invasive screening tool for elevated brain A β [107]. Similarly, Coolmans et al. used the SIMOA platform to compare plasma pTau181 and tau PET in predicting cognitive stage, preclinical A β status, and cognitive functioning [110]. They found comparable high AUC values for plasma pTau181 and tau PET in discriminating preclinical A β status, although tau PET outperformed plasma pTau181 in distinguishing MCI/AD from subjective cognitive decline individuals. Tau PET showed stronger correlations with cognitive decline

and a wider range of cognitive tests compared to plasma pTau181. Furthermore, while both plasma pTau181 and tau PET increased more steeply over time in MCI/AD compared to individuals with subjective cognitive decline, only tau PET annual changes were associated with cognitive decline. This may indicate its superiority in monitoring disease stage and clinical progression [110]. Moreover, Janelidze et al. revealed elevated plasma pTau181 concentrations in preclinical AD, further increasing at MCI and dementia stages, predicting positive Tau PET scans [111]. They observed a correlation between plasma and CSF pTau181, and reported that plasma pTau181 differentiated AD dementia from non-AD neurodegenerative diseases with high accuracy, like Tau PET and CSF pTau181. Elevated plasma pTau181 has also been associated with subsequent development of AD dementia in CU and MCI subjects [111]. The authors concluded that plasma pTau181 serves as a non-invasive diagnostic and prognostic biomarker for AD, with potential utility in clinical practice, trials, and predicting future progression to AD dementia in subjects without dementia [111]. Furthermore, plasma pTau181 effectively distinguishes subjects with abnormal A β and tau PET scans and differentiates AD dementia from non-AD neurodegenerative disorders [111].

Recent studies focusing on multiple phosphorylation sites of the tau protein have suggested that CSF tau phosphorylated at threonine 217 (pTau217) may provide a more accurate reflection of AD-related tau pathology compared to pTau181 [112, 113]. Palmqvist et al. demonstrated that plasma levels of pTau217 show changes concurrently with CSF levels, thereby better distinguishing neuropathologically confirmed AD subjects from those without neuropathological evidence of AD compared to plasma pTau181 [114]. Additionally, the authors reported that plasma pTau217 effectively differentiated clinically diagnosed AD dementia from non-AD neurodegenerative disorders, with an accuracy comparable to that of CSF pTau and tau PET [118]. Moreover, they concluded that plasma pTau217 levels begin to rise c.20 years before the onset of MCI in autosomal-dominant AD cases [114]. In their study, Palmqvist et al. compared currently available methods for determining pTau in the blood to ascertain which methods are sufficiently accurate for implementation in clinical practice [114]. Plasma pTau181, pTau217, and pTau231 levels were evaluated using immunoassays to assess abnormal brain A β status and predict future progression to AD [116]. The authors noted that MS pTau217 exhibited significantly better performance than all other plasma pTau biomarkers in detecting abnormal A β status (AUC = 0.947) or predicting progression to AD (AUC = 0.932) [116]. Despite variations in the performance of plasma pTau assays, these findings underline relatively high and consistent accuracy across several pTau immunoassays, emphasising their potential clinical utility [116].

Clinical investigations using immunoassay technology such as Meso Scale Discovery (MSD) have unveiled associations between plasma pTau217 and tau PET signals in one

of the earliest regions of AD-related tau pathology in CU individuals. Through the measurement of changes in plasma pTau217, CSF pTau217, and various tau PET measures [78], researchers have assessed the correlation between plasma pTau217 and longitudinal changes in tau PET in CU and MCI patients. They observed that plasma levels of pTau217 exhibit changes early in AD compared to well-established CSF and PET biomarkers of AD pathology. Specifically, pTau217 levels were elevated in CU participants with abnormal A β -PET, but normal tau PET, in the entorhinal cortex and increased in the early preclinical stages of AD when insoluble tau aggregates were not yet detectable by tau PET. Moreover, similarly to CSF pTau217, the non-invasive and cost-effective nature of plasma pTau217 suggests its potential as a more practical biomarker than tau PET in the earliest stages of AD. The authors conclude that plasma pTau217 holds promise as a biomarker for early AD brain pathology, and could serve as a valuable tool for individual selection and as an outcome measure to monitor drug responses in clinical trials involving individuals with preclinical AD [78].

In Palmqvist et al.'s study, a comparison between plasma pTau181 and pTau217 assays using Elecsys prototype immunoassays revealed interesting insights [81]. The pTau217 N-terminal assay demonstrated higher accuracies for A β positivity than the mid-domain assay. However, it is noteworthy that in many cases, concentrations of the analysed molecules fell below the lower level of detection. Despite previous studies indicating better performance of pTau217 in identifying AD pathology, the Elecsys prototype immunoassay for pTau181 appears to be more suitable than pTau217 [82, 117–121]. Therefore, further studies are imperative to validate these findings [81].

Chatterjee et al. assessed plasma tTau, pTau181, and pTau231 levels using SIMOA assays in CU older adults, stratified based on the absence (A β -) or presence (A β +) of brain amyloidosis [122]. Their findings revealed higher plasma levels of tTau, pTau181, and pTau231 in A β + CU individuals compared to A β - CU individuals. Moreover, longitudinal analyses showed an increase in pTau181 levels in A β + CU participants over 12 months. Additionally, correlations were observed between pTau181 and pTau231 levels with cognition, although no significant associations were found with hippocampal volume [122]. The authors underlined the diagnostic and longitudinal monitoring potential of pTau for preclinical AD [122]. Furthermore, while pTau181 demonstrated equivalent or superior performance compared to other biomarkers in predicting A β status, a combination of biomarkers may offer enhanced predictive capability across the AD continuum [123]. Similarly, Ashton et al. assessed plasma pTau231 concentrations using the SIMOA assay and highlighted its high diagnostic value based on tau PET, CSF amyloid and tau classification, as well as findings from post mortems [118]. Plasma pTau231 effectively identified AD patients, and differentiated them from A β negative CU older adults, with high accuracy (AUC = 0.92–0.94). Additionally,

this biomarker distinguished AD patients from individuals with non-AD neurodegenerative disorders (AUC = 0.93) and A β negative MCI patients (AUC = 0.89) [118]. Furthermore, plasma pTau231 demonstrated the ability to differentiate subjects across the Braak stage spectrum, unlike plasma pTau181. The authors concluded that novel plasma pTau231 assay identifies the clinical stages of AD and increases earlier, even in the presence of subtle A β deposition, prior to reaching the threshold for amyloid- β PET positivity, and in response to early brain tau deposition.

Thus, plasma pTau231 could be a promising biomarker of emerging AD pathology with the potential to facilitate clinical trials targeting vulnerable populations below the PET threshold of amyloid- β positivity or apparent entorhinal tau deposition [118]. Similar findings were reported by Smirnov et al., who also observed that plasma pTau231 levels appear to increase earlier than plasma pTau181 and could be used to determine the severity of tau pathology. The investigators demonstrated that plasma pTau231 exhibited an earlier increase during intermediate stages of neuritic plaque pathology compared to plasma pTau181 [94].

Investigators have determined plasma pTau181, pTau217, pTau231 compared to tau PET in individuals from memory clinics with subjective cognitive decline, CU status, or dementia. Plasma pTau217, characterised by 95% sensitivity, resulted in nearly halving the number of tau PET scans needed. The investigators concluded that plasma pTau217 could effectively guide the selection of patients for tau PET scans [124].

Plasma markers of inflammation as candidate biomarkers of Alzheimer's Disease

The neuroinflammatory response of the immune system can potentially promote protein aggregation and interact with pattern recognition receptors on microglia and astroglia, leading to the initiation of an immune response characterised by the release of inflammatory mediators [125, 126]. The proteins involved in the neurodegeneration process are responsible for the development of AD [125]. Some authors have explored plasma levels of biomarkers associated with neuroinflammation in a cohort of preclinical AD patients, comparing these levels to those in healthy elderly individuals defined by A β 42 CSF status. Several clinical studies have suggested that certain inflammatory biomarkers in plasma, such as GFAP, YKL-40, MCP-1, and eotaxin-1, could potentially identify preclinical AD subjects at high risk of developing AD [127].

It has been suggested that YKL-40 is a more sensitive marker of the incipient inflammatory process that occurs in response to the A β misfolding and aggregation, which is confirmed by the reduced A β 1-42 levels in CSF [127]. The same authors also showed that plasma YKL-40 levels increase with age, like CSF YKL-40. Elevated plasma YKL-40 concentrations seem to be correlated with male sex, older age, APOE ϵ 4 status, and cerebral accumulation of A β measured with PET [128, 129].

Another inflammatory protein considered as a plasma candidate biomarker for AD is GFAP. Elevated levels of GFAP in the plasma of preclinical AD patients suggest that astrocytic damage or activation may initiate during the preclinical phase of AD [122]. Prins et al., using the SIMOA technique, found significantly increased plasma GFAP concentrations in preclinical AD patients compared to healthy elderly individuals [127]. Additionally, plasma GFAP levels were notably higher in the A β + group compared to the A β - group. Verberk et al. suggested that plasma GFAP may serve as a valuable prognostic biomarker for predicting incident dementia [130]. Moreover, plasma GFAP demonstrated the highest impact and AUC in distinguishing between A β -positive cases and A β -negative cases CU, which was higher than AUC of plasma tTau. These findings suggest that integrating plasma GFAP into current theoretical models of AD pathogenesis could serve as a non-invasive and readily accessible method for detecting early astrocytosis secondary to A β pathology [127, 130, 131]. Additionally, Pereira et al. [132] concluded that plasma GFAP serves as an early indicator associated with brain A β pathology, though not tau aggregation, even in cognitively normal individuals with normal A β levels. The authors found a significant correlation between plasma GFAP concentrations and higher A β -PET signals in all A β -positive patients, as well as in cognitively normal subjects with normal A β values, even after adjusting for tau PET signals. They proposed that plasma GFAP should be incorporated into existing theoretical models of AD pathogenesis and used as a non-invasive and easily accessible means to detect early astrocytosis secondary to amyloid- β pathology [132].

According to recent research, chemokines such as MCP-1 or eotaxin-1 might serve as neuromodulators and are associated with increased memory impairment in MCI and AD [133]. Morgan et al. investigated plasma concentrations of various inflammatory markers, including complement components (C3, C4, C5), complement regulators (FH, FI), a soluble form of complement receptor (sCR1), a classical inflammation marker (CRP), and three chemokines (eotaxin-1, MCP-1, and MIP-1b) in AD, MCI, and elderly controls using the ELISA and MSD methods [134]. Using logistic regression, it was found that the most effective model for distinguishing between AD and elderly controls included sCR1, FB, FH, eotaxin-1, and MCP-1, alongside covariates such as age and APOE status with AUC = 0.79, while sCR1, MCP-1, eotaxin-1 optimally differentiated between AD and MCI, with AUC = 0.74. These models were confirmed in an independent cohort with AUC = 0.81 and 0.67, respectively [134].

These findings suggest that GFAP and YKL-40 serve as more sensitive indicators of inflammation in response to A β misfolding and aggregation, as evidenced by decreased A β 1-42 levels in the CSF. Using neuroinflammatory biomarkers to characterise individuals with preclinical AD is crucial for selecting subjects for new disease-modifying clinical trials. The authors suggest that evaluating these neuroimmune

response-related biomarkers during preclinical AD stages could aid in predicting which cognitively healthy elderly individuals are at higher risk of developing AD. Additionally, assessing plasma levels of GFAP and YKL-40 in individuals with preclinical AD may enhance differentiation between patients with reduced CSF A β 42 and otherwise healthy elderly individuals, thereby refining the definition of preclinical AD status. However, further research is necessary to determine whether these inflammatory plasma candidate biomarkers are specific to (preclinical) AD.

Neurofilament light (NfL) is a component of the axonal cytoskeleton and a marker of large calibre axonal degeneration that reflects pathological alteration in neurodegenerative diseases such as AD [135–137]. Some clinical investigations have demonstrated that NfL concentrations, and its rate of change in plasma, were higher in sporadic and familial AD and correlated with clinical symptoms and progression of the disease [138]. A growing body of evidence suggests that NfL could be a novel biomarker for early neurodegeneration in AD [135, 139–144]. Researchers have reported that in A β -positive cognitively normal and MCI participants, baseline NfL shows a significant predictive value in assessing tau burden in the left medial orbitofrontal cortex and para-hippocampus. They demonstrated an association between plasma NfL and multi-modal neuroimaging features in AD-vulnerable regions and its predictive value for future tau deposition [135]. Chatterjee et al. investigated plasma NfL in preclinical AD patients using SIMOA assays in CU, with absence (A β -) or presence (A β +) of brain amyloidosis. Increased plasma NfL was indicated in MCI A β + and AD A β +, compared to CU A β - and MCI A β - [123, 145]. Moreover, plasma NfL levels were found to be elevated in AD compared to CU individuals. Reduced plasma A β 1–42/A β 1–40 ratio and higher plasma NfL levels were correlated with a prospective cognitive decline. Increasing evidence shows that, in contrast to tau protein, the correlation between plasma and CSF levels of NfL protein is strong [146]. Mattsson et al. reported a significant increase in plasma NfL levels in AD cases, with an AUC = 0.87, which is comparable to the core AD CSF biomarkers [139]. In addition, plasma NfL levels were elevated in MCI cases with positive amyloid PET scans, and predicted faster cognitive deterioration, and higher rates of future hypometabolism and brain atrophy [139]. Moreover, blood NfL concentrations were higher in symptomatic FAD (familial AD) patients, but also in presymptomatic mutation carriers, with levels correlating with expected year of symptom onset as well as both cognitive and MRI measures of disease stage [147].

These findings suggest that plasma NfL reflects neurodegeneration also in the preclinical stage of AD. They suggest a future application for plasma NfL as a screening test at the first clinical evaluation of patients with cognitive disturbances. Thus, plasma NfL might serve as a simple, non-invasive and cheap screening tool, especially to exclude neurodegeneration [16].

Limitations

Although blood is more accessible than CSF, using a less invasive, low cost procedure, many investigators underline that measurement of blood biomarkers of AD patients has proven difficult [16]. CSF is connected with the brain extracellular fluid, via exchange of molecules from the brain to the CSF. However, only a fraction of brain proteins enters the bloodstream [16]. Furthermore, it has been proved that blood is a more challenging matrix than CSF for brain biomarkers, for several reasons. The small levels of brain proteins in the bloodstream have to be analysed in a matrix containing very high levels of plasma proteins, including albumin and immunoglobulins, entailing a high risk of interference in the analytical methods [148]. In addition, brain proteins that are released into the blood could be degraded by different proteases, and then metabolised in the liver or cleared by the kidneys. These processes are unrelated to brain changes, and are very difficult to regulate [16]. Moreover, in CSF, levels of A β 42 and A β 42/40 appear to decrease before amyloid PET reaches significant thresholds [22, 149, 150]. However, it is a great challenge to determine thresholds and identify cutoff points for plasma A β 42/40, which produces a much smaller effect than CSF A β 42/40 [101]. Additionally, this procedure could be more difficult among older patients, due to intermediate but sub-threshold amyloid pathology [98]. However, plasma candidate biomarkers useful in predicting future AD dementia are sorely needed.

Conclusions

A growing body of evidence suggests that the core AD CSF biomarkers such as tTau, pTau, A β 42 and the A β 42/40 ratio confirm a high diagnostic accuracy not only for AD dementia, but also for prodromal AD. Despite large strides in our understanding of AD pathogenesis, validated blood biomarkers for early detection and accurate diagnosis of AD patients are much needed. Many investigators believe that blood biomarkers may be implemented as screening tools in the initial clinical evaluation of this group of patients. Plasma A β 42/40 ratio seems to be the best candidate biomarker for discriminating A β -positive from A β -negative individuals. In addition, pTau217 differentiates clinically diagnosed AD dementia from non-AD neurodegenerative disorders with an accuracy comparable to that of CSF pTau and tau PET, while pTau231 distinguishes AD patients from patients with non-AD neurodegenerative disorders, as well as from A β negative MCI patients.

Among other non-specific-for-AD biomarkers, plasma GFAP levels correlate with an increased risk of progression to dementia and steeper cognitive decline, suggesting its role as a valuable prognostic biomarker predicting incident dementia. Moreover, researchers have found a significant increase of plasma NfL levels in AD cases, comparable to the core AD

CSF biomarkers, indicating the significance of plasma NfL as a screening test at the first clinical evaluation of patients with cognitive disturbance. Some researchers suggest that plasma AD biomarkers are more suitable than CSF or PET for implementation in primary care settings worldwide, and could lessen the cost of clinical trials by improving selection and stratification of participants and monitoring of treatment response.

However, a clinical implementation process, and validation of the assays' accuracy and robustness using predefined cut-offs across more diverse patient populations is crucial to further establish the utility of plasma candidate biomarkers in routine practice.

Article information

Acknowledgements: *PL has received consultation and/or lecture honoraria from IBL International, Fujirebio Europe, AJ Roboscreen, and Roche, and acts as a consultant to the European Commission; BM has received consultation and/or lecture honoraria from Abbott, Wiener, Roche, Cormay, Biameditek and TK Biotech; MLZ has received lecture honoraria from Roche.*

Authors' contributions: *All authors made significant contributions to conception, study design, acquisition of data, and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to submit to the current journal; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.*

Conflicts of interest: *The authors report no conflict of interest in this work.*

Funding: *None declared.*

References

- World Health Organization, Dementia. <https://www.who.int/news-room/fact-sheets/detail/dementia> (18 Aug 2021).
- Feigin VL, Vos T, Nichols E, et al. The global burden of neurological disorders: translating evidence into policy. *Lancet Neurol.* 2020; 19(3): 255–265, doi: [10.1016/S1474-4422\(19\)30411-9](https://doi.org/10.1016/S1474-4422(19)30411-9), indexed in Pubmed: [31813850](https://pubmed.ncbi.nlm.nih.gov/31813850/).
- Rocca WA, Petersen RC, Knopman DS, et al. Trends in the incidence and prevalence of Alzheimer's disease, dementia, and cognitive impairment in the United States. *Alzheimers Dement.* 2011; 7(1): 80–93, doi: [10.1016/j.jalz.2010.11.002](https://doi.org/10.1016/j.jalz.2010.11.002), indexed in Pubmed: [21255746](https://pubmed.ncbi.nlm.nih.gov/21255746/).
- Reitz C, Brayne C, Mayeux R. Epidemiology of Alzheimer disease. *Nat Rev Neurol.* 2011; 7(3): 137–152, doi: [10.1038/nrneurol.2011.2](https://doi.org/10.1038/nrneurol.2011.2), indexed in Pubmed: [21304480](https://pubmed.ncbi.nlm.nih.gov/21304480/).
- Dubois B, Feldman HH, Jacova C, et al. Research criteria for the diagnosis of Alzheimer's disease: revising the NINCDS-ADRDA criteria. *Lancet Neurol.* 2007; 6(8): 734–746, doi: [10.1016/S1474-4422\(07\)70178-3](https://doi.org/10.1016/S1474-4422(07)70178-3), indexed in Pubmed: [17616482](https://pubmed.ncbi.nlm.nih.gov/17616482/).
- Petersen RC, Aisen PS, Beckett LA, et al. Alzheimer's Disease Neuroimaging Initiative (ADNI): clinical characterization. *Neurology.* 2010; 74(3): 201–209, doi: [10.1212/WNL.0b013e3181cb3e25](https://doi.org/10.1212/WNL.0b013e3181cb3e25), indexed in Pubmed: [20042704](https://pubmed.ncbi.nlm.nih.gov/20042704/).
- Hansson O, Lehmann S, Otto M, et al. Advantages and disadvantages of the use of the CSF Amyloid β (A β) 42/40 ratio in the diagnosis of Alzheimer's Disease. *Alzheimers Res Ther.* 2019; 11(1): 34, doi: [10.1186/s13195-019-0485-0](https://doi.org/10.1186/s13195-019-0485-0), indexed in Pubmed: [31010420](https://pubmed.ncbi.nlm.nih.gov/31010420/).
- Lewczuk P, Riederer P, O'Bryant SE, et al. Members of the WFSBP Task Force Working on this Topic: Peter Riederer, Carla Gallo, Dimitrios Kapogiannis, Andrea Lopez Mato, Florence Thibaut. Cerebrospinal fluid and blood biomarkers for neurodegenerative dementias: An update of the Consensus of the Task Force on Biological Markers in Psychiatry of the World Federation of Societies of Biological Psychiatry. *World J Biol Psychiatry.* 2018; 19(4): 244–328, doi: [10.1080/15622975.2017.1375556](https://doi.org/10.1080/15622975.2017.1375556), indexed in Pubmed: [29076399](https://pubmed.ncbi.nlm.nih.gov/29076399/).
- Carroll CM, Li YM. Physiological and pathological roles of the γ -secretase complex. *Brain Res Bull.* 2016; 126(Pt 2): 199–206, doi: [10.1016/j.brainresbull.2016.04.019](https://doi.org/10.1016/j.brainresbull.2016.04.019), indexed in Pubmed: [27133790](https://pubmed.ncbi.nlm.nih.gov/27133790/).
- Portelius E, Price E, Brinkmalm G, et al. A novel pathway for amyloid precursor protein processing. *Neurobiol Aging.* 2011; 32(6): 1090–1098, doi: [10.1016/j.neurobiolaging.2009.06.002](https://doi.org/10.1016/j.neurobiolaging.2009.06.002), indexed in Pubmed: [19604603](https://pubmed.ncbi.nlm.nih.gov/19604603/).
- Lewczuk P, Matzen A, Blennow K, et al. Cerebrospinal Fluid A β 42/40 Corresponds Better than A β 42 to Amyloid PET in Alzheimer's Disease. *J Alzheimers Dis.* 2017; 55(2): 813–822, doi: [10.3233/JAD-160722](https://doi.org/10.3233/JAD-160722), indexed in Pubmed: [27792012](https://pubmed.ncbi.nlm.nih.gov/27792012/).
- Olsson B, Lautner R, Andreasson U, et al. CSF and blood biomarkers for the diagnosis of Alzheimer's disease: a systematic review and meta-analysis. *Lancet Neurol.* 2016; 15(7): 673–684, doi: [10.1016/S1474-4422\(16\)00070-3](https://doi.org/10.1016/S1474-4422(16)00070-3), indexed in Pubmed: [27068280](https://pubmed.ncbi.nlm.nih.gov/27068280/).
- Hulstaert F, Blennow K, Ivanoiu A, et al. Improved discrimination of AD patients using β -amyloid₍₁₋₄₂₎ and tau levels in CSF. *Neurology.* 1999; 52(8): 1555–1555, doi: [10.1212/wnl.52.8.1555](https://doi.org/10.1212/wnl.52.8.1555).
- Galasko D, Chang L, Motter R, et al. High cerebrospinal fluid tau and low amyloid beta42 levels in the clinical diagnosis of Alzheimer disease and relation to apolipoprotein E genotype. *Arch Neurol.* 1998; 55(7): 937–945, doi: [10.1001/archneur.55.7.937](https://doi.org/10.1001/archneur.55.7.937), indexed in Pubmed: [9678311](https://pubmed.ncbi.nlm.nih.gov/9678311/).
- Lewczuk P, Esselmann H, Bibl M, et al. Tau protein phosphorylated at threonine 181 in CSF as a neurochemical biomarker in Alzheimer's disease: original data and review of the literature. *J Mol Neurosci.* 2004; 23(1-2): 115–122, doi: [10.1385/JMN:23:1-2:115](https://doi.org/10.1385/JMN:23:1-2:115), indexed in Pubmed: [15126697](https://pubmed.ncbi.nlm.nih.gov/15126697/).
- Blennow K, Zetterberg H. Biomarkers for Alzheimer's disease: current status and prospects for the future. *J Intern Med.* 2018; 284(6): 643–663, doi: [10.1111/joim.12816](https://doi.org/10.1111/joim.12816), indexed in Pubmed: [30051512](https://pubmed.ncbi.nlm.nih.gov/30051512/).
- Lewczuk P, Esselmann H, Otto M, et al. Neurochemical diagnosis of Alzheimer's dementia by CSF A β 42, A β 42/A β 40 ratio and total tau. *Neurobiol Aging.* 2004; 25(3): 273–281, doi: [10.1016/S0197-4580\(03\)00086-1](https://doi.org/10.1016/S0197-4580(03)00086-1), indexed in Pubmed: [15123331](https://pubmed.ncbi.nlm.nih.gov/15123331/).
- Wiltfang J, Esselmann H, Bibl M, et al. Amyloid β peptide ratio 42/40 but not A β 42 correlates with phospho-Tau in patients with low- and high-CSF A β 40 load. *Journal of Neurochemistry.* 2006; 101(4): 1053–1059, doi: [10.1111/j.1471-4159.2006.04404.x](https://doi.org/10.1111/j.1471-4159.2006.04404.x).
- Lewczuk P, Wiltfang J, Kornhuber J, et al. Distributions of A β 42 and A β 42/40 in the Cerebrospinal Fluid in View of the Probability Theory. *Diagnostics (Basel).* 2021; 11(12), doi: [10.3390/diagnostics11122372](https://doi.org/10.3390/diagnostics11122372), indexed in Pubmed: [34943609](https://pubmed.ncbi.nlm.nih.gov/34943609/).

20. Palmqvist S, Zetterberg H, Blennow K, et al. Accuracy of brain amyloid detection in clinical practice using cerebrospinal fluid β -amyloid 42: a cross-validation study against amyloid positron emission tomography. *JAMA Neurol.* 2014; 71(10): 1282–1289, doi: [10.1001/jamaneurol.2014.1358](https://doi.org/10.1001/jamaneurol.2014.1358), indexed in Pubmed: [25155658](https://pubmed.ncbi.nlm.nih.gov/25155658/).
21. Palmqvist S, Zetterberg H, Mattsson N, et al. Alzheimer's Disease Neuroimaging Initiative, Swedish BioFINDER Study Group. Detailed comparison of amyloid PET and CSF biomarkers for identifying early Alzheimer disease. *Neurology.* 2015; 85(14): 1240–1249, doi: [10.1212/WNL.0000000000001991](https://doi.org/10.1212/WNL.0000000000001991), indexed in Pubmed: [26354982](https://pubmed.ncbi.nlm.nih.gov/26354982/).
22. Palmqvist S, Mattsson N, Hansson O, et al. Alzheimer's Disease Neuroimaging Initiative. Cerebrospinal fluid analysis detects cerebral amyloid- β accumulation earlier than positron emission tomography. *Brain.* 2016; 139(Pt 4): 1226–1236, doi: [10.1093/brain/aww015](https://doi.org/10.1093/brain/aww015), indexed in Pubmed: [26936941](https://pubmed.ncbi.nlm.nih.gov/26936941/).
23. Lewczuk P, Kornhuber J. Do we still need positron emission tomography for early Alzheimer's disease diagnosis? *Brain.* 2016; 139(11): e60, doi: [10.1093/brain/aww168](https://doi.org/10.1093/brain/aww168), indexed in Pubmed: [27383528](https://pubmed.ncbi.nlm.nih.gov/27383528/).
24. Henjum K, Almdahl IS, Årskog V, et al. Cerebrospinal fluid soluble TREM2 in aging and Alzheimer's disease. *Alzheimers Res Ther.* 2016; 8(1): 17, doi: [10.1186/s13195-016-0182-1](https://doi.org/10.1186/s13195-016-0182-1), indexed in Pubmed: [27121148](https://pubmed.ncbi.nlm.nih.gov/27121148/).
25. Mattsson N, Insel PS, Palmqvist S, et al. Increased amyloidogenic APP processing in APOE ϵ 4-negative individuals with cerebral β -amyloidosis. *Nat Commun.* 2016; 7: 10918, doi: [10.1038/ncomms10918](https://doi.org/10.1038/ncomms10918), indexed in Pubmed: [26948379](https://pubmed.ncbi.nlm.nih.gov/26948379/).
26. Janelidze S, Zetterberg H, Mattsson N, et al. Swedish BioFINDER study group. CSF A β 42/A β 40 and A β 42/A β 38 ratios: better diagnostic markers of Alzheimer disease. *Ann Clin Transl Neurol.* 2016; 3(3): 154–165, doi: [10.1002/acn3.274](https://doi.org/10.1002/acn3.274), indexed in Pubmed: [27042676](https://pubmed.ncbi.nlm.nih.gov/27042676/).
27. Mulugeta E, Londos E, Ballard C, et al. CSF amyloid β 38 as a novel diagnostic marker for dementia with Lewy bodies. *J Neurol Neurosurg Psychiatry.* 2011; 82(2): 160–164, doi: [10.1136/jnnp.2009.199398](https://doi.org/10.1136/jnnp.2009.199398), indexed in Pubmed: [21047883](https://pubmed.ncbi.nlm.nih.gov/21047883/).
28. Lista S, Toschi N, Baldacci F, et al. Alzheimer Precision Medicine Initiative (APMI). Cerebrospinal Fluid Neurogranin as a Biomarker of Neurodegenerative Diseases: A Cross-Sectional Study. *J Alzheimers Dis.* 2017; 59(4): 1327–1334, doi: [10.3233/JAD-170368](https://doi.org/10.3233/JAD-170368), indexed in Pubmed: [28731449](https://pubmed.ncbi.nlm.nih.gov/28731449/).
29. Herskovits AZ, Locascio JJ, Peskind ER, et al. A Luminex assay detects amyloid β oligomers in Alzheimer's disease cerebrospinal fluid. *PLoS One.* 2013; 8(7): e67898, doi: [10.1371/journal.pone.0067898](https://doi.org/10.1371/journal.pone.0067898), indexed in Pubmed: [23844122](https://pubmed.ncbi.nlm.nih.gov/23844122/).
30. Fukumoto H, Tokuda T, Kasai T, et al. High-molecular-weight beta-amyloid oligomers are elevated in cerebrospinal fluid of Alzheimer patients. *FASEB J.* 2010; 24(8): 2716–2726, doi: [10.1096/fj.09-150359](https://doi.org/10.1096/fj.09-150359), indexed in Pubmed: [20339023](https://pubmed.ncbi.nlm.nih.gov/20339023/).
31. Gao CM, Yam AY, Wang X, et al. A β 40 oligomers identified as a potential biomarker for the diagnosis of Alzheimer's disease. *PLoS One.* 2010; 5(12): e15725, doi: [10.1371/journal.pone.0015725](https://doi.org/10.1371/journal.pone.0015725), indexed in Pubmed: [21209907](https://pubmed.ncbi.nlm.nih.gov/21209907/).
32. Mandelkow E, von Bergen M, Biernat J, et al. Structural principles of tau and the paired helical filaments of Alzheimer's disease. *Brain Pathol.* 2007; 17(1): 83–90, doi: [10.1111/j.1750-3639.2007.00053.x](https://doi.org/10.1111/j.1750-3639.2007.00053.x), indexed in Pubmed: [17493042](https://pubmed.ncbi.nlm.nih.gov/17493042/).
33. Sunderland T, Linker G, Mirza N, et al. Decreased beta-amyloid1-42 and increased tau levels in cerebrospinal fluid of patients with Alzheimer disease. *JAMA.* 2003; 289(16): 2094–2103, doi: [10.1001/jama.289.16.2094](https://doi.org/10.1001/jama.289.16.2094), indexed in Pubmed: [12709467](https://pubmed.ncbi.nlm.nih.gov/12709467/).
34. Kohnken R, Buerger K, Zinkowski R, et al. Detection of tau phosphorylated at threonine 231 in cerebrospinal fluid of Alzheimer's disease patients. *Neurosci Lett.* 2000; 287(3): 187–190, doi: [10.1016/S0304-3940\(00\)01178-2](https://doi.org/10.1016/S0304-3940(00)01178-2), indexed in Pubmed: [10863026](https://pubmed.ncbi.nlm.nih.gov/10863026/).
35. Hampel H, Buerger K, Zinkowski R, et al. Measurement of phosphorylated tau epitopes in the differential diagnosis of Alzheimer disease: a comparative cerebrospinal fluid study. *Arch Gen Psychiatry.* 2004; 61(1): 95–102, doi: [10.1001/archpsyc.61.1.95](https://doi.org/10.1001/archpsyc.61.1.95), indexed in Pubmed: [14706948](https://pubmed.ncbi.nlm.nih.gov/14706948/).
36. Parnetti L, Lanari A, Amici S, et al. Phospho-Tau International Study Group. CSF phosphorylated tau is a possible marker for discriminating Alzheimer's disease from dementia with Lewy bodies. *Phospho-Tau International Study Group. Neurol Sci.* 2001; 22(1): 77–78, doi: [10.1007/s100720170055](https://doi.org/10.1007/s100720170055), indexed in Pubmed: [11487210](https://pubmed.ncbi.nlm.nih.gov/11487210/).
37. Arai H, Ishiguro K, Ohno H, et al. CSF phosphorylated tau protein and mild cognitive impairment: a prospective study. *Exp Neurol.* 2000; 166(1): 201–203, doi: [10.1006/exnr.2000.7501](https://doi.org/10.1006/exnr.2000.7501), indexed in Pubmed: [11031097](https://pubmed.ncbi.nlm.nih.gov/11031097/).
38. Suárez-Calvet M, Karikari TK, Ashton NJ, et al. ALFA Study. Novel tau biomarkers phosphorylated at T181, T217 or T231 rise in the initial stages of the preclinical Alzheimer's continuum when only subtle changes in A β pathology are detected. *EMBO Mol Med.* 2020; 12(12): e12921, doi: [10.15252/emmm.202012921](https://doi.org/10.15252/emmm.202012921), indexed in Pubmed: [33169916](https://pubmed.ncbi.nlm.nih.gov/33169916/).
39. Leuzy A, Janelidze S, Mattsson-Carligen N, et al. Comparing the Clinical Utility and Diagnostic Performance of CSF P-Tau181, P-Tau217, and P-Tau231 Assays. *Neurology.* 2021; 97(17): e1681–e1694, doi: [10.1212/WNL.00000000000012727](https://doi.org/10.1212/WNL.00000000000012727), indexed in Pubmed: [34493616](https://pubmed.ncbi.nlm.nih.gov/34493616/).
40. Buchhave P, Minthon L, Zetterberg H, et al. Cerebrospinal fluid levels of β -amyloid 1-42, but not of tau, are fully changed already 5 to 10 years before the onset of Alzheimer dementia. *Arch Gen Psychiatry.* 2012; 69(1): 98–106, doi: [10.1001/archgenpsychiatry.2011.155](https://doi.org/10.1001/archgenpsychiatry.2011.155), indexed in Pubmed: [22213792](https://pubmed.ncbi.nlm.nih.gov/22213792/).
41. de Leon MJ, Pirraglia E, Osorio RS, et al. Alzheimer's Disease Neuroimaging Initiative, National Alzheimer's Coordinating Center. The nonlinear relationship between cerebrospinal fluid A β 42 and tau in preclinical Alzheimer's disease. *PLoS One.* 2018; 13(2): e0191240, doi: [10.1371/journal.pone.0191240](https://doi.org/10.1371/journal.pone.0191240), indexed in Pubmed: [29415068](https://pubmed.ncbi.nlm.nih.gov/29415068/).
42. Molinuevo JL, Ayton S, Batrla R, et al. Current state of Alzheimer's fluid biomarkers. *Acta Neuropathol.* 2018; 136(6): 821–853, doi: [10.1007/s00401-018-1932-x](https://doi.org/10.1007/s00401-018-1932-x), indexed in Pubmed: [30488277](https://pubmed.ncbi.nlm.nih.gov/30488277/).
43. Shen Y, Wang H, Sun Q, et al. Increased Plasma Beta-Secretase 1 May Predict Conversion to Alzheimer's Disease Dementia in Individuals With Mild Cognitive Impairment. *Biol Psychiatry.* 2018; 83(5): 447–455, doi: [10.1016/j.biopsych.2017.02.007](https://doi.org/10.1016/j.biopsych.2017.02.007), indexed in Pubmed: [28359566](https://pubmed.ncbi.nlm.nih.gov/28359566/).
44. Leung YY, Toledo JB, Nefedov A, et al. Identifying amyloid pathology-related cerebrospinal fluid biomarkers for Alzheimer's disease in a multicohort study. *Alzheimers Dement (Amst).* 2015; 1(3): 339–348, doi: [10.1016/j.dadm.2015.06.008](https://doi.org/10.1016/j.dadm.2015.06.008), indexed in Pubmed: [26693175](https://pubmed.ncbi.nlm.nih.gov/26693175/).

45. Guo LH, Alexopoulos P, Perneczky R. Heart-type fatty acid binding protein and vascular endothelial growth factor: cerebrospinal fluid biomarker candidates for Alzheimer's disease. *Eur Arch Psychiatry Clin Neurosci*. 2013; 263(7): 553–560, doi: [10.1007/s00406-013-0405-4](https://doi.org/10.1007/s00406-013-0405-4), indexed in Pubmed: [23591828](https://pubmed.ncbi.nlm.nih.gov/23591828/).
46. Heslegrave A, Heywood W, Paterson R, et al. Increased cerebrospinal fluid soluble TREM2 concentration in Alzheimer's disease. *Mol Neurodegener*. 2016; 11: 3, doi: [10.1186/s13024-016-0071-x](https://doi.org/10.1186/s13024-016-0071-x), indexed in Pubmed: [26754172](https://pubmed.ncbi.nlm.nih.gov/26754172/).
47. Baldacci F, Lista S, Cavado E, et al. Diagnostic function of the neuroinflammatory biomarker YKL-40 in Alzheimer's disease and other neurodegenerative diseases. *Expert Rev Proteomics*. 2017; 14(4): 285–299, doi: [10.1080/14789450.2017.1304217](https://doi.org/10.1080/14789450.2017.1304217), indexed in Pubmed: [28281838](https://pubmed.ncbi.nlm.nih.gov/28281838/).
48. Muszyński P, Kulczyńska-Przybyk A, Borawska R, et al. The Relationship between Markers of Inflammation and Degeneration in the Central Nervous System and the Blood-Brain Barrier Impairment in Alzheimer's Disease. *J Alzheimers Dis*. 2017; 59(3): 903–912, doi: [10.3233/JAD-170220](https://doi.org/10.3233/JAD-170220), indexed in Pubmed: [28697565](https://pubmed.ncbi.nlm.nih.gov/28697565/).
49. Iturria-Medina Y, Sotero RC, Toussaint PJ, et al. Alzheimer's Disease Neuroimaging Initiative. Early role of vascular dysregulation on late-onset Alzheimer's disease based on multifactorial data-driven analysis. *Nat Commun*. 2016; 7: 11934.
50. Pereira JB, Westman E, Hansson O. Alzheimer's Disease Neuroimaging Initiative. Association between cerebrospinal fluid and plasma neurodegeneration biomarkers with brain atrophy in Alzheimer's disease. *Neurobiol Aging*. 2017; 58: 14–29.
51. Öhrfelt A, Brinkmalm A, Dumurgier J, et al. The pre-synaptic vesicle protein synaptotagmin is a novel biomarker for Alzheimer's disease. *Alzheimers Res Ther*. 2016; 8(1): 41, doi: [10.1186/s13195-016-0208-8](https://doi.org/10.1186/s13195-016-0208-8), indexed in Pubmed: [27716408](https://pubmed.ncbi.nlm.nih.gov/27716408/).
52. Chiasserini D, Biscetti L, Eusebi P, et al. Differential role of CSF fatty acid binding protein 3, α -synuclein, and Alzheimer's disease core biomarkers in Lewy body disorders and Alzheimer's dementia. *Alzheimers Res Ther*. 2017; 9(1): 52, doi: [10.1186/s13195-017-0276-4](https://doi.org/10.1186/s13195-017-0276-4), indexed in Pubmed: [28750675](https://pubmed.ncbi.nlm.nih.gov/28750675/).
53. Feneberg E, Steinacker P, Lehnert S, et al. Limited role of free TDP-43 as a diagnostic tool in neurodegenerative diseases. *Amyotroph Lateral Scler Frontotemporal Degener*. 2014; 15(5-6): 351–356, doi: [10.3109/21678421.2014.905606](https://doi.org/10.3109/21678421.2014.905606), indexed in Pubmed: [24834468](https://pubmed.ncbi.nlm.nih.gov/24834468/).
54. Mroczko B, Groblewska M, Zboch M, et al. Evaluation of visinin-like protein 1 concentrations in the cerebrospinal fluid of patients with mild cognitive impairment as a dynamic biomarker of Alzheimer's disease. *J Alzheimers Dis*. 2015; 43(3): 1031–1037, doi: [10.3233/JAD-141050](https://doi.org/10.3233/JAD-141050), indexed in Pubmed: [25159667](https://pubmed.ncbi.nlm.nih.gov/25159667/).
55. Zetterberg H, Skillbäck T, Mattsson N, et al. Alzheimer's Disease Neuroimaging Initiative. Association of Cerebrospinal Fluid Neurofilament Light Concentration With Alzheimer Disease Progression. *JAMA Neurol*. 2016; 73(1): 60–67, doi: [10.1001/jamaneurol.2015.3037](https://doi.org/10.1001/jamaneurol.2015.3037), indexed in Pubmed: [26524180](https://pubmed.ncbi.nlm.nih.gov/26524180/).
56. Mroczko B, Groblewska M, Zboch M, et al. Concentrations of matrix metalloproteinases and their tissue inhibitors in the cerebrospinal fluid of patients with Alzheimer's disease. *J Alzheimers Dis*. 2014; 40(2): 351–357, doi: [10.3233/JAD-131634](https://doi.org/10.3233/JAD-131634), indexed in Pubmed: [24448781](https://pubmed.ncbi.nlm.nih.gov/24448781/).
57. Rohrer JD, Woollacott IOC, Dick KM, et al. Serum neurofilament light chain protein is a measure of disease intensity in frontotemporal dementia. *Neurology*. 2016; 87(13): 1329–1336, doi: [10.1212/WNL.0000000000003154](https://doi.org/10.1212/WNL.0000000000003154), indexed in Pubmed: [27581216](https://pubmed.ncbi.nlm.nih.gov/27581216/).
58. Barage SH, Sonawane KD. Amyloid cascade hypothesis: Pathogenesis and therapeutic strategies in Alzheimer's disease. *Neuropeptides*. 2015; 52: 1–18, doi: [10.1016/j.npep.2015.06.008](https://doi.org/10.1016/j.npep.2015.06.008), indexed in Pubmed: [26149638](https://pubmed.ncbi.nlm.nih.gov/26149638/).
59. Morris GP, Clark IA, Vissel B. Inconsistencies and controversies surrounding the amyloid hypothesis of Alzheimer's disease. *Acta Neuropathol Commun*. 2014; 2: 135, doi: [10.1186/s40478-014-0135-5](https://doi.org/10.1186/s40478-014-0135-5), indexed in Pubmed: [25231068](https://pubmed.ncbi.nlm.nih.gov/25231068/).
60. Pedersen JT, Sigurdsson EM. Tau immunotherapy for Alzheimer's disease. *Trends Mol Med*. 2015; 21(6): 394–402, doi: [10.1016/j.molmed.2015.03.003](https://doi.org/10.1016/j.molmed.2015.03.003), indexed in Pubmed: [25846560](https://pubmed.ncbi.nlm.nih.gov/25846560/).
61. Lewczuk P, Leleental N, Lachmann I, et al. Non-Phosphorylated Tau as a Potential Biomarker of Alzheimer's Disease: Analytical and Diagnostic Characterization. *J Alzheimers Dis*. 2017; 55(1): 159–170, doi: [10.3233/JAD-160448](https://doi.org/10.3233/JAD-160448), indexed in Pubmed: [27662295](https://pubmed.ncbi.nlm.nih.gov/27662295/).
62. Ermann N, Lewczuk P, Schmitz M, et al. CSF nonphosphorylated Tau as a biomarker for the discrimination of AD from CJD. *Ann Clin Transl Neurol*. 2018; 5(7): 883–887, doi: [10.1002/acn3.584](https://doi.org/10.1002/acn3.584), indexed in Pubmed: [30009207](https://pubmed.ncbi.nlm.nih.gov/30009207/).
63. Foiani MS, Cicognola C, Ermann N, et al. Searching for novel cerebrospinal fluid biomarkers of tau pathology in frontotemporal dementia: an elusive quest. *J Neurol Neurosurg Psychiatry*. 2019; 90(7): 740–746, doi: [10.1136/jnnp-2018-319266](https://doi.org/10.1136/jnnp-2018-319266), indexed in Pubmed: [30981993](https://pubmed.ncbi.nlm.nih.gov/30981993/).
64. Parnetti L, Cicognola C, Eusebi P, et al. Value of cerebrospinal fluid α -synuclein species as biomarker in Parkinson's diagnosis and prognosis. *Biomark Med*. 2016; 10(1): 35–49, doi: [10.2217/bmm.15.107](https://doi.org/10.2217/bmm.15.107), indexed in Pubmed: [26643452](https://pubmed.ncbi.nlm.nih.gov/26643452/).
65. Eusebi P, Giannandrea D, Biscetti L, et al. Diagnostic utility of CSF α -synuclein species in Parkinson's disease: protocol for a systematic review and meta-analysis. *BMJ Open*. 2016; 6(6): e011113, doi: [10.1136/bmjopen-2016-011113](https://doi.org/10.1136/bmjopen-2016-011113), indexed in Pubmed: [27297011](https://pubmed.ncbi.nlm.nih.gov/27297011/).
66. Twohig D, Nielsen HM. α -synuclein in the pathophysiology of Alzheimer's disease. *Mol Neurodegener*. 2019; 14(1): 23, doi: [10.1186/s13024-019-0320-x](https://doi.org/10.1186/s13024-019-0320-x), indexed in Pubmed: [31186026](https://pubmed.ncbi.nlm.nih.gov/31186026/).
67. Slaets S, Vanmechelen E, Le Bastard N, et al. Increased CSF α -synuclein levels in Alzheimer's disease: correlation with tau levels. *Alzheimers Dement*. 2014; 10(5 Suppl): S290–S298, doi: [10.1016/j.jalz.2013.10.004](https://doi.org/10.1016/j.jalz.2013.10.004), indexed in Pubmed: [24439167](https://pubmed.ncbi.nlm.nih.gov/24439167/).
68. Engelborghs S, Niemantsverdriet E, Struyfs H, et al. Consensus guidelines for lumbar puncture in patients with neurological diseases. *Alzheimers Dement (Amst)*. 2017; 8: 111–126, doi: [10.1016/j.dadm.2017.04.007](https://doi.org/10.1016/j.dadm.2017.04.007), indexed in Pubmed: [28603768](https://pubmed.ncbi.nlm.nih.gov/28603768/).
69. Hansson O, Mikulskis A, Fagan AM, et al. The impact of preanalytical variables on measuring cerebrospinal fluid biomarkers for Alzheimer's disease diagnosis: A review. *Alzheimers Dement*. 2018; 14(10): 1313–1333, doi: [10.1016/j.jalz.2018.05.008](https://doi.org/10.1016/j.jalz.2018.05.008), indexed in Pubmed: [29940161](https://pubmed.ncbi.nlm.nih.gov/29940161/).
70. Bjerke M, Andreasson U, Kuhlmann J, et al. Assessing the commutability of reference material formats for the harmonization of amyloid- β measurements. *Clin Chem Lab Med*. 2016; 54(7): 1177–1191, doi: [10.1515/cclm-2015-0733](https://doi.org/10.1515/cclm-2015-0733), indexed in Pubmed: [26495928](https://pubmed.ncbi.nlm.nih.gov/26495928/).
71. Lewczuk P, Zimmermann R, Wiltfang J, et al. Neurochemical dementia diagnostics: a simple algorithm for interpretation of the CSF biomarkers. *J Neural Transm (Vienna)*. 2009; 116(9): 1163–1167, doi: [10.1007/s00702-009-0277-y](https://doi.org/10.1007/s00702-009-0277-y), indexed in Pubmed: [19653063](https://pubmed.ncbi.nlm.nih.gov/19653063/).
72. Lehmann S, Delaby C, Boursier G, et al. Relevance of A β 42/40 Ratio for Detection of Alzheimer Disease Pathology in Clinical Routine:

- The PLM Scale. *Front Aging Neurosci.* 2018; 10: 138, doi: [10.3389/fnagi.2018.00138](https://doi.org/10.3389/fnagi.2018.00138), indexed in Pubmed: [29892221](https://pubmed.ncbi.nlm.nih.gov/29892221/).
73. Jack CR, Bennett DA, Blennow K, et al. A/T/N: An unbiased descriptive classification scheme for Alzheimer disease biomarkers. *Neurology.* 2016; 87(5): 539–547, doi: [10.1212/WNL.0000000000002923](https://doi.org/10.1212/WNL.0000000000002923), indexed in Pubmed: [27371494](https://pubmed.ncbi.nlm.nih.gov/27371494/).
 74. Lewczuk P, Kornhuber J, Toledo JB, et al. German Dementia Competence Network, US-ADNI. Validation of the Erlangen Score Algorithm for the Prediction of the Development of Dementia due to Alzheimer's Disease in Pre-Dementia Subjects. *J Alzheimers Dis.* 2015; 48(2): 433–441, doi: [10.3233/JAD-150342](https://doi.org/10.3233/JAD-150342), indexed in Pubmed: [26402007](https://pubmed.ncbi.nlm.nih.gov/26402007/).
 75. Baldeiras I, Santana I, Leitão MJ, et al. Erlangen Score as a tool to predict progression from mild cognitive impairment to dementia in Alzheimer's disease. *Alzheimers Res Ther.* 2019; 11(1): 2, doi: [10.1186/s13195-018-0456-x](https://doi.org/10.1186/s13195-018-0456-x), indexed in Pubmed: [30611311](https://pubmed.ncbi.nlm.nih.gov/30611311/).
 76. Skillbäck T, Kornhuber J, Blennow K, et al. Alzheimer's Disease Neuroimaging Initiative. Erlangen Score Predicts Cognitive and Neuroimaging Progression in Mild Cognitive Impairment Stage of Alzheimer's Disease. *J Alzheimers Dis.* 2019; 69(2): 551–559, doi: [10.3233/JAD-190067](https://doi.org/10.3233/JAD-190067), indexed in Pubmed: [31104027](https://pubmed.ncbi.nlm.nih.gov/31104027/).
 77. Somers C, Lewczuk P, Sieben A, et al. Validation of the Erlangen Score Algorithm for Differential Dementia Diagnosis in Autopsy-Confirmed Subjects. *J Alzheimers Dis.* 2019; 68(3): 1151–1159, doi: [10.3233/JAD-180563](https://doi.org/10.3233/JAD-180563), indexed in Pubmed: [30883344](https://pubmed.ncbi.nlm.nih.gov/30883344/).
 78. Janelidze S, Berron D, Smith R, et al. Associations of Plasma Phospho-Tau217 Levels With Tau Positron Emission Tomography in Early Alzheimer Disease. *JAMA Neurol.* 2021; 78(2): 149–156, doi: [10.1001/jamaneurol.2020.4201](https://doi.org/10.1001/jamaneurol.2020.4201), indexed in Pubmed: [33165506](https://pubmed.ncbi.nlm.nih.gov/33165506/).
 79. Beach TG, Monsell SE, Phillips LE, et al. Accuracy of the clinical diagnosis of Alzheimer disease at National Institute on Aging Alzheimer Disease Centers, 2005-2010. *J Neuropathol Exp Neurol.* 2012; 71(4): 266–273, doi: [10.1097/NEN.0b013e31824b211b](https://doi.org/10.1097/NEN.0b013e31824b211b), indexed in Pubmed: [22437338](https://pubmed.ncbi.nlm.nih.gov/22437338/).
 80. Jack CR, Bennett DA, Blennow K, et al. Contributors. NIA-AA Research Framework: Toward a biological definition of Alzheimer's disease. *Alzheimers Dement.* 2018; 14(4): 535–562, doi: [10.1016/j.jalz.2018.02.018](https://doi.org/10.1016/j.jalz.2018.02.018), indexed in Pubmed: [29653606](https://pubmed.ncbi.nlm.nih.gov/29653606/).
 81. Palmqvist S, Stomrud E, Cullen N, et al. An accurate fully automated panel of plasma biomarkers for Alzheimer's disease. *Alzheimers Dement.* 2023; 19(4): 1204–1215, doi: [10.1002/alz.12751](https://doi.org/10.1002/alz.12751), indexed in Pubmed: [35950735](https://pubmed.ncbi.nlm.nih.gov/35950735/).
 82. Dubois B, Villain N, Frisoni GB, et al. Clinical diagnosis of Alzheimer's disease: recommendations of the International Working Group. *Lancet Neurol.* 2021; 20(6): 484–496, doi: [10.1016/S1474-4422\(21\)00066-1](https://doi.org/10.1016/S1474-4422(21)00066-1), indexed in Pubmed: [33933186](https://pubmed.ncbi.nlm.nih.gov/33933186/).
 83. Rabinovici GD, Carrillo MC, Apgar C, et al. Association of Amyloid Positron Emission Tomography With Subsequent Change in Clinical Management Among Medicare Beneficiaries With Mild Cognitive Impairment or Dementia. *JAMA.* 2019; 321(13): 1286–1294, doi: [10.1001/jama.2019.2000](https://doi.org/10.1001/jama.2019.2000), indexed in Pubmed: [30938796](https://pubmed.ncbi.nlm.nih.gov/30938796/).
 84. Brand AL, Lawler PE, Bollinger JG, et al. The performance of plasma amyloid beta measurements in identifying amyloid plaques in Alzheimer's disease: a literature review. *Alzheimers Res Ther.* 2022; 14(1): 195, doi: [10.1186/s13195-022-01117-1](https://doi.org/10.1186/s13195-022-01117-1), indexed in Pubmed: [36575454](https://pubmed.ncbi.nlm.nih.gov/36575454/).
 85. Bateman RJ, Munsell LY, Morris JC, et al. Human amyloid-beta synthesis and clearance rates as measured in cerebrospinal fluid in vivo. *Nat Med.* 2006; 12(7): 856–861, doi: [10.1038/nm1438](https://doi.org/10.1038/nm1438), indexed in Pubmed: [16799555](https://pubmed.ncbi.nlm.nih.gov/16799555/).
 86. Hansson O. Biomarkers for neurodegenerative diseases. *Nat Med.* 2021; 27(6): 954–963, doi: [10.1038/s41591-021-01382-x](https://doi.org/10.1038/s41591-021-01382-x), indexed in Pubmed: [34083813](https://pubmed.ncbi.nlm.nih.gov/34083813/).
 87. Ovod V, Ramsey KN, Mawuenyega KG, et al. Amyloid β concentrations and stable isotope labeling kinetics of human plasma specific to central nervous system amyloidosis. *Alzheimers Dement.* 2017; 13(8): 841–849, doi: [10.1016/j.jalz.2017.06.2266](https://doi.org/10.1016/j.jalz.2017.06.2266), indexed in Pubmed: [28734653](https://pubmed.ncbi.nlm.nih.gov/28734653/).
 88. Teunissen CE, Verberk IMW, Thijssen EH, et al. Blood-based biomarkers for Alzheimer's disease: towards clinical implementation. *Lancet Neurol.* 2022; 21(1): 66–77, doi: [10.1016/S1474-4422\(21\)00361-6](https://doi.org/10.1016/S1474-4422(21)00361-6), indexed in Pubmed: [34838239](https://pubmed.ncbi.nlm.nih.gov/34838239/).
 89. O'Bryant SE, Mielke MM, Rissman RA, et al. Biofluid Based Biomarker Professional Interest Area. Blood-based biomarkers in Alzheimer disease: Current state of the science and a novel collaborative paradigm for advancing from discovery to clinic. *Alzheimers Dement.* 2017; 13(1): 45–58, doi: [10.1016/j.jalz.2016.09.014](https://doi.org/10.1016/j.jalz.2016.09.014), indexed in Pubmed: [27870940](https://pubmed.ncbi.nlm.nih.gov/27870940/).
 90. Pannee J, Törnqvist U, Westerlund A, et al. The amyloid- β degradation pattern in plasma—a possible tool for clinical trials in Alzheimer's disease. *Neurosci Lett.* 2014; 573: 7–12, doi: [10.1016/j.neulet.2014.04.041](https://doi.org/10.1016/j.neulet.2014.04.041), indexed in Pubmed: [24796810](https://pubmed.ncbi.nlm.nih.gov/24796810/).
 91. Schindler SE, Bollinger JG, Ovod V, et al. High-precision plasma β -amyloid 42/40 predicts current and future brain amyloidosis. *Neurology.* 2019; 93(17): e1647–e1659, doi: [10.1212/WNL.0000000000008081](https://doi.org/10.1212/WNL.0000000000008081), indexed in Pubmed: [31371569](https://pubmed.ncbi.nlm.nih.gov/31371569/).
 92. Pérez-Grijalva V, Arbizu J, Romero J, et al. AB255 Study Group. Plasma A β 42/40 ratio alone or combined with FDG-PET can accurately predict amyloid-PET positivity: a cross-sectional analysis from the AB255 Study. *Alzheimers Res Ther.* 2019; 11(1): 96, doi: [10.1186/s13195-019-0549-1](https://doi.org/10.1186/s13195-019-0549-1), indexed in Pubmed: [31787105](https://pubmed.ncbi.nlm.nih.gov/31787105/).
 93. Tanaka T, Ruifen JC, Nai YH, et al. Head-to-head comparison of amplified plasmonic exosome A β 42 platform and single-molecule array immunoassay in a memory clinic cohort. *Eur J Neurol.* 2021; 28(5): 1479–1489, doi: [10.1111/ene.14704](https://doi.org/10.1111/ene.14704), indexed in Pubmed: [33370497](https://pubmed.ncbi.nlm.nih.gov/33370497/).
 94. Smirnov DS, Ashton NJ, Blennow K, et al. Plasma biomarkers for Alzheimer's Disease in relation to neuropathology and cognitive change. *Acta Neuropathol.* 2022; 143(4): 487–503, doi: [10.1007/s00401-022-02408-5](https://doi.org/10.1007/s00401-022-02408-5), indexed in Pubmed: [35195758](https://pubmed.ncbi.nlm.nih.gov/35195758/).
 95. Keshavan A, Pannee J, Karikari T, et al. Population-based blood screening for pre-clinical Alzheimer's disease: a British birth cohort at age 70. *Brain.* 2021; 144(2): 434–449, doi: [0.1093/brain/awaa403](https://doi.org/10.1093/brain/awaa403), indexed in Pubmed: [33479777](https://pubmed.ncbi.nlm.nih.gov/33479777/).
 96. Nakamura A, Kaneko N, Villemagne VL, et al. High performance plasma amyloid- β biomarkers for Alzheimer's disease. *Nature.* 2018; 554(7691): 249–254, doi: [10.1038/nature25456](https://doi.org/10.1038/nature25456), indexed in Pubmed: [29420472](https://pubmed.ncbi.nlm.nih.gov/29420472/).
 97. West T, Kirmess KM, Meyer MR, et al. A blood-based diagnostic test incorporating plasma A β 42/40 ratio, ApoE proteotype, and age accurately identifies brain amyloid status: findings from a multi cohort validity analysis. *Mol Neurodegener.* 2021; 16(1): 30, doi: [10.1186/s13024-021-00451-6](https://doi.org/10.1186/s13024-021-00451-6), indexed in Pubmed: [33933117](https://pubmed.ncbi.nlm.nih.gov/33933117/).
 98. Janelidze S, Stomrud E, Palmqvist S, et al. Plasma β -amyloid in Alzheimer's disease and vascular disease. *Sci Rep.* 2016; 6: 26801, doi: [10.1038/srep26801](https://doi.org/10.1038/srep26801), indexed in Pubmed: [27241045](https://pubmed.ncbi.nlm.nih.gov/27241045/).
 99. Hansson O, Zetterberg H, Vanmechelen E, et al. Evaluation of plasma A β (40) and A β (42) as predictors of conversion to Alzheimer's

- disease in patients with mild cognitive impairment. *Neurobiol Aging*. 2010; 31: 357–67.
100. Kuo YM, Emmerling MR, Lampert HC, et al. High levels of circulating Abeta42 are sequestered by plasma proteins in Alzheimer's disease. *Biochem Biophys Res Commun*. 1999; 257(3): 787–791, doi: [10.1006/bbrc.1999.0552](https://doi.org/10.1006/bbrc.1999.0552), indexed in Pubmed: [10208861](https://pubmed.ncbi.nlm.nih.gov/10208861/).
101. Mattsson N, Schöll M, Strandberg O, et al. ¹⁸F-AV-1451 and CSF T-tau and P-tau as biomarkers in Alzheimer's disease. *EMBO Molecular Medicine*. 2017; 9(9): 1212–1223, doi: [10.15252/emmm.201707809](https://doi.org/10.15252/emmm.201707809).
102. Mattsson-Carlgrén N, Andersson E, Janelidze S, et al. Aβ deposition is associated with increases in soluble and phosphorylated tau that precede a positive Tau PET in Alzheimer's disease. *Sci Adv*. 2020; 6(16): eaaz2387, doi: [10.1126/sciadv.aaz2387](https://doi.org/10.1126/sciadv.aaz2387), indexed in Pubmed: [32426454](https://pubmed.ncbi.nlm.nih.gov/32426454/).
103. Meyer PF, Pichet Binette A, Gonneaud J, et al. Characterization of Alzheimer Disease Biomarker Discrepancies Using Cerebrospinal Fluid Phosphorylated Tau and AV1451 Positron Emission Tomography. *JAMA Neurol*. 2020; 77(4): 508–516, doi: [10.1001/jamaneurol.2019.4749](https://doi.org/10.1001/jamaneurol.2019.4749), indexed in Pubmed: [31961372](https://pubmed.ncbi.nlm.nih.gov/31961372/).
104. Bejanin A, Schonhaut DR, La Joie R, et al. Tau pathology and neurodegeneration contribute to cognitive impairment in Alzheimer's disease. *Brain*. 2017; 140(12): 3286–3300, doi: [10.1093/brain/awx243](https://doi.org/10.1093/brain/awx243), indexed in Pubmed: [29053874](https://pubmed.ncbi.nlm.nih.gov/29053874/).
105. Ossenkoppele R, Smith R, Ohlsson T, et al. Associations between tau, Aβ, and cortical thickness with cognition in Alzheimer disease. *Neurology*. 2019; 92(6), doi: [10.1212/wnl.0000000000006875](https://doi.org/10.1212/wnl.0000000000006875).
106. Mattsson N, Zetterberg H, Janelidze S, et al. ADNI Investigators. Plasma tau in Alzheimer disease. *Neurology*. 2016; 87(17): 1827–1835, doi: [10.1212/WNL.0000000000003246](https://doi.org/10.1212/WNL.0000000000003246), indexed in Pubmed: [27694257](https://pubmed.ncbi.nlm.nih.gov/27694257/).
107. Mielke MM, Hagen CE, Xu J, et al. Plasma phospho-tau181 increases with Alzheimer's disease clinical severity and is associated with tau- and amyloid-positron emission tomography. *Alzheimers Dement*. 2018; 14(8): 989–997, doi: [10.1016/j.jalz.2018.02.013](https://doi.org/10.1016/j.jalz.2018.02.013), indexed in Pubmed: [29626426](https://pubmed.ncbi.nlm.nih.gov/29626426/).
108. Tatebe H, Kasai T, Ohmichi T, et al. Quantification of plasma phosphorylated tau to use as a biomarker for brain Alzheimer pathology: pilot case-control studies including patients with Alzheimer's disease and down syndrome. *Mol Neurodegener*. 2017; 12(1): 63, doi: [10.1186/s13024-017-0206-8](https://doi.org/10.1186/s13024-017-0206-8), indexed in Pubmed: [28866979](https://pubmed.ncbi.nlm.nih.gov/28866979/).
109. Yang CC, Chiu MJ, Chen TF, et al. Assay of Plasma Phosphorylated Tau Protein (Threonine 181) and Total Tau Protein in Early-Stage Alzheimer's Disease. *J Alzheimers Dis*. 2018; 61(4): 1323–1332, doi: [10.3233/JAD-170810](https://doi.org/10.3233/JAD-170810), indexed in Pubmed: [29376870](https://pubmed.ncbi.nlm.nih.gov/29376870/).
110. Coomans EM, Verberk IMW, Ossenkoppele R, et al. A Head-to-Head Comparison Between Plasma pTau181 and Tau PET Along the Alzheimer's Disease Continuum. *J Nucl Med*. 2023; 64(3): 437–443, doi: [10.2967/jnumed.122.264279](https://doi.org/10.2967/jnumed.122.264279), indexed in Pubmed: [36229187](https://pubmed.ncbi.nlm.nih.gov/36229187/).
111. Janelidze S, Mattsson N, Palmqvist S, et al. Plasma P-tau181 in Alzheimer's disease: relationship to other biomarkers, differential diagnosis, neuropathology and longitudinal progression to Alzheimer's dementia. *Nat Med*. 2020; 26(3): 379–386, doi: [10.1038/s41591-020-0755-1](https://doi.org/10.1038/s41591-020-0755-1), indexed in Pubmed: [32123385](https://pubmed.ncbi.nlm.nih.gov/32123385/).
112. Barthélemy NR, Bateman RJ, Hirtz C, et al. Cerebrospinal fluid phospho-tau T217 outperforms T181 as a biomarker for the differential diagnosis of Alzheimer's disease and PET amyloid-positive patient identification. *Alzheimers Res Ther*. 2020; 12(1): 26, doi: [10.1186/s13195-020-00596-4](https://doi.org/10.1186/s13195-020-00596-4), indexed in Pubmed: [32183883](https://pubmed.ncbi.nlm.nih.gov/32183883/).
113. Janelidze S, Stomrud E, Smith R, et al. Cerebrospinal fluid p-tau217 performs better than p-tau181 as a biomarker of Alzheimer's disease. *Nat Commun*. 2020; 11(1): 1683, doi: [10.1038/s41467-020-15436-0](https://doi.org/10.1038/s41467-020-15436-0), indexed in Pubmed: [32246036](https://pubmed.ncbi.nlm.nih.gov/32246036/).
114. Palmqvist S, Insel PS, Stomrud E, et al. Cerebrospinal fluid and plasma biomarker trajectories with increasing amyloid deposition in Alzheimer's disease. *EMBO Mol Med*. 2019; 11(12): e11170, doi: [10.15252/emmm.201911170](https://doi.org/10.15252/emmm.201911170), indexed in Pubmed: [31709776](https://pubmed.ncbi.nlm.nih.gov/31709776/).
115. Palmqvist S, Janelidze S, Quiroz YT, et al. Discriminative Accuracy of Plasma Phospho-tau217 for Alzheimer Disease vs Other Neurodegenerative Disorders. *JAMA*. 2020; 324(8): 772–781, doi: [10.1001/jama.2020.12134](https://doi.org/10.1001/jama.2020.12134), indexed in Pubmed: [32722745](https://pubmed.ncbi.nlm.nih.gov/32722745/).
116. Janelidze S, Bali D, Ashton NJ, et al. Head-to-head comparison of 10 plasma phospho-tau assays in prodromal Alzheimer's disease. *Brain*. 2023; 146(4): 1592–1601, doi: [10.1093/brain/awac333](https://doi.org/10.1093/brain/awac333), indexed in Pubmed: [36087307](https://pubmed.ncbi.nlm.nih.gov/36087307/).
117. Palmqvist S, Tideman P, Cullen N, et al. Alzheimer's Disease Neuroimaging Initiative. Prediction of future Alzheimer's disease dementia using plasma phospho-tau combined with other accessible measures. *Nat Med*. 2021; 27(6): 1034–1042, doi: [10.1038/s41591-021-01348-z](https://doi.org/10.1038/s41591-021-01348-z), indexed in Pubmed: [34031605](https://pubmed.ncbi.nlm.nih.gov/34031605/).
118. Ashton NJ, Pascoal TA, Karikari TK, et al. Plasma p-tau231: a new biomarker for incipient Alzheimer's disease pathology. *Acta Neuropathol*. 2021; 141(5): 709–724, doi: [10.1007/s00401-021-02275-6](https://doi.org/10.1007/s00401-021-02275-6), indexed in Pubmed: [33585983](https://pubmed.ncbi.nlm.nih.gov/33585983/).
119. Brickman AM, Manly JJ, Honig LS, et al. Plasma p-tau181, p-tau217, and other blood-based Alzheimer's disease biomarkers in a multi-ethnic, community study. *Alzheimers Dement*. 2021; 17(8): 1353–1364, doi: [10.1002/alz.12301](https://doi.org/10.1002/alz.12301), indexed in Pubmed: [33580742](https://pubmed.ncbi.nlm.nih.gov/33580742/).
120. Karikari TK, Emeršič A, Vrillon A, et al. Head-to-head comparison of clinical performance of CSF phospho-tau T181 and T217 biomarkers for Alzheimer's disease diagnosis. *Alzheimers Dement*. 2021; 17(5): 755–767, doi: [10.1002/alz.12236](https://doi.org/10.1002/alz.12236), indexed in Pubmed: [33252199](https://pubmed.ncbi.nlm.nih.gov/33252199/).
121. Thijssen EH, La Joie R, Strom A, et al. Advancing Research and Treatment for Frontotemporal Lobar Degeneration investigators. Plasma phosphorylated tau 217 and phosphorylated tau 181 as biomarkers in Alzheimer's disease and frontotemporal lobar degeneration: a retrospective diagnostic performance study. *Lancet Neurol*. 2021; 20(9): 739–752, doi: [10.1016/S1474-4422\(21\)00214-3](https://doi.org/10.1016/S1474-4422(21)00214-3), indexed in Pubmed: [34418401](https://pubmed.ncbi.nlm.nih.gov/34418401/).
122. Chatterjee P, Pedrini S, Stoops E, et al. Plasma glial fibrillary acidic protein is elevated in cognitively normal older adults at risk of Alzheimer's disease. *Translational Psychiatry*. 2021; 11(1), doi: [10.1038/s41398-020-01137-1](https://doi.org/10.1038/s41398-020-01137-1).
123. Chatterjee P, Pedrini S, Doecke JD, et al. AIBL Research Group. Plasma Aβ42/40 ratio, p-tau181, GFAP, and NFL across the Alzheimer's disease continuum: A cross-sectional and longitudinal study in the AIBL cohort. *Alzheimers Dement*. 2023; 19(4): 1117–1134, doi: [10.1002/alz.12724](https://doi.org/10.1002/alz.12724), indexed in Pubmed: [36574591](https://pubmed.ncbi.nlm.nih.gov/36574591/).
124. Brum WS, Cullen NC, Therriault J, et al. A blood-based biomarker workflow for optimal tau-PET referral in memory clinic settings. *Nat Commun*. 2024; 15(1): 2311, doi: [10.1038/s41467-024-46603-2](https://doi.org/10.1038/s41467-024-46603-2), indexed in Pubmed: [38486040](https://pubmed.ncbi.nlm.nih.gov/38486040/).
125. Saitgareeva A, Bulygin K, Gareev I, et al. The role of microglia in the development of neurodegeneration. *Neurological Sciences*. 2020; 41(12): 3609–3615, doi: [10.1007/s10072-020-04468-5](https://doi.org/10.1007/s10072-020-04468-5).

126. Heneka M, Carson M, Khoury J, et al. Neuroinflammation in Alzheimer's disease. *Lancet Neurol.* 2015; 14(4): 388–405, doi: [10.1016/S1474-4422\(15\)70016-5](https://doi.org/10.1016/S1474-4422(15)70016-5).
127. Prins S, de Kam ML, Teunissen CE, et al. Inflammatory plasma biomarkers in subjects with preclinical Alzheimer's disease. *Alzheimers Res Ther.* 2022; 14(1): 106, doi: [10.1186/s13195-022-01051-2](https://doi.org/10.1186/s13195-022-01051-2), indexed in Pubmed: [35922871](https://pubmed.ncbi.nlm.nih.gov/35922871/).
128. Vergallo A, Lista S, Lemercier P, et al. INSIGHT-preAD study group and the Alzheimer Precision Medicine Initiative (APMI), INSIGHT-preAD study group, Alzheimer Precision Medicine Initiative (APMI). Association of plasma YKL-40 with brain amyloid- β levels, memory performance, and sex in subjective memory complainers. *Neurobiol Aging.* 2020; 96: 22–32, doi: [10.1016/j.neurobiolaging.2020.07.009](https://doi.org/10.1016/j.neurobiolaging.2020.07.009), indexed in Pubmed: [32920471](https://pubmed.ncbi.nlm.nih.gov/32920471/).
129. Kronborg G, Ostergaard C, Weis N, et al. Serum level of YKL-40 is elevated in patients with *Streptococcus pneumoniae* bacteremia and is associated with the outcome of the disease. *Scand J Infect Dis.* 2002; 34(5): 323–326, doi: [10.1080/00365540110080233](https://doi.org/10.1080/00365540110080233), indexed in Pubmed: [12069012](https://pubmed.ncbi.nlm.nih.gov/12069012/).
130. Verberk IMW, Laarhuis MB, van den Bosch KA, et al. Serum markers glial fibrillary acidic protein and neurofilament light for prognosis and monitoring in cognitively normal older people: a prospective memory clinic-based cohort study. *Lancet Healthy Longev.* 2021; 2(2): e87–e95, doi: [10.1016/S2666-7568\(20\)30061-1](https://doi.org/10.1016/S2666-7568(20)30061-1), indexed in Pubmed: [36098162](https://pubmed.ncbi.nlm.nih.gov/36098162/).
131. Prins S, Zhuparris A, Hart EP, et al. A cross-sectional study in healthy elderly subjects aimed at development of an algorithm to increase identification of Alzheimer pathology for the purpose of clinical trial participation. *Alzheimers Res Ther.* 2021; 13(1): 132, doi: [10.1186/s13195-021-00874-9](https://doi.org/10.1186/s13195-021-00874-9), indexed in Pubmed: [34274005](https://pubmed.ncbi.nlm.nih.gov/34274005/).
132. Pereira J, Janelidze S, Smith R, et al. Plasma GFAP is an early marker of amyloid- β but not tau pathology in Alzheimer's disease. *Brain.* 2021; 144(11): 3505–3516, doi: [10.1093/brain/awab223](https://doi.org/10.1093/brain/awab223).
133. Bettcher BM, Fitch R, Wynn MJ, et al. MCP-1 and eotaxin-1 selectively and negatively associate with memory in MCI and Alzheimer's disease dementia phenotypes. *Alzheimers Dement (Amst).* 2016; 3: 91–97, doi: [10.1016/j.dadm.2016.05.004](https://doi.org/10.1016/j.dadm.2016.05.004), indexed in Pubmed: [27453930](https://pubmed.ncbi.nlm.nih.gov/27453930/).
134. Morgan AR, Touchard S, Leckey C, et al. NIMA Consortium, Annex: NIMA-Wellcome Trust Consortium for Neuroimmunology of Mood Disorders and Alzheimer's Disease. Inflammatory biomarkers in Alzheimer's disease plasma. *Alzheimers Dement.* 2019; 15(6): 776–787, doi: [10.1016/j.jalz.2019.03.007](https://doi.org/10.1016/j.jalz.2019.03.007), indexed in Pubmed: [31047856](https://pubmed.ncbi.nlm.nih.gov/31047856/).
135. Hu Q, Shi M, Li Y, et al. Elevated plasma neurofilament light was associated with multi-modal neuroimaging features in Alzheimer's Disease signature regions and predicted future tau deposition. *Res Sq.* 2024, doi: [10.21203/rs.3.rs-3946421/v1](https://doi.org/10.21203/rs.3.rs-3946421/v1), indexed in Pubmed: [38464117](https://pubmed.ncbi.nlm.nih.gov/38464117/).
136. Yuan A, Rao M, Nixon R. Neurofilaments and Neurofilament Proteins in Health and Disease. *Cold Spring Harbor Perspectives in Biology.* 2017; 9(4): a018309, doi: [10.1101/cshperspect.a018309](https://doi.org/10.1101/cshperspect.a018309).
137. Norgren N, Rosengren L, Stigbrand T. Elevated neurofilament levels in neurological diseases. *Brain Research.* 2003; 987(1): 25–31, doi: [10.1016/S0006-8993\(03\)03219-0](https://doi.org/10.1016/S0006-8993(03)03219-0).
138. Preische O, Schultz SA, Apel A, et al. Dominantly Inherited Alzheimer Network. Serum neurofilament dynamics predicts neurodegeneration and clinical progression in presymptomatic Alzheimer's disease. *Nat Med.* 2019; 25(2): 277–283, doi: [10.1038/s41591-018-0304-3](https://doi.org/10.1038/s41591-018-0304-3), indexed in Pubmed: [30664784](https://pubmed.ncbi.nlm.nih.gov/30664784/).
139. Mattsson N, Andreasson U, Zetterberg H, et al. Alzheimer's Disease Neuroimaging Initiative. Association of Plasma Neurofilament Light With Neurodegeneration in Patients With Alzheimer Disease. *JAMA Neurol.* 2017; 74(5): 557–566, doi: [10.1001/jamaneuro.2016.6117](https://doi.org/10.1001/jamaneuro.2016.6117), indexed in Pubmed: [28346578](https://pubmed.ncbi.nlm.nih.gov/28346578/).
140. Mattsson N, Cullen NC, Andreasson U, et al. Association Between Longitudinal Plasma Neurofilament Light and Neurodegeneration in Patients With Alzheimer Disease. *JAMA Neurol.* 2019; 76(7): 791–799, doi: [10.1001/jamaneuro.2019.0765](https://doi.org/10.1001/jamaneuro.2019.0765), indexed in Pubmed: [31009028](https://pubmed.ncbi.nlm.nih.gov/31009028/).
141. Benedet A, Leuzy A, Pascoal T, et al. Stage-specific links between plasma neurofilament light and imaging biomarkers of Alzheimer's disease. *Brain.* 2020; 143(12): 3793–3804, doi: [10.1093/brain/awaa342](https://doi.org/10.1093/brain/awaa342).
142. Nabizadeh F, Balabandian M, Rostami MR, et al. Alzheimer's Disease Neuroimaging Initiative (ADNI). Plasma neurofilament light levels correlate with white matter damage prior to Alzheimer's disease: results from ADNI. *Aging Clin Exp Res.* 2022; 34(10): 2363–2372, doi: [10.1007/s40520-022-02095-y](https://doi.org/10.1007/s40520-022-02095-y), indexed in Pubmed: [35226303](https://pubmed.ncbi.nlm.nih.gov/35226303/).
143. Lewczuk P, Ermann N, Andreasson U, et al. Plasma neurofilament light as a potential biomarker of neurodegeneration in Alzheimer's disease. *Alzheimers Res Ther.* 2018; 10(1): 71, doi: [10.1186/s13195-018-0404-9](https://doi.org/10.1186/s13195-018-0404-9), indexed in Pubmed: [30055655](https://pubmed.ncbi.nlm.nih.gov/30055655/).
144. Clark C, Lewczuk P, Kornhuber J, et al. Plasma neurofilament light and phosphorylated tau 181 as biomarkers of Alzheimer's disease pathology and clinical disease progression. *Alzheimers Res Ther.* 2021; 13(1): 65, doi: [10.1186/s13195-021-00805-8](https://doi.org/10.1186/s13195-021-00805-8), indexed in Pubmed: [33766131](https://pubmed.ncbi.nlm.nih.gov/33766131/).
145. Chatterjee P, Pedrini S, Ashton NJ, et al. Diagnostic and prognostic plasma biomarkers for preclinical Alzheimer's disease. *Alzheimers Dement.* 2022; 18(6): 1141–1154, doi: [10.1002/alz.12447](https://doi.org/10.1002/alz.12447), indexed in Pubmed: [34494715](https://pubmed.ncbi.nlm.nih.gov/34494715/).
146. Gisslén M, Price RW, Andreasson U, et al. Plasma Concentration of the Neurofilament Light Protein (NFL) is a Biomarker of CNS Injury in HIV Infection: A Cross-Sectional Study. *EBioMedicine.* 2016; 3: 135–140, doi: [10.1016/j.ebiom.2015.11.036](https://doi.org/10.1016/j.ebiom.2015.11.036), indexed in Pubmed: [26870824](https://pubmed.ncbi.nlm.nih.gov/26870824/).
147. Weston PSJ, Poole T, Ryan NS, et al. Serum neurofilament light in familial Alzheimer disease: A marker of early neurodegeneration. *Neurology.* 2017; 89(21): 2167–2175, doi: [10.1212/WNL.0000000000004667](https://doi.org/10.1212/WNL.0000000000004667), indexed in Pubmed: [29070659](https://pubmed.ncbi.nlm.nih.gov/29070659/).
148. Blennow K, Zetterberg H. Understanding biomarkers of neurodegeneration: Ultrasensitive detection techniques pave the way for mechanistic understanding. *Nat Med.* 2015; 21(3): 217–219, doi: [10.1038/nm.3810](https://doi.org/10.1038/nm.3810), indexed in Pubmed: [25742456](https://pubmed.ncbi.nlm.nih.gov/25742456/).
149. Guo T, Shaw LM, Trojanowski JQ, et al. Alzheimer's Disease Neuroimaging Initiative. Association of CSF A β , amyloid PET, and cognition in cognitively unimpaired elderly adults. *Neurology.* 2020; 95(15): e2075–e2085, doi: [10.1212/WNL.0000000000010596](https://doi.org/10.1212/WNL.0000000000010596), indexed in Pubmed: [32759202](https://pubmed.ncbi.nlm.nih.gov/32759202/).
150. Luo J, Agboola F, Grant E, et al. Sequence of Alzheimer disease biomarker changes in cognitively normal adults: A cross-sectional study. *Neurology.* 2020; 95(23): e3104–e3116, doi: [10.1212/WNL.0000000000010747](https://doi.org/10.1212/WNL.0000000000010747), indexed in Pubmed: [32873693](https://pubmed.ncbi.nlm.nih.gov/32873693/).