

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

journal homepage: <http://www.elsevier.com/locate/pjnns>

## Letter to the Editor

### An unusual radiological presentation of posterior reversible encephalopathy syndrome



#### Keywords:

PRES

Hypertension

Brain MRI

Radiological improvement

Dear Editor-in-Chief,

The patient was a 17-year-old girl who developed blurred vision, first in the left eye followed by the right eye, since one month ago. She started to have weakness of the lower limbs two weeks and a headache one week before she was visited. The headache was pulsatile and in the occipital region with nausea and vomiting. The patient developed restlessness three days before visiting.

On systemic examination, her blood pressure was 260/200 millimeter of mercury (mm Hg) and her pulse rate was 110/min. The systemic examination was otherwise unremarkable.

On neurologic examination, the patient was completely oriented but restless. Ophthalmoscopy revealed bilateral papilledema.

Moreover, left central facial palsy was also noted. Upon examination, the proximal force was decreased to 4/5 in the lower limbs. Blood tests on patient arrival showed potassium of 4.3 milliequivalents per liter (mEq/L), sodium of 122 mEq/L, and creatinine of 1.9 milligrams per deciliter (mg/dL). Other biochemical tests, complete blood count (CBC), and vasculitis tests were normal. Urinalysis showed 4–6 white

blood cells (WBCs), 8–10 red blood cells (RBCs), and 3+ proteins.

On renal ultrasound, the echo of the renal parenchyma was slightly increased. Doppler ultrasound of the renal arteries was also normal.

Brain magnetic resonance imaging (MRI) showed hyper-signal lesions in the pons and bilateral basal ganglia which did not enhance after contrast administration (Fig. 1a and b). Diffusion weighted (DW) images did not reveal any restriction in favor of stroke.

The patient received intravenous labetalol which caused a decrease in blood pressure. The neurologic condition of the patient improved following the decrease in blood pressure and her restlessness, blurred vision, facial palsy, and lower limbs weakness resolved. The hypersignal lesions were mostly resolved on the second brain MRI (Fig. 1c and d).


With regard to the considerable clinical and radiological improvement of the patient following decreasing blood pressure, a diagnosis of posterior reversible encephalopathy syndrome (PRES) was made.

PRES, which can occur in different contexts such as hypertension, renal failure, and administration of immunosuppressant, is characterized by a condition that is both clinically and radiologically reversible [1].

Radiologically, due to the vasogenic edema, involvement of the white matter, especially in the posterior region, is observed [2].

Atypical radiologic manifestations include isolated pons, frontal, thalamus, or basal ganglia involvement [3,4].

According to review of the literature, this is the first report of the radiologic manifestations of PRES as severe involvement of the pons and bilateral basal ganglia.



**Fig. 1 – (a and b) In axial Flair's view, hypersignal lesions in the pons and bilateral basal ganglia were seen. (c and d) After one month the hypersignal lesions were mostly resolved.**

---

### Conflict of interest

None declared.

---

### Financial support

None declared.

---

### Ethics

The work described in this article has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans; EU Directive 2010/63/EU for animal experiments;

Uniform Requirements for manuscripts submitted to Biomedical journals.

---

### REFERENCES

- [1] Karakis I, MacDonald JA, Stefanidou M, Kase CS. Clinical and radiological features of brainstem variant of hypertensive encephalopathy. *J Vasc Interv Neurol* 2009;2(2):172-6.
- [2] Yerdelen D, Giray S, Tan M, Yildirim T. Hypertensive encephalopathy with atypical MRI leukoencephalopathy affecting brain stem and cerebellum. *Acta Neurol Belg* 2009;109(2):142-5.
- [3] Gamanagatti S, Subramanian S. Hypertensive encephalopathy: isolated pons involvement mimicking central pontine myelinolysis. *Korean J Radiol* 2006;7(3): 218-9.

- [4] McKinney AM, Short J, Truwit CL, McKinney ZJ, Kozak OS, SantaCruz KS, et al. Posterior reversible encephalopathy syndrome: incidence of atypical regions of involvement and imaging findings. *AJR Am J Roentgenol* 2007;189(4):904–12.

Abdorreza Naser Moghadasi<sup>a,b\*</sup>  
Khadijeh Bozorgvari<sup>b</sup>  
Mahsa Arzani<sup>b</sup>

<sup>a</sup>MS Research Center, Neuroscience Institute,  
Tehran University of Medical Science, Tehran, Iran

<sup>b</sup>Department of Neurology, Sina Hospital,  
Tehran University of Medical Science, Tehran, Iran

\*Corresponding author at: MS Research Center, Sina Hospital,  
Tehran University of Medical Science, Hasan Abad Sq.,

Tehran 11367-46911, Iran. Tel.: +0098 21 66348571;  
fax: +0098 21 66348570  
E-mail address: [abdorrezamoghadasi@gmail.com](mailto:abdorrezamoghadasi@gmail.com)  
(A. Naser Moghadasi)

Received 16 January 2015

Accepted 26 March 2015

Available online 9 April 2015

0028-3843/

© 2015 Polish Neurological Society. Published by Elsevier Sp. z  
o.o. All rights reserved.

<http://dx.doi.org/10.1016/j.pjnns.2015.03.006>