

Migration of atrial catheter of ventriculo-cysto-atrial shunt into heart and pulmonary artery — case report and literature review

Václav Vybíhal^{1,2} , Martin Plevko^{1,2} , Pavel Fadrus^{1,2} , Martin Smrčka^{1,2} , Marek Sova^{1,2} ,
Martin Poloczek^{3,4}, Miloš Keřkovský^{5,6}

¹Department of Neurosurgery, Faculty of Medicine, Masaryk University, Brno, Czech Republic

²Department of Neurosurgery, University Hospital Brno, Czech Republic

³Internal Cardiology Clinic, Faculty of Medicine, Masaryk University, Brno, Czech Republic

⁴Internal Cardiology Clinic, University Hospital Brno, Czech Republic

⁵Department of Radiology and Nuclear Medicine, Faculty of Medicine, Masaryk University, Brno, Czech Republic

⁶Department of Radiology and Nuclear Medicine, University Hospital Brno, Czech Republic

ABSTRACT

Ventriculoatrial shunts are the alternative treatments when it is impossible to use ventriculoperitoneal shunts. Limited indication for ventriculoatrial shunt is due to the possibility of very serious complications inherent with this procedure. We present a case report of a young patient who suffered from disconnection of an atrial catheter from the valve after an accidental blow to his neck. The atrial catheter was dislocated to the heart and pulmonary artery and it was extracted through the femoral vein in the groin area using an endovascular technique. The procedure went without complications. A new atrial catheter was introduced under ultrasonic guidance during surgical revision.

Keywords: ventriculoatrial shunt, rare complication, atrial catheter, hydrocephalus

(*Neurol Neurochir Pol* 2024; 58 (4): 459–464)

Introduction

Shunt implantation is a universal method of treating patients who suffer from hydrocephalus [1]. Ventriculoperitoneal shunts (VPSs) are the current treatment of choice for hydrocephalus. If this system cannot be introduced, implantation of a ventriculoatrial shunt (VAS) is the next option in most centres. Limited indication for VAS is due to the possibility of very serious complications of this procedure [2]. In this case report, we present a rare case of migration of an atrial catheter into the heart and pulmonary artery.

Case report

A male in his 20s was admitted to the neurological department at the regional hospital after an accidental blow to his neck when he was opening the door of a bookshelf at home. Because there was an implanted VAS, our clinic was consulted and we were asked to admit the patient into our care.

The patient was found to have an enlarged head circumference in his first months of life. A temporal arachnoid cyst on the right side was diagnosed on magnetic resonance imaging. The patient underwent the implantation of a cysto-peritoneal

Address for correspondence: Martin Plevko, Department of Neurosurgery, University Hospital Brno Bohunice, Faculty of Medicine, Masaryk University, Brno, Czech Republic; e-mail: plevko.martin@fnbrno.cz

Received: 01.02.2024 Accepted: 07.04.2024 Early publication date: 12.06.2024

This article is available in open access under Creative Commons Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.




Figure 1. CT scan of brain with implanted ventriculo-peritoneal shunt (on left) and a cysto-peritoneal shunt (on right) at age of 12. First catheter is introduced into arachnoid cyst of right temporal lobe and second one into frontal horn of left ventricle. There are asymptomatic slit ventricles

shunt. After a few years, a programmable VPS was introduced because of a ventricular enlargement. The patient was checked by a paediatric neurologist and at the outpatient department of neurosurgery at the Children’s Hospital (Fig. 1). In all, the patient underwent six surgical revisions during his childhood.

After reaching adulthood, the patient began to be followed up in an adult neurosurgery department. Due to a malfunction of the cysto-peritoneal (CP) shunt, a revision was performed when the patient was in his early 20s. The CP shunt was extracted and a new catheter was inserted into the arachnoid cyst and connected to the VP shunt via a T-connector (Fig. 2). In the last revision in his late 20s, the ventriculo-cysto-peritoneal shunt was converted to a ventriculo-cysto-atrial shunt. The reason was a pseudocyst filled with cerebrospinal fluid in the abdominal cavity and a high amount of derived cerebrospinal fluid (Fig. 3).

On the present admission, the patient complained of neck pain but did not report any other problems. The patient was conscious, fully oriented and cooperative, without focal neurological deficit. The scars of the post-shunt implantation areas were healed without signs of inflammation. The valve chamber was palpable and filled promptly.

Laboratory samples and an X-ray image of the course of the shunt were performed, which showed the disconnection of the atrial catheter and its dislocation to the heart and pulmonary artery (Fig. 4). An invasive cardiologist was contacted immediately. It was decided to try to extract the displaced atrial catheter. The cardiac surgeon was also informed and was ready to perform open surgery to remove the atrial catheter if the endovascular procedure should fail.

The patient was taken to the interventional angiography department at the Cardiology clinic. After local anaesthesia with 1% mesocaine, access to the right femoral vein was obtained by the Seldinger technique. A GooseNeck 10 F catheter with a 10 mm loop was inserted through an introducer. Using this, the end of the atrial catheter was secured and extracted slowly out through the femoral vein in the groin area (Fig. 5).




Figure 2. X-ray scan of skull of patient after revision. Catheter introduced into temporal arachnoid cyst (yellow arrows) on right side was connected through T-connector to ventricular catheter (red arrows)



Figure 3. CT exam of abdominal cavity. CT exam of abdominal cavity with pseudocyst filled with cerebrospinal fluid around distal end of ventriculoperitoneal shunt (yellow arrows). For this reason, peritoneal catheter was exchanged for atrial catheter

The procedure went without complications. The puncture canal was treated with a suture.

Subsequently, a surgical revision was performed under general anaesthesia, during which a new atrial catheter was inserted into the patient under ultrasonic guidance. The procedure went without complications. Even before the revision, the patient began to report headaches as a sign of shunt dependency. After the revision, the symptoms subsided.




Figure 4. X-ray images of disconnected atrial catheter. X-ray images of ventriculo-cysto-atrial shunt show disconnected atrial catheter from valve moving towards heart and pulmonary artery. Last two images display difference in position of atrial catheter from initial image and just before its endovascular removal when distal end of catheter was located in pulmonary artery

The patient was discharged home on the fifth postoperative day. Surgical wounds were healed by first intention.

Discussion

After the introduction of the Spitz-Holter valve in 1952, VASs became a treatment choice for hydrocephalus [3]. Because over subsequent years various complications of VASs were recognised, VPSs were introduced in the late 1960s and showed fewer long-term complications compared to VASs [4]. As VASs drain cerebrospinal fluid directly into the bloodstream and into the heart they have shown an increased risk of serious complications compared to other techniques, but they remain an important alternative to VPSs in selected cases [5, 6].

Obstructions, malpositions and shunt infections are frequent complications in VASs [7, 8]. The most serious complications of VASs are shunt infections, thromboemboli and misplacement or blockage of the atrial catheter. Migration of the distal segment of a broken atrial catheter is a less common but very serious complication [9].

Atrial catheter migration can be classified either as proximal (retrograde) i.e. retreating back to the superior vena cava or internal jugular vein, or distal (anterograde) i.e. projecting into the tricuspid valve, the right ventricle, or even the




Figure 5. Endovascular removal of atrial catheter. Proximal end of atrial catheter was captured using a teflon-coated metal loop (yellow arrows) and was gradually extruded out of body through groin via venous system

Table 1. Literature review of atrial catheter migration into heart and pulmonary artery

Study	Age (years) /sex	Symptoms	Catheter location	Complication	Treatment	Outcome
Mori et al., 1993 [9]	21/M	Stupor	PA	None	Endovascular	GR
James et al., 1997 [25]	20/M	Headache, nausea, vomiting	PA	None	Endovascular	GR
So and Shirani, 2009 [26]	53/M	Severe chest discomfort	PA	None	Endovascular	GR
Irie et al., 2009 [27]	50/M	Chest distress	PA	Cardiopulmonary arrest (ischaemic cardiac failure)	Medical	Died
Matsubara et al., 2012 [28]	67/M	Disorientation, memory disturbance and gait disturbance	H	None	Endovascular	GR
Aloddadi et al., 2018 [29]	9/M	Progressive neck swelling	PA	None	Endovascular	GR
Xu et al., 2021 [30]	12/F	Poor feeding, fatigue	H	None	Endovascular	GR
López Sánchez et al., 2022 [31]	36/F	General discomfort, headache, neck pain	PA	None	Endovascular	GR
Zervos et al., 2022 [32]	72/F	NPH symptoms	H	None	Endovascular	GR
	79/M	NPH symptoms	H	None	Endovascular	GR
	32/F	Progressive headaches	H	None	Endovascular	GR

The PubMed database was searched for original articles published between 1990 and 2022 with the search term "ventriculoatrial". Eligibility criteria were (i) availability of English abstract and (ii) complications clearly attributed to distal catheter migration into the heart or pulmonary artery. M — male, F — female, PA — pulmonary artery, H — hearth, GR — good recovery

pulmonary arteries [10] (Tab. 1). Retrograde migration has been reported in both paediatric and adult populations and can occur when the VA shunt was inserted in infancy, and as the patient grows, it leads to catheter migration by upward pulling of the tubing. It can cause hepatic vein thrombosis, cervical subcutaneous abscess, superior vena cava syndrome, and thrombosis [11–13], but it is sometimes without consequence.

Anterograde migration can lead to right atrial thrombus, pulmonary embolism, pulmonary hypertension, cardiac tamponade, endocarditis and arrhythmias due to direct irritation of the sinoatrial and/or atrioventricular nodes in cases when the catheter tip is located too distally [10].

A broken segment of atrial catheter can be retrieved by a venous catheterisation technique via the transjugular approach [14] or more commonly via the transfemoral approach [9] by the cardiologist, and eventually by the interventional radiologist. If a long time has elapsed since the insertion, potential adhesion of the shunt catheter to intracardiac structures can be problematic for percutaneous removal [15, 16]. If this mini-invasive technique is unsuccessful, it is then necessary to remove the catheter or part of it from the open thoracotomy [17]. In the case of adhesion of the atrial catheter to the right ventricle, the pulmonary valve, pulmonary artery, or superior vena cava, the open heart approach is the safer option [16].

Migration of a VPS into the heart or pulmonary artery has been reported [18, 19]. It is assumed that the subcutaneous catheter guide perforates the internal or external jugular vein during the shunt procedure, and negative pressure in the

vein draws the catheter into the heart, despite no evidence of subcutaneous haematoma or signs of vessel injury during the original shunt procedure [18].

Appropriate vein selection and precise atrial catheter placement are important to reduce the complications of VASs. Therefore, new placement strategies and monitoring methods have been introduced to improve success and reduce complications in atrial catheter placement. Chuang et al. [20] reported the use of percutaneous placement with real-time transoesophageal echocardiogram monitoring because it was quick, safe, more accurate, and less invasive. The endovascular technique was presented by Gonzales et al. [21]. This technique can identify venous system easily, can demonstrate jugular vein patency, and can determine quickly the true placement or malposition [21]. Percutaneous placement of an atrial catheter with ultrasonography guidance was reported by Metellus et al. [22].

Minimising the risk of distal VAS disconnection is particularly important due to the risk of catheter embolisation. This rare event occurs in c.1.56% of implanted VA shunts [23]. There are no clear treatment guidelines in the literature to help prevent and manage this complication [24].

Conclusions

VASs are the usual alternative when it is impossible to use VPSs. Their use is connected with rare but serious complications. Appropriate vein selection, precise atrial catheter placement, and minimising the risk of atrial catheter disconnection

are important requirements to reduce the VAS complication rate. Endovascular removal of a displaced atrial catheter is the method of choice and is minimally invasive. Quick resolution of this complication is essential.

Authors' contributions: *All authors agree with the content of the manuscript, and all gave explicit consent after careful revision. Material preparation (including review), data collection and analysis were performed by Václav Vybíhal M.D. The final draft of the manuscript was written by Václav Vybíhal and Martin Plevko and all authors commented on previous versions of the manuscript.*

Funding: *Supported by Ministry of Health, Czech Republic — conceptual development of research organisation (FNBr, 65269705).*

Statements and declarations: *All authors certify that they have no affiliations with nor involvement in any organisation with any financial or nonfinancial interest in the subject matter or materials presented in the manuscript. The funder also confirms that it had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.*

Ethics approval: *Ethical approval was not necessary for the preparation of this article.*

References

- Nowosławska E, Moszura T, Mikołajczyk-Wieczorek W, et al. A hybrid technique for ventriculoatrial shunt implantation—technical note. *Childs Nerv Syst.* 2014; 30(10): 1729–1732, doi: [10.1007/s00381-014-2520-6](https://doi.org/10.1007/s00381-014-2520-6), indexed in Pubmed: [25119152](https://pubmed.ncbi.nlm.nih.gov/25119152/).
- Yavuz C, Demirtas S, Caliskan A, et al. Reasons, procedures, and outcomes in ventriculoatrial shunts: A single-center experience. *Surg Neurol Int.* 2013; 4: 10, doi: [10.4103/2152-7806.106284](https://doi.org/10.4103/2152-7806.106284), indexed in Pubmed: [23493480](https://pubmed.ncbi.nlm.nih.gov/23493480/).
- Gopal VV, Peethambaran AK. Rare sequelae following ventriculoatrial shunt: Case report and review of literature. *Asian J Neurosurg.* 2016; 11(2): 173, doi: [10.4103/1793-5482.175635](https://doi.org/10.4103/1793-5482.175635), indexed in Pubmed: [27057231](https://pubmed.ncbi.nlm.nih.gov/27057231/).
- Vernet O, Rilliet B. Late complications of ventriculoatrial or ventriculoperitoneal shunts. *Lancet.* 2001; 358(9293): 1569–1570, doi: [10.1016/S0140-6736\(01\)06670-3](https://doi.org/10.1016/S0140-6736(01)06670-3), indexed in Pubmed: [11716880](https://pubmed.ncbi.nlm.nih.gov/11716880/).
- Elhammady MS, Benglis DM, Bhatia S, et al. Ventriculoatrial shunt catheter displacement in a child with partial anomalous pulmonary venous return: case report. *J Neurosurg Pediatr.* 2008; 2(1): 68–70, doi: [10.3171/PED/2008/2/7/068](https://doi.org/10.3171/PED/2008/2/7/068), indexed in Pubmed: [18590399](https://pubmed.ncbi.nlm.nih.gov/18590399/).
- Hanak BW, Bonow RH, Harris CA, et al. Cerebrospinal fluid shunting complications in children. *pediatr neurosurg.* 2017; 52(6): 381–400, doi: [10.1159/000452840](https://doi.org/10.1159/000452840), indexed in Pubmed: [28249297](https://pubmed.ncbi.nlm.nih.gov/28249297/).
- Nash D, Schmidt K. Complications and management of ventriculoatrial (VA) shunts, a case report. *Cerebrospinal Fluid Research.* 2010; 7(S1), doi: [10.1186/1743-8454-7-s1-s42](https://doi.org/10.1186/1743-8454-7-s1-s42).
- Natarajan A, Mazhar S. Right heart complications of ventriculoatrial shunt. *Eur Heart J.* 2011; 32(17): 2134, doi: [10.1093/eurheartj/ehr164](https://doi.org/10.1093/eurheartj/ehr164), indexed in Pubmed: [21609972](https://pubmed.ncbi.nlm.nih.gov/21609972/).
- Mori T, Arisawa M, Fukuoka M, et al. Management of a broken atrial catheter migrated into the heart: a rare complication of ventriculoatrial shunt—case report. *Neurol Med Chir (Tokyo).* 1993; 33(10): 713–715, doi: [10.2176/nmc.33.713](https://doi.org/10.2176/nmc.33.713), indexed in Pubmed: [7505902](https://pubmed.ncbi.nlm.nih.gov/7505902/).
- Pradini-Santos L, Craven CL, Watkins LD, et al. Ventriculoatrial shunt catheter tip migration causing tricuspid regurgitation: case report and review of the literature. *World Neurosurg.* 2020; 136: 83–89, doi: [10.1016/j.wneu.2020.01.016](https://doi.org/10.1016/j.wneu.2020.01.016), indexed in Pubmed: [31931241](https://pubmed.ncbi.nlm.nih.gov/31931241/).
- O'Shea PA. Inferior vena cava and hepatic vein thrombosis as a rare complication of ventriculoatrial shunt. Case report. *J Neurosurg.* 1978; 48(1): 143–145, doi: [10.3171/jns.1978.48.1.0143](https://doi.org/10.3171/jns.1978.48.1.0143), indexed in Pubmed: [619017](https://pubmed.ncbi.nlm.nih.gov/619017/).
- Hung AL, Vivas-Buitrago T, Adam A, et al. Ventriculoatrial versus ventriculoperitoneal shunt complications in idiopathic normal pressure hydrocephalus. *Clin Neurol Neurosurg.* 2017; 157: 1–6, doi: [10.1016/j.clineuro.2017.03.014](https://doi.org/10.1016/j.clineuro.2017.03.014), indexed in Pubmed: [28347957](https://pubmed.ncbi.nlm.nih.gov/28347957/).
- Yurtseven T, Erşahin Y, Kitiş O, et al. Thrombosis and thrombophlebitis of the internal jugular vein as a very rare complication of the ventriculoatrial shunt. *Clin Neurol Neurosurg.* 2005; 107(2): 144–146, doi: [10.1016/j.clineuro.2004.05.002](https://doi.org/10.1016/j.clineuro.2004.05.002), indexed in Pubmed: [15708232](https://pubmed.ncbi.nlm.nih.gov/15708232/).
- Tatsumi T, Howland WJ. Retrieval of a ventriculoatrial shunt catheter from the heart by a venous catheterization technique. Technical note. *J Neurosurg.* 1970; 32(5): 593–596, doi: [10.3171/jns.1970.32.5.0593](https://doi.org/10.3171/jns.1970.32.5.0593), indexed in Pubmed: [5438102](https://pubmed.ncbi.nlm.nih.gov/5438102/).
- Fewel ME, Garton HJL. Migration of distal ventriculoperitoneal shunt catheter into the heart. Case report and review of the literature. *J Neurosurg.* 2004; 100(2 Suppl Pediatrics): 206–211, doi: [10.3171/ped.2004.100.2.0206](https://doi.org/10.3171/ped.2004.100.2.0206), indexed in Pubmed: [14758952](https://pubmed.ncbi.nlm.nih.gov/14758952/).
- Morell RC, Bell WO, Hertz GE, et al. Migration of a ventriculoperitoneal shunt into the pulmonary artery. *J Neurosurg Anesthesiol.* 1994; 6(2): 132–134, doi: [10.1097/00008506-199404000-00010](https://doi.org/10.1097/00008506-199404000-00010), indexed in Pubmed: [8012173](https://pubmed.ncbi.nlm.nih.gov/8012173/).
- Holden TM, Crow ML. Free intracardiac foreign body: A complication of ventriculovenous shunt for hydrocephalus. *J Thorac Cardiovasc Surg.* 1963; 45: 138–140, doi: [10.1016/S0022-5223\(19\)32904-6](https://doi.org/10.1016/S0022-5223(19)32904-6).
- Imamura H, Nomura M. Migration of ventriculoperitoneal shunt into the heart—case report. *Neurol Med Chir (Tokyo).* 2002; 42(4): 181–183, doi: [10.2176/nmc.42.181](https://doi.org/10.2176/nmc.42.181), indexed in Pubmed: [12013672](https://pubmed.ncbi.nlm.nih.gov/12013672/).
- Kubo S, Takimoto H, Takakura S, et al. Peritoneal shunt migration into the pulmonary artery—case report. *Neurol Med Chir (Tokyo).* 2002; 42(12): 572–574, doi: [10.2176/nmc.42.572](https://doi.org/10.2176/nmc.42.572), indexed in Pubmed: [12513032](https://pubmed.ncbi.nlm.nih.gov/12513032/).
- Chuang HL, Chang CN, Hsu JC. Minimally invasive procedure for ventriculoatrial shunt—combining a percutaneous approach with real-time transesophageal echocardiogram monitoring: report of six cases. *Chang Gung Med J.* 2002; 25(1): 62–66, indexed in Pubmed: [11926589](https://pubmed.ncbi.nlm.nih.gov/11926589/).
- Gonzalez LF, Kim L, ReKate HL, et al. Endovascular placement of a ventriculoatrial shunt. Technical note. *J Neurosurg.* 2007; 106(4 Suppl): 319–321, doi: [10.3171/ped.2007.106.4.319](https://doi.org/10.3171/ped.2007.106.4.319), indexed in Pubmed: [17465370](https://pubmed.ncbi.nlm.nih.gov/17465370/).
- Metellus P, Hsu W, Kharkar S, et al. Accuracy of percutaneous placement of a ventriculoatrial shunt under ultrasonography guidance: a retrospective study at a single institution. *J Neurosurg.* 2009; 110(5): 867–870, doi: [10.3171/2008.10.17674](https://doi.org/10.3171/2008.10.17674), indexed in Pubmed: [19099376](https://pubmed.ncbi.nlm.nih.gov/19099376/).

23. Vernet O, Rilliet B. Late complications of ventriculoatrial or ventriculoperitoneal shunts. *Lancet*. 2001; 358(9293): 1569–1570, doi: [10.1016/S0140-6736\(01\)06670-3](https://doi.org/10.1016/S0140-6736(01)06670-3), indexed in Pubmed: [11716880](https://pubmed.ncbi.nlm.nih.gov/11716880/).
24. Akhtar N, Khan AA, Yousaf M. Experience and outcome of ventricular atrial shunt: A multi centre study. *J Ayub Med Coll Abbottabad*. 2015; 27(4): 817–820, indexed in Pubmed: [27004330](https://pubmed.ncbi.nlm.nih.gov/27004330/).
25. James CA, McFarland DR, Wormuth CJ, et al. Snare retrieval of migrated ventriculoatrial shunt. *Pediatr Radiol*. 1997; 27(4): 330–332, doi: [10.1007/s002470050142](https://doi.org/10.1007/s002470050142), indexed in Pubmed: [9094241](https://pubmed.ncbi.nlm.nih.gov/9094241/).
26. So A, Shirani J. Pulmonary artery embolization of ventriculoatrial shunt fragment. *Tex Heart Inst J*. 2009; 36(2): 184–185, indexed in Pubmed: [19436823](https://pubmed.ncbi.nlm.nih.gov/19436823/).
27. Irie W, Furukawa M, Murakami C, et al. A case of V-A shunt catheters migration into the pulmonary artery. *Leg Med (Tokyo)*. 2009; 11(1): 25–29, doi: [10.1016/j.legalmed.2008.06.005](https://doi.org/10.1016/j.legalmed.2008.06.005), indexed in Pubmed: [18789874](https://pubmed.ncbi.nlm.nih.gov/18789874/).
28. Matsubara N, Miyachi S, Tsukamoto N. [Intracardiac migration of a ventriculoatrial shunt catheter treated by endovascular transvenous retrieval]. *No Shinkei Geka*. 2012; 40(6): 539–545, indexed in Pubmed: [22647514](https://pubmed.ncbi.nlm.nih.gov/22647514/).
29. Aloddadi M, Alshahrani S, Alnaami I. Endovascular retrieval of detached ventriculoatrial shunt into pulmonary artery in pediatric patient: case report. *J Pediatr Neurosci*. 2018; 13(1): 78–80, doi: [10.4103/JPN.JPN_143_17](https://doi.org/10.4103/JPN.JPN_143_17), indexed in Pubmed: [29899776](https://pubmed.ncbi.nlm.nih.gov/29899776/).
30. Xu J, Morchi G, Magge SN. Endovascular retrieval of a broken distal ventriculoatrial shunt catheter within the heart: illustrative case. *J Neurosurg Case Lessons*. 2021; 2(21): CASE21452, doi: [10.3171/CASE21452](https://doi.org/10.3171/CASE21452), indexed in Pubmed: [36060425](https://pubmed.ncbi.nlm.nih.gov/36060425/).
31. López Sánchez M, Domínguez Artiga MJ, Ortiz Lasa M, et al. Ventriculoatrial catheter migration. *Med Intensiva (Engl Ed)*. 2020 [Epub ahead of print]; 46(5): 296, doi: [10.1016/j.medin.2020.01.006](https://doi.org/10.1016/j.medin.2020.01.006), indexed in Pubmed: [32107034](https://pubmed.ncbi.nlm.nih.gov/32107034/).
32. Zervos TM, Kutschman K, Frisoli T, et al. Techniques for management and avoidance of ventriculoatrial shunt disconnection: illustrative cases. *J Neurosurg Case Lessons*. 2022; 3(22): CASE21654, doi: [10.3171/CASE21654](https://doi.org/10.3171/CASE21654), indexed in Pubmed: [35734610](https://pubmed.ncbi.nlm.nih.gov/35734610/).