

This is a provisional PDF only.



ISSN: 0029-540X

e-ISSN: 2300-2115

Current role of chemoembolization in the treatment of HCC

Authors: Rafał Kidziński, Grzegorz Kade, Krzysztof Pyra

DOI: 10.5603/njo.98621

Article type: Review paper

Submitted: 2023-12-20

How to cite:

Kidziński R, Kade G, Pyra K. Current role of chemoembolization in the treatment of HCC. NOWOTWORY J Oncol 2024; 74 (Ahead of print).

Accepted: 2024-03-06

Published online: 2024-04-09

This article has been peer reviewed and published immediately upon acceptance. It is an open access article, which means that it can be downloaded, printed, and distributed freely, provided the work is properly cited.

Current role of chemoembolization in the treatment of HCC

Rafał Kidziński¹, Grzegorz Kade¹, Krzysztof Pyra²

¹ *Clinical Hospital of the Ministry of Internal Affairs and Administration with the Warmia-Mazury Oncology Centre, Olsztyn, Poland*

² *Department of Interventional Radiology Medical University of Lublin, Lublin, Poland*

Hepatocellular carcinoma (HCC) accounts for 75% to 85% of primary liver cancers. Recent years have shown a significant increase in the incidence of HCC in Europe and the United States. The algorithm used most commonly in the treatment of HCC is the one developed in 1999 by Barcelona Clinic Liver Cancer (BCLC), updated from clinical trials. The last update is from 2022. Among the available treatments, depending on the stage of HCC, are liver transplantation, resection, thermal ablation, transarterial embolisation (TAE), transarterial chemoembolization (TACE), transarterial radioembolisation (TARE), stereotactic body radiation therapy (SBRT) as well as systemic treatment. The use of irreversible electroporation (IRE), a method involving disruption of cell membrane integrity is currently undergoing research. According to the BCLC, TACE is recommended for patients with BCLC stage-B (more than three lesions, preserved portal vein flow, preserved Child-Pugh A-B liver function and no extrahepatic lesions) and with BCLC stage 0 and stage 1 as an option after failure or not feasible for the first treatment option. In this article, we will try to explain in more detail what the chemoembolization method is and what the indications for its implementation are.

Key words: liver, embolisation, chemoembolization, transarterial chemoembolization, Barcelona Clinic Liver Cancer

Introduction

Hepatocellular carcinoma (HCC) accounts for 75% to 85% of primary liver cancers [1]. In Poland, there are between 2,000 and 3,000 new cases per year, while globally in 2020, HCC will account for around 900,000 new cases and around 830,000 deaths [2, 3]. HCC is the sixth most common cancer and -third/fourth most common cause of death among cancers [4, 5].

It is three times more frequent in men. Recent years have shown a significant increase in the incidence of HCC in Europe and the United States. Between 2000 and 2016, mortality from HCC in the United States increased by 43% [6]. HCC is associated with chronic liver disease and cirrhosis in 80–90% of cases. Major risk factors include hepatitis B and C, alcohol abuse, non-alcoholic steatohepatitis

(NASH), as well as diabetes, obesity and aflatoxin B1. It is estimated that approximately one-third of patients with cirrhosis may develop HCC with a one-year rate of 1–8% [7]. Elevated α -fetoprotein levels are found in 70–80% of patients with HCC.

There is also a variant of HCC – fibrolamellar carcinoma (FLC) – unrelated to cirrhosis, occurring mainly in young people with a slight predominance in women. This form has a different pathology and histopathology, and also a different prognosis. α -fetoprotein levels remain normal. The algorithm used most commonly in the treatment of HCC is one developed in 1999 by The Barcelona Clinic of Liver Cancer (BCLC), updated from clinical trials. The last update from 2022 is presented in figure 1 [8]. Among the available treatments, depending on the stage of HCC, are liver transplantation, resection, thermal ablation (microwave MWA, radiofrequency RFA and laser ablation), transarterial embolisation (TAE), transarterial chemoembolization (TACE), transarterial radioembolisation (TARE), stereotactic body radiation therapy (SBRT) as well as systemic treatment. The use of irreversible electroporation (IRE), a method involving disruption of cell membrane integrity [9, 10], is currently undergoing research. According to the BCLC, TACE (transarterial chemoembolization) is recommended for patients with BCLC stage-B (more than three lesions, preserved portal vein flow, preserved Child-Pugh A–B liver function and no extrahepatic lesions) [11–15], and with BCLC stage 0 and stage 1 as a option after failure or not feasible for the first treatment option [8].

Treatment strategy for HCC

An understanding of the liver's vascularity and HCC is necessary for the correct choice of treatment strategy for HCC. Healthy liver parenchyma is nourished approximately 75% from the portal vein branch, with only the remainder coming from the hepatic artery branch [16]. The terminal branches of the hepatic artery are divided into two sections. The first section accompanies the portal vessels and supplies the peribiliary vascular plexus (PBP), the interstitial tissue of the portal system and the walls of the portal vessels. The second section, named the isolated artery, penetrates the liver parenchyma independently of the portal vein branch. In the cirrhotic liver, PBPs are more developed which provides a link between the arterial and portal systems, and favors tumour survival in the event of arterial occlusion. The development of HCC in a cirrhotic liver progresses in several stages from a regenerative nodule undergoing transformation initially to a dysplastic nodule with a low and then high degree of dysplasia. In subsequent stages, a foci of HCC, known as nodule-in-nodule, appears within the dysplastic nodule to eventually progress to a large HCC. With this process, the proportions of vascularization change – the role of the portal system gradually declines in favor of the arterial system. In poorly and moderately differentiated HCC, portal vascularization almost completely disappears [17]. HCC can grow in an expanding, infiltrating or mixed form. The first form is

encapsulated and compresses the surrounding parenchyma, while the second form is poorly differentiated without a capsule with blurred outlines. This differentiation makes some HCCs, especially encapsulated, well-differentiated and extracapsular infiltrating HCCs having partially preserved portal vascularization.

The mode of enhancement has an obvious impact on HCC characteristics in imaging studies. In patients at risk, LI-RADS criteria are used in the assessment of liver lesions. These take into account lesion size, non-rim arterial phase hyperenhancement (APHE), non-peripheral washout, enhancing capsule and threshold growth. Using the above as a basis, the lesion can be assigned to one of the groups from LI-RADS 1, defined as definitely benign, to LI-RADS 5, defined as definitely HCC [18].

The first reports of hepatic artery embolisation in the treatment of hepatic cancers date back to 1974. In the 1970s, the first doses of chemotherapeutic agents were administered via the hepatic artery, and results showed that even single procedures gave better results than multiple cycles of systemic therapy [19–22]. There are currently two types of TACE procedures resulting from the embolisation material used. Conventional TACE (cTACE) in which the chemotherapeutic agent is mixed with Lipiodol – an oily, thick contrast agent to act as a drug carrier.

Drug eluting bead TACE (DEB TACE) – drug-soaked microspheres which, when injected into the vasculature, close the vasculature and then release the chemotherapeutic agent into the tumour in a controlled manner. The microspheres require the addition of a contrast agent to visualize the mixture.

Both procedures can be performed using a standard microcatheter or with a balloon-tipped microcatheter that, when inflated, changes regional hemodynamic conditions in the catheterized vessel or can be used as a safeguard against reflux. This method is called balloon occluded TACE (b-TACE). There is a difference in the distribution of embolisation material in cTACE and DEB-TACE. In cTACE, the emulsion, formed at a ratio of one part chemotherapeutic agent to two parts Lipiodol, selectively injected into the arteries is initially deposited in the tumor's sinusoids and then passes into the tumor's draining vessels on the side of the portal system and, via PBP, enters the portal system of healthy liver tissue in the tumor's immediate vicinity and into the arterial anastomoses [23–25]. This results in the prevention of flow reversal in the outflow pathway, the tumors necrosis and the increased margin of healthy liver surrounding the tumour. There is also an opportunity to potentially identify other tumour feeding routes that were not originally visible [26]. 31 In the case of richly vascularized lesions, where the mixture's full dose is not sufficient to close the tumor's vascular bed, embolisation can be completed using particles or Spongostan. As this mechanism also causes necrosis of healthy hepatocytes surrounding the tumour, ultra-selective embolisation of the feeding vessels to minimize liver damage is very important [27].

DEB-TACE involves injecting embolisation material saturated with a chemotherapeutic agent (usually doxorubicin but also epirubicin, mitomycin, cisplatin) through a catheter directly into the branch of the hepatic artery feeding the tumour. In DEB-TACE, it is possible to select the size of microspheres (from 40 μ m to 900 μ m). Smaller microspheres result in more peripheral vascular closure (i.e. closer to the tumor's centre) resulting in better deposition of the chemotherapeutic agent, but also significantly more necrosis of the liver parenchyma compared to the procedure performed with larger particles [28]. With smaller microspheres, there may also be an increased risk of biliary necrosis and blockages outside the liver. At the same time, microspheres are unable to block outflow from the tumour in DEB-TACE. Closing only the arterial vessels enables reverse flow to be generated from the surrounding hepatic sinuses and portal veins to the tumor's peripheral part. Arterial micro-anastomoses can also be difficult to block. Peripheral tumour tissues can therefore survive due to retained vascularization. Admittedly, the chemotherapeutics released from the microspheres in DEB-TACE can induce necrosis of surviving tumour cells, but this requires depositing them close to the living part of the tumour. DEB TACE also causes more arterial damage than cTACE and a higher risk of arteriovenous fistulae [29–31].

The procedure is performed under local anaesthesia with fluoroscopy guidance. After a percutaneous puncture of the femoral or radial artery, the interventional radiologist inserts a vascular sheath 5 Fr (2 mm in diameter) to prevent blood loss while providing access for subsequent instruments. A guidewire and catheter of appropriate curvature are inserted through the sheath, with fluoroscopy guidance, obtained with an angiographic apparatus. As an a-traumatic tool, the guidewire allows for safe navigation through the vascular system while providing guidance for the catheter, through which the contrast agent is administered. Aortic nephrography is performed first to assess possible routes for feeding the lesion. The visceral trunk is catheterized first, followed by the common hepatic artery. Angiography is performed by administering 25 ml of contrast for 5 seconds. This allows for an accurate assessment of the liver's vascular bed and the tumor's vascularization. If the vascularization is not complete, arteriography of the superior mesenteric artery is also performed in search of the right hepatic artery. This is the most common anatomical variation. Once the vessels feeding the HCC have been identified, the catheter tip is inserted as close to the tumour as possible using a micro-catheter, while avoiding the vessels feeding the healthy liver parenchyma. Once the micro-catheter's correct location is confirmed, a slow infusion of embolisation material (beads soaked in a cytostatic agent) mixed with contrast begins, thereby enabling observation of the material's distribution. Chemoembolization using slow-release drug particles produces a synergistic effect: it closes or reduces the arterial blood supply to the tumour with simultaneous deposition of the chemotherapeutic agent in the tumour area and reduced washout.

Depending on the number, size and degree of vascularization of the lesions, the authors perform 1 to 3 procedures at intervals of 4–6 weeks per TACE cycle. A follow-up examination is performed after the last procedure, preferably using the same technique as the eligibility examination. MRI is the preferred method. If there is no enhancement after embolisation and the tumour regresses, a follow-up examination is performed after another 3 months.

If enhancement of the residual tumour tissue is visualized, thermal ablation is used or further TACE sessions are performed, depending on the tissue's extent and availability. Two thermal ablation systems can be used: Emprint Medtronic (tmVA) and Echo Laser Elesta. In BCLC stage A patients, a complementary TACE procedure, after thermal ablation of lesions with borderline indications, is used. The efficacy of such combination therapy is confirmed in the available literature [32–37].

The causes of TACE failure and incomplete tumor necrosis can be divided into two groups. The first group includes reasons related to the technical side of the procedure. These include: incomplete, overly rapid embolisation which results in compaction of the embolisation material and blood supply into the vessels proximally feeding the lesion. Another reason may be the catheterization of the abnormal vessel (this occurs when tumors have a poor vascularization) or embolisation of not all the vessels feeding the lesion, particularly marginal, subcostal lesions, where additional feeding may come from arterial anastomoses or from extrahepatic arteries, e.g. from the internal thoracic or diaphragmatic artery, which is usually given off directly from the aorta. The second group can be described as dependent on the form of HCC. A proportion of HCCs, especially encapsulated, well-differentiated and extracapsular infiltrating HCCs have partially preserved portal vascularization.

In other cases, arterial inflow closure may result in portal vascularization of the tumour due to reversed flow in the small vessels on the portal system side and in the surrounding hepatic sinusoids [38–40]. Although TACE enables obtaining high concentrations of chemotherapeutic agents in the tumor not achievable with systemic treatment and relatively low concentrations outside the tumour area, it is the ischaemia caused by embolisation that contributes significantly to HCC necrosis [41].

The mRECIST criteria, in which areas undergoing contrast enhancement are considered as a viable tumour, are adopted to assess the response to treatment [42]. This is of great importance, as necrosis caused by TACE often leads to tumour swelling and an increase in tumour size which can be incorrectly treated as progression. Unintentional chemoembolization of a healthy part of the liver, and a concentrated dose of the cytostatic agent can lead to local liver damage and the formation of perfusion lesions in the healthy part of the liver, or lesions that mimic new foci.

Hence, it is extremely important that imaging examinations are evaluated by radiologists who are familiar with the specifics of the procedures and are members of multidisciplinary teams.

The efficacy of both TACE and also TAE methods has been evaluated in a number of studies.

In a five-year follow-up of 173 patients treated with DEB-TACE with Child-Pugh class A/B (102/71 [59/41%]), and mean lesion diameter 7.6 ± 2.1 cm, Malagari and her team obtained the following results: Overall survival at 1, 2, 3, 4, and 5 years was 93.6, 83.8, 62, 41.04, and 22.5 %, with higher rates achieved in Child class A compared with Child class B patients. Mean overall survival was 43.8 months (range 1.2–64.8). Cumulative survival was better for Child class A compared to Child class B patients ($p = 0.029$). For patients with dominant lesions ≥ 5 cm 1-, 2-, 3-, 4-, and 5-year survival rates were 100, 95.2, 71.4, 66.6, and 47.6 % for Child class A and 94.1, 88.2, 58.8, 41.2, 29.4, and 23.5 % for Child class B patients. Regarding DEB-DOX treatment, multivariate analysis identified a number of lesions ($p = 0.033$), lesion vascularity ($p < 0.0001$), initially achieved complete response ($p < 0.0001$), and objective response ($p = 0.046$) as significant and independent determinants of 5-year survival (43).

The PRECISION V study compared cTACE with DEB-TACE. The microsphere treated group showed higher rates of complete response (27% vs. 22%), objective response (52% vs. 44%) and disease control (63% vs. 52%) compared to the cTACE treated group. The hypothesis of a DEB TACE advantage was not confirmed (unilateral $p = 0.11$). Nevertheless, patients with cirrhosis and Child-Pugh class B, ECOG 1 performance, lesions in both lobes of the liver and disease recurrence showed a significant increase in objective response ($p = 0.038$) compared to cTACE. The use of microspheres was associated with improved tolerability, a significant reduction in severe liver toxicity ($p < 0.001$) and a significantly lower rate of doxorubicin-related side effects ($p = 0.0001$) [44].

In a randomized controlled trial (RCT) conducted between 1996 and 2000, Llovet and his team compared the efficacy of TAE, TACE and conservative treatment. Of the 903 patients, 112 were eligible for the study. Survival probabilities at 1 year and 2 years were 75% and 50% for embolisation; 82% and 63% for chemoembolization, and 63% and 27% for control (chemoembolization vs control $p = 0.009$). chemoembolization induced objective responses sustained for at least 6 months in 35% [14] of cases, and was associated with a significantly lower rate of portal-vein invasion than conservative treatment [45].

The systematic review and meta-analysis presented by Bzeizi and co-authors included 34 studies involving 4,841 patients with HCC, and an average follow-up period of 1.5 to 18 months. There were no significant differences between DEB-TACE and C-TACE in terms of complete response, partial response and disease stability. However, disease control (OR: 1.42, 95% CI: 1.03, 1.96) and objective response (OR: 1.33, 95% CI: 0.99, 1.79) were significantly more successful with DEB-TACE treatment with fewer major complications and overall mortality. A pooled analysis showed no superiority of DEB-TACE in terms of complete or partial response, disease stability, disease progression control or mortality at 30 days or at the end of the study [46].

However, the results showed that DEB-TACE was associated with better objective response, disease control and lower overall mortality compared to C-TACE treatment with fewer major complications. DEB-TACE shows less systemic exposure to the chemotherapeutic agent. Furthermore, it shows a standardized release of the chemotherapeutic agent from microspheres, resulting in prolonged retention in the tumor as well as lower liver toxicity. An important aspect is the ability to select the size of the microspheres.

Our extensive experience also shows the advantage of DEB-TACE in terms of controlling the rate and volume of microspheres administered [44].

In vivo studies performed on pigs have shown the spread of doxorubicin to a distance of 600 μ m from the edge of the microsphere, with a very rapid decrease in the first 100–200 μ m around the particle, and a very slow decrease in the next 400 μ m. A sudden drop in drug concentration suggests the presence of barriers to drug diffusion [47]. Particles released 43% of the initial doxorubicin load within the first month and 89% within 3 months of the procedure, consistent with *in vitro* tests predicting a 50% release within 2–3 months [48, 49].

However, it should be noted that the above study took place on healthy pig livers without a tumor. HCC occurring in a cirrhotic liver has a different vascularization from healthy tissue, and the permeability and sensitivity of tumor cells to doxorubicin is also different [50–52]. The above work suggests that when deciding on the type of TACE (cTACE vs. DEB-TACE), an in-depth analysis of imaging examinations, in particular, is required to optimally select the procedure technique due to the heterogeneity of the BCLC B group. Despite the clear advantages of DEB-TACE, some authors identify groups of patients in whom they prioritize cTACE.

Adverse effects associated with TACE include post-embolism syndrome, which is the body's natural response to tumor embolisation. It can manifest in a number of ways: abdominal pain, raised body temperature, vomiting or temporary deterioration of liver function. The duration of symptoms is highly individual, ranging from 2–3 days to 2 weeks. The incidence ranges from 5 to about 22% [53]. It is important to adequately provide patients with painkillers. More serious complications include liver abscesses requiring drainage, acute pancreatitis or acute cholecystitis, liver failure, kidney failure. Their incidence ranges from 2% to 4% [53]. Vascular dissection and punctures are even rarer.

Monier et al. in their study assess adverse effects of forming biloma, portal vein thrombosis, portal vein branch narrowing, and bile duct dilatation. They assess incidence range up to 5% and for global hepatic damages up to 15% [54]. In order to detect potential side effects quickly, patients require regular monitoring after TACE, especially of liver parameters. Due to the contrast agent used during the procedure, contrast-induced nephropathy should be excluded in patients at risk. It is defined as an increase in creatinine concentration by ≥ 0.5 mg/dl or more than 25% from the baseline within 2–3 days of contrast agent administration. The evaluation of liver function is done according to

the Child-Pugh scale correlated with the pre-treatment results. The ALBI score also shows great usefulness in post-treatment evaluation [55].

Conclusions

In case of TACE failure and disease progression at BCLC stage C and Child-Pugh liver stage A–B, the patient receives systemic treatment. Systemic therapy should be considered as a first line over TACE in patients where: HCC exceeds “up to seven” criteria, tumor(s) is/are larger than 5 cm, contiguous multinodular tumors, poorly differentiated or undifferentiated HCC and if there is no objective response after two consecutive TACE treatments [56].

Sorafenib was initially used, being the first multi-kinase inhibitor available for the treatment of advanced HCC. Currently, a atezolizumab and bevacizumab combination is the preferred treatment method, superior to Sorafenib, and demonstrating prolonged overall survival. On the other hand, in the presence of sorafenib contraindications, lenvatinib remains the preferred drug of choice. Second-line treatment includes using regorafenib, cabozantinib, nivolumab, pembrolizumab, ramucirumab and combination therapies [57].

Article information and declarations

Conflicts of Interest

None declared

Krzysztof Pyra

Medical University of Lublin

Department of Interventional Radiology Medical University of Lublin

ul. Jaczewskiego 8

20-954 Lublin, Poland

e-mail: k.pyra@poczta.fm

Received: 20 Dec. 2023

Accepted: 6 Mar 2024

References

1. Smoter P, Kotulski M, Smoter K et al. Liver transplantation in primary liver tumors. *NOWOTWORY J Oncol* 2023; 73: 248–254.
2. Ferlay J, Ervik M, Lam F et al. *Global Cancer Observatory: Cancer Today*; International Agency for Research on Cancer: Lyon, France, 2018; pp. 1–6.
3. Sung H, Ferlay J, Siegel R.L et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J. Clin.* 2021, 71, 209–249.
4. Villanueva A Hepatocellular carcinoma. *N. Engl. J. Med.* 2019, 380, 1450–1462.
5. Forner et al. "Hepatocellular carcinoma *Lancet* 2018; 391: 1301–14 Published Online January 4, 2018 [http://dx.doi.org/10.1016/S0140-6736\(18\)30010-2](http://dx.doi.org/10.1016/S0140-6736(18)30010-2)
6. Centers for Disease Control and Prevention, National Center for Health Statistics. Trends in liver cancer mortality among adults aged 25 and over in the United States, 2000–2016. July 2018 (<https://www.cdc.gov/nchs/products/databriefs/db314.htm>).
7. Ioannou G.N, Green P, Lowy E et al. Differences in hepatocellular carcinoma risk, predictors and trends over time according to etiology of cirrhosis. *PLoS ONE* 2018, 13, e0204412.
8. Reig M, Forner A, Rimola J et al. BCLC strategy for prognosis prediction and treatment recommendation: The 2022 update *Journal of Hepatology* 2022 vol.76 681–693
9. Bhutiani N, Philips P, Scoggins CR et al. Evaluation of tolerability and efficacy of irreversible electroporation (IRE) in treatment of Child-Pugh B (7/8) hepatocellular carcinoma (HCC). *HPB* 2016;18(7):593e9.
10. Lencioni R, Cioni D, Della Pina MC et al. New options for image-guided ablation. *J Hepatobiliary Pancreat Sci* 2010;17(4):399e403.
11. Llovet JM, Real MI, Montana X et al. (2002) Barcelona Liver Cancer Group. Arterial embolisation or chemoembolization versus symptomatic treatment in patients with unresectable hepatocellular carcinoma: a randomized controlled trial. *Lancet* 359(9319):1734–1739
12. Lo C.-M, Ngan H, Tso W-K et al. Randomized controlled trial of transarterial lipiodol chemoembolization for unresectable hepatocellular carcinoma. *Hepatology* 2002, 35, 1164–1171. [CrossRef]
13. Cammà C, Schepis F, Orlando A et al. Transarterial chemoembolization for unresectable hepatocellular carcinoma: Meta-analysis of randomized controlled trials. *Radiology* 2002, 224, 47–54.

14. Llovet J.M, Bruix J Systematic review of randomized trials for unresectable hepatocellular carcinoma: Chemoembolization improves survival. *Hepatology* 2003, 37, 429–442.
15. Huang Y-H, Chen C-H, Chang T-T et al. The role of transcatheter arterial embolization for patients with unresectable hepatocellular carcinoma: A nationwide, multicentre study evaluated by cancer stage. *Aliment. Pharmacol. Ther.* 2005, 21, 687–694.
16. Matsui O, Kobayashi S, Sanada J et al. Hepatocellular nodules in liver cirrhosis: Hemodynamic evaluation (angiography-assisted CT) with special reference to multi-step hepatocarcinogenesis. *Abdom. Imaging* 2011, 36, 264–272.
17. Miyayama S, Treatment Strategy of Transarterial Chemoembolization for Hepatocellular Carcinoma. *Appl. Sci.* 2020, 10, 7337.
18. American College of Radiology Committee on LI-RADS® (Liver) Available at:
<https://www.acr.org/-/media/ACR/Files/Clinical-Resources/LIRADS/LI-RADS-2018-Manual-5Dec18.pdf>
19. Doyon D et al. "Hepatic, arterial embolization in patients with malignant liver tumours, *Annales de Radiologie* 17, 6, 593–603, 1974
20. Friedman M et al. "Therapy for hepatocellular cancer with intrahepatic arterial adriamycin and 5-fluorouracil combined with whole-liver irradiation: a Northern California Oncology Group study, *Cancer Treatment Reports*, vol. 63, no. 11-12, pp. 1885–1888, 1979.
21. Misra N. C et al. "Intrahepatic arterial infusion of combination of mitomycin C and 5 fluorouracil in treatment of primary and metastatic liver carcinoma, *Cancer*, vol. 39, no. 4, pp. 1425–1429, 1977.
22. Hirose H. et al. Chemotherapy of hepatocellular carcinoma—with special reference to one-shot intra-arterial infusion of a high dose of Adriamycin. *Gan To Kagaku Ryoho*, vol. 9, no. 12, pp. 2216–2221, 1982.
23. Terayama N, Matsui O, Gabata T et al. Accumulation of iodized oil within the nonneoplastic liver adjacent to hepatocellular carcinoma via the drainage routes of the tumor after transcatheter arterial embolization. *Cardiovasc. Intervent. Radiol.* 2001, 24, 383–387.
[CrossRef]
24. Miyayama S, Matsui O, Yamashiro M et al. Ultraslective transcatheter arterial chemoembolization with a 2-F tip microcatheter for small hepatocellular carcinomas: Relationship between local tumor recurrence and visualization of the portal vein with iodized oil. *J. Vasc. Interv. Radiol.* 2007, 18, 365–376.

25. Miyayama S, Yamashiro M, Okuda M et al. Anastomosis between the hepatic artery and the extrahepatic collateral or between extrahepatic collaterals: Observation on angiography. *J. Med. Imag. Radiat. Oncol.* 2009, 53, 271–282. [CrossRef] [PubMed]
26. Miyayama S, Matsui O Superselective conventional transarterial chemoembolization for hepatocellular carcinoma: Rationale, technique, and outcome. *J. Vasc. Interv. Radiol.* 2016, 27, 1269–1278
27. Yoshida K, Matsui O, Miyayama S, et al Isolated arteries originating from the intrahepatic arteries: Anatomy, function, and importance in intervention. *J. Vasc. Interv. Radiol.* 2018, 29, 531–537.e1.
28. Namur J, Wassef M, Millot JM et al. A. Drug-eluting beads for liver embolization: concentration of doxorubicin in tissue and in beads in a pig model. *J Vasc Interv Radiol.* 2010 Feb;21(2):259-67. doi: 10.1016/j.jvir.2009.10.026. Erratum in: *J Vasc Interv Radiol.* 2010 Apr;21(4):596. PMID: 20123210.
29. Minamiguchi K, Tanaka T, Nishiofuku H et al. Comparison of embolic effect between water-in-oil emulsion and microspheres in transarterial embolization for rat hepatocellular carcinoma model. *Hepatol. Res.* 2020.
30. Miyayma S, Matsui O Applying superselective conventional TACE. *Endovasc. Today* 2017, 16, 52–56.
31. Shimose S, Iwamoto H, Tanaka M et al. Increased arterio-portal shunt formation after drug-eluting beads TACE for hepatocellular carcinoma. *Oncology* 2020, 98, 558–565.
32. Helmberger T, Dogan S, Straub G et al. Liver resection or combined chemoembolization and radiofrequency ablation improve survival in patients with Hepatocellular carcinoma. *Digestion* 2007;75(2e3):104e12.
33. Lu Z, Wen F, Guo Q et al. Radiofrequency ablation plus chemoembolization versus radiofrequency ablation alone for hepatocellular carcinoma: a meta-analysis of randomized-controlled trials. *Eur J Gastroenterol Hepatol* 2013;25(2):187e94.
34. Lencioni R, Crocetti L, Petruzzi P et al. Doxorubicin-eluting bead-enhanced radiofrequency ablation of hepatocellular carcinoma: a pilot clinical study. *J Hepatol* 2008;49(2):217e22.
35. Iezzi R, Pompili M, Posa A et al. Combined locoregional treatment of patients with hepatocellular carcinoma: State of the art. *World J Gastroenterol* 2016;22(6):1935e42.
36. Thamtorawat S, Hicks RM, Yu J et al. Preliminary outcome of microwave ablation of hepatocellular carcinoma: breaking the 3-cm barrier? *J Vasc Interv Radiol* 2016;27(5):623e30.

37. Zhang NN, Lu W, Cheng XJ et al. High-powered microwave ablation of larger hepatocellular carcinoma: evaluation of recurrence rate and factors related to recurrence. *Clin Radiol* 2015;70(11):1237e43.
38. Goseki N, Nosaka T, Endo M et al. Nourishment of hepatocellular carcinoma cells through the portal blood flow with and without transcatheter arterial embolization. *Cancer* 1995, 76, 736–742. [CrossRef]
39. Ekelund L, Lin G, Jeppsson B Blood supply of experimental liver tumors after intraarterial embolization with Gelfoam powder and absolute ethanol. *Cardiovasc. Intervent. Radiol.* 1984, 7, 234–239.
40. Yoshida K, Matsui O, Miyayama S et al. Isolated arteries originating from the intrahepatic arteries: Anatomy, function, and importance in intervention. *J. Vasc. Interv. Radiol.* 2018, 29, 531–537.e1.
41. Kluger M et al. Bland Embolization Versus Chemoembolization of Hepatocellular Carcinoma Before Transplantation *Liver Transpl.* 2014 May ; 20(5): 536–543. doi:10.1002/lt.23846.
42. Gregory J, Dioguardi Burgio M, Corrias G et al. Evaluation of liver tumour response by imaging. *JHEP Rep.* 2020 Apr 28;2(3):100100. doi: 10.1016/j.jhepr.2020.100100. PMID: 32514496; PMCID: PMC7267412.
43. Malagari K, Pomoni M, Moschouris H et al. Chemoembolization With Doxorubicin-Eluting Beads for Unresectable Hepatocellular Carcinoma: Five-Year Survival Analysis. *Cardiovasc Intervent Radiol* 35, 1119–1128 (2012). <https://doi.org/10.1007/s00270-012-0394-0>
44. Lammer J, Malagari K, Vogl T et al. Prospective randomized study of doxorubicin-eluting-bead embolization in the treatment of hepatocellular carcinoma: Results of the PRECISION V study. *Cardiovasc. Intervent. Radiol.* 2010, 33, 41–52
45. Llovet JM, Real MI, Montana X et al. (2002) Barcelona Liver Cancer Group. Arterial embolisation or chemoembolization versus symptomatic treatment in patients with unresectable hepato- cellular carcinoma: a randomized controlled trial. *Lancet* 359(9319):1734–1739
46. Bzeizi K.I, Arabi M, Jamshidi N et al. Conventional Transarterial Chemoembolization Versus Drug-Eluting Beads in Patients with Hepatocellular Carcinoma: A Systematic Review and Meta-Analysis. *Cancers* 2021, 13, 6172. <https://doi.org/10.3390/cancers13246172>

47. Namur J, Wassef W, Millot J_M et al. Drug-Eluting Beads for Liver Embolization: Concentration of Doxorubicin in Tissue and in Beads in a Pig Model. *Journal of Vascular and Interventional Radiology* 21, nr 2 (luty 2010): 259–67.
48. Gonzalez MV, Tang Y, Phillips GJ et al. Doxorubicin eluting beads—2: methods for evaluating drug elution and in-vitro: in-vivo correlation. *J Mater Sci Mater Med* 2008; 19:767–775.
49. Lewis AL, Gonzalez MV, Lloyd AW et al. DC bead: in vitro characterization of a drug-delivery device for transarterial chemoembolization. *J Vasc Interv Radiol* 2006; 17:335–342.
50. Nakashima T, Kojiro M Pathologic characteristics of hepatocellular carcinoma. *Semin Liver Dis* 1986; 6:259 –266.
51. Jain RK Transport of molecules in 45. the tumor interstitium: a review. *Cancer Res* 1987; 47:3039–3051.
52. Minchinton AI, Tannock IF Drug penetration in solid tumours *Nat Rev Cancer* 2006; 6:583–592.
53. Marcacuzco Quinto A, Nutu O-A, San Roman Manso R et al. Complicaciones de la quimioembolizacion transarterial (QETA) en el tratamiento de los tumores hepaticos. *Cir Esp.* 2018; 96:560–567
54. Monier, A., Guiu, B., Duran, R. et al. Liver and biliary damages following transarterial chemoembolization of hepatocellular carcinoma: comparison between drug-eluting beads and lipiodol emulsion. *Eur Radiol* 27, 1431–1439 (2017). <https://doi.org/10.1007/s00330-016-4488-y>
55. Johnson P.J, Berhane S, Kagebayashi C et al. Assessment of liver function in patients with hepatocellular carcinoma: A new evidence-based approach— The ALBI grade. *J. Clin. Oncol.* 2015, 33, 550–558.
56. Silk T, Silk M, Wu J. Up to seven criteria in selection of systemic therapy for hepatocellular carcinoma. *World J Gastroenterol.* 2022;28(23):2561-2568. doi:10.3748/wjg.v28.i23.2561
57. Finn R.S, Qin S, Ikeda M et al. Atezolizumab plus Bevacizumab in Unresectable Hepatocellular Carcinoma. *N. Engl. J. Med.* 2020, 382, 1894–1905.




Figure 1. Barcelona Clinic Liver Cancer (BCLC) – updated from clinical trials (2022)