

World Tobacco Quitting Day 2020 – the united voice of Polish experts on tobacco prevention and control

Paweł Koczkodaj¹, Joanna Didkowska^{1,2}, Łukasz Balwicki³, Magdalena Cedzyńska¹, Marzena Dominiak^{4,5}, Wojciech Hanke^{6,7}, Piotr Jankowski⁸, Jacek Jassem⁹, Grzegorz Juszczyk^{10,11}, Maciej Krzakowski¹², Jarosław Pinkas^{13,14}, Irena Przepiórka¹, Krzysztof Przewoźniak¹, Piotr Rutkowski^{15,16}, Daniel Śliż¹⁷, Stefan Wesołowski¹⁸, Tomasz Zdrojewski¹⁹, Jan Walewski²⁰

¹Department of Epidemiology and Primary Cancer Prevention, Maria Skłodowska-Curie National Research Institute of Oncology, Warsaw, Poland

²National Cancer Registry, Maria Skłodowska-Curie National Research Institute of Oncology, Warsaw, Poland

³Department of Public Health & Social Medicine, Medical University of Gdansk, Poland

⁴Department of Oral Surgery, Wrocław Medical University, Poland

⁵Polish Dental Society, Wrocław, Poland

⁶Department of Environmental Epidemiology, Nofer Institute of Occupational Medicine in Lodz, Poland

⁷Public Health Committee, Polish Academy of Science, Warsaw, Poland

⁸Department of Cardiology, Interventional Electrophysiology and Hypertension, Institute of Cardiology, Jagiellonian University Medical College, Krakow, Poland

⁹Department of Oncology and Radiotherapy at the Medical University of Gdansk, Poland

¹⁰National Institute of Public Health-National Institute of Hygiene, Warsaw, Poland

¹¹Department of Public Health, Medical University of Warsaw, Poland

¹²Department of Lung Cancer and Thoracic Tumors, Maria Skłodowska-Curie National Research Institute of Oncology, Warsaw, Poland

¹³Chief Sanitary Inspectorate of the State Sanitary Inspection, Warsaw, Poland

¹⁴School of Public Health, Centre of Postgraduate Medical Education, Warsaw, Poland

¹⁵Department of Soft Tissue/Bone Sarcoma and Melanoma, Maria Skłodowska-Curie National Research Institute of Oncology, Warsaw, Poland

¹⁶Committee for the National Oncological Strategy, Ministry of Health, Warsaw, Poland

¹⁷Department of Internal Diseases and Cardiology, Medical University of Warsaw, Poland

¹⁸National Institute of Tuberculosis and Lung Diseases, Warsaw, Poland

¹⁹Department of Preventive Medicine and Education, Medical University in Gdansk, Poland

²⁰Department of Lymphoid Malignancies, Maria Skłodowska-Curie National Research Institute of Oncology, Warsaw, Poland

On November 19th, 2020 we will celebrate annual World Tobacco Quitting Day. On this occasion, we would like call to action for a stronger and united tobacco prevention activities in Poland.

Despite many successes in tobacco prevention and control, use of tobacco products in Poland is still a great public health challenge. Whereas the prevalence of traditional cigarettes smoking is declining, new tobacco products are more and more popular – particularly among young people. It has been estimated that about 28% of boys and

18.6% of girls aged 13–15 years in Poland use e-cigarettes. Moreover, in the same age group 15.6% of boys and 14.9% of girls smoke traditional cigarettes [1]. Considering adult population, e-cigarettes are less popular – about 4% of men and about 1% of women use this particular tobacco product [2]. However, the use of traditional cigarettes is much higher – 26% and 17% of Polish men and women, respectively, are regular smokers [3].

Being aware of overwhelming influence of SARS-CoV-2 pandemic on health care systems, we cannot ignore impact

How to cite:

Koczkodaj P, Didkowska J, Balwicki Ł, Cedzyńska M, Dominiak M, Hanke W, Jankowski P, Jassem J, Juszczyk G, Krzakowski M, Pinkas J, Przepiórka I, Przewoźniak K, Rutkowski P, Śliż D, Wesołowski S, Zdrojewski T, Walewski J. *World Tobacco Quitting Day 2020 – the united voice of Polish experts on tobacco prevention and control*. NOWOTWORY J Oncol 2020; 70: 278–279.

of other pandemic, which lasts for decades and is evolving just in front of our eyes. Only between 2011 and 2014, e-cigarettes use among young Poles has increased from 6% to 29.9% [4]. Compared to other EU-countries, Poland has one of the highest rates of e-cigarettes use among teenagers [1]. Moreover, Poland is among a few European Union (EU) countries with a very high level (17.5% to <20%) of tobacco attributable Disability-Adjusted Life Years (DALYs) [5]. For comparison, Romania and Czech Republic are in the group of countries with DALY between 15% to <17.5%, Germany and Spain 12.5% to <15%, and France and Italy 10% to <12.5%.

Considering these data, we appeal to all involved parties – researchers, health professionals, and stakeholder and policymakers to advocate for a stronger and more tailored anti tobacco law, health education and stable long lasting financing of such actions. We believe that we need to stand together to stop, or at least to slow down tobacco epidemic in Poland.

Conflict of interest: none declared

Paweł Koczkodaj

*Maria Skłodowska-Curie National Research Institute of Oncology
Cancer Epidemiology and Primary Prevention Department
ul. Wawelska 15B
02-034, Warszawa, Poland
e-mail: pawel.koczkodaj@pib-nio.pl*

Received and accepted: 6 Nov 2020

References

1. Centers for Disease Control and Prevention. National Center for Chronic Disease Prevention and Health Promotion, Office of Smoking and Health, Global Tobacco Surveillance System Data (GTSSData). <https://www.cdc.gov/tobacco/global/gtss/gtssdata/index.html> (2.11.2020).
2. Trząsalska A, Staszyńska M, Krassowska U. Raport z ogólnopolskiego badania ankietowego na temat postaw wobec palenia tytoniu. Kantar Public dla Głównego Inspektoratu Sanitarnego, Warszawa 2017.
3. Centrum Badań Opinii Społecznej. Komunikat z Badań nr 104/2019. Palenie papierosów. 2019; sierpień.
4. Balwicki L, Smith D, Balwicka-Szczyrba M, et al. Youth Access to Electronic Cigarettes in an Unrestricted Market: A Cross-Sectional Study from Poland. *Int J Environ Res Public Health*. 2018; 15(7), doi: 10.3390/ijerph15071465, indexed in Pubmed: 29997350.
5. Murray C, Aravkin A, Zheng P, et al. Global burden of 87 risk factors in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet*. 2020; 396(10258): 1223–1249, doi: 10.1016/s0140-6736(20)30752-2.