


## Solitary cutaneous focal mucinosis during immunotherapy for melanoma

Przemysław Baran<sup>1</sup>, Wojciech M. Wysocki<sup>1, 2, 3</sup>

<sup>1</sup>Department of General and Oncological Surgery, 5<sup>th</sup> Military Clinical Hospital, Krakow, Poland

<sup>2</sup>Chair of Surgery, Faculty of Medicine and Health Sciences, Andrzej Frycz Modrzewski Krakow University, Krakow, Poland

<sup>3</sup>Maria Skłodowska-Curie National Research Institute of Oncology, Warsaw, Poland



**Figure 1.** Microscopic presentation of hematoxylin and eosin-stained sections of a solitary cutaneous focal mucinosis on the right lower leg of a 40-year-old woman. Pallor of the dermis is a consequence of increased amounts of interstitial mucin. (A) x40, (B) x200

A 40-year-old woman was diagnosed with pT3b melanoma of the right thigh (fig. 1). The patient underwent wide local excision and sentinel node biopsy with clinically occult metastases; subsequently an inguinal lymphadenectomy was performed and adjuvant pembrolizumab introduced. Ten months since the start of immunotherapy, an asymptomatic, plateau-shaped, white-colored lesion of the right lower leg was noted. An excisional biopsy was taken and a diagnosis of “cutaneous focal mucinosis” was established. “Solitary cutaneous focal mucinosis” is a rare skin lesion characterized by the increased presence of mucin in the dermis and is mostly an idiopathic condition. The lesion typically presents as an asymptomatic dome-shaped papule or nodule on the extremities. The color varies from flesh-colored to white to red. It occurs in adults with a male predominance.

The lesion has been reported in less than 200 individuals. It is rarely clinically diagnosed due to variable morphologic presentation and the necessity of a surgical biopsy to establish the diagnosis [1]. Association of this infrequent skin lesion with immunotherapy has been previously observed in a very few cases [2]. Typically, a surgical biopsy provides adequate treatment of the solitary cutaneous focal mucinosis; additionally the lesion does not tend to reoccur. There are additional studies necessary.

### References

1. Gutierrez N, Erickson C, Calame A, et al. Solitary Cutaneous Focal Mucinosis. *Cureus*. 2021; 13(10): e18618, doi: 10.7759/cureus.18618, indexed in Pubmed: 34786224.
2. Lesiak A, Włodarczyk M, Sobolewska A, et al. Can biologic treatment induce cutaneous focal mucinosis? *Postepy Dermatol Alergol*. 2014; 31(6): 413–416, doi: 10.5114/pdia.2014.40944, indexed in Pubmed: 25610359.

### How to cite:

Baran P, Wysocki WM. *Solitary cutaneous focal mucinosis during immunotherapy for melanoma*. *NOWOTWORY J Oncol* 2023; 73: 109.

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.