

Increased uptake of technetium-99m methylene diphosphonate in muscles in the course of polymyositis

Zofia Niemir¹, Robert Oleksa², Rafał Czepczyński², Jerzy Sowiński²

¹Department of Nephrology, Poznan University of Medical Sciences, Poznań, Poland

²Nuclear Medicine Unit, Department of Endocrinology

[Received: 27 VI 2005; Accepted 19 VIII 2005]

Abstract

A case of a woman aged 46 years with signs of rhabdomyolysis and acute renal failure is presented. Coxsackie serum test was positive. Increased uptake of Technetium-99m methylene diphosphonate (99mTc-MDP) by muscles of thighs and calves was observed. After 1 year no increased accumulation of radiotracer in the muscles was found.

Keywords: bone scintigraphy, polymyositis

Case presentation

A woman aged 46 years was admitted to a department of nephrology because of acute renal failure. Serum level of creatinine was 9 mg/dl. She presented signs of rhabdomyolysis: creatine kinase (CK) — 100 000 IU/l (normal value: 26–140). After 5 weeks of treatment, serum level of creatinine was 1.2 mg/dl and serum level of CK was 120 IU/l. Coxsackie serum test was positive. Due to the elevated serum level of calcium, bone scintigraphy was performed. Increased uptake of Technetium-99m methylene diphosphonate (^{99m}Tc-MDP) by muscles of thighs and calves was observed (Figure 1). Diagnosis: polymyositis due to Cox-

Correspondence to: Robert Oleksa
 Nuclear Medicine Unit, Department of Endocrinology and Metabolism
 Przybyszewskiego 49, 60–355 Poznań, Poland
 Tel: (+48 61) 869 13 56, fax: (+48 61) 869 16 82
 e-mail: roleksa@amp.edu.pl




Figure 1. Bone scintigraphy using technetium-99m methylene diphosphonate (99mTc-MDP). Increased uptake of the radiotracer in the muscles of lower extremities.




Figure 2. Bone scintigraphy using technetium-99m methylene diphosphonate (^{99m}Tc -MDP) performed 1 year after the previous one. Activity of the radiotracer in the muscles is markedly diminished.

sackie infection. A follow-up scan was obtained after one year of oral steroid treatment. No increased accumulation of radiotracer in the muscles was found (Figure 2). Serum level of creatinine was 1.1 mg/dl, CK — 110 IU/l.

Increased uptake of radiotracer in the muscles can be found due to calcification, inflammation, certain intramuscular drug injections (e.g. iron supplementation, pethidine).

In this report, a case of inflammatory rhabdomyolysis was described. Increased uptake of ^{99m}Tc -MDP was found in the muscles of the lower extremities. This scintigraphic sign disappeared with a successful anti-inflammatory treatment.

As postulated by Nakayama et al., location of inflammatory myopathy can be detected using ^{99m}Tc -MDP scintigraphy. The scan can indicate the site of muscle biopsy [1].

Increased accumulation of ^{99m}Tc -MDP in the muscles was described in the case of muscle injury after cardiac resuscitation [2], rhabdomyolysis caused by drug intake (statins, heroin) [3–4].

Malignant disease may be connected with muscle uptake of the bone-seeking agents as well [5–7].

It should be remembered that excessive exertion may also cause unexpected uptake of ^{99m}Tc -MDP in the muscles [8–10].

References

1. Nakayama T, Saitoh Y, Yatabe K, Sueishi M, Kawai M. Diagnosis of inflammatory myopathy; usefulness of ^{99m}Tc MDP scintigraphy and muscle MRI for determination of affected sites. *Rinsho-Shinkeigaku* 1999; 39: 1114–1117.
2. Provan I, Murray C, Mansberg VJ, Rossleigh MA. Intense muscle uptake of Tc-99m MDP and Ga-67 citrate in massive rhabdomyolysis. *Clin Nucl Med* 1997; 22: 463–466.
3. Rhodes CA, Balon HR. Simvastatin-associated rhabdomyolysis demonstrated on bone scan. *Clin Nucl Med* 2002; 27: 793–794.
4. Lee BF, Chiu NT, Chen WH, Liu GC, Yu HS. Heroin-induced rhabdomyolysis as a cause of reflex sympathetic dystrophy. *Clin Nucl Med* 2001; 26: 289–292.
5. Low RD, Hicks RJ, Arkles LB, Gill G, Adam W. Progressive soft tissue uptake of Tc-99m MDP reflecting metastatic microcalcification. *Clin Nucl Med* 1992; 17: 658–662.
6. Batte WG Jr, Yeh SD, Rosenblum MK, Larson SM. Intense muscle uptake of Ga-67 citrate and Tc-99m MDP in a patient with aplastic anemia. *Clin Nucl Med* 1991; 16: 421–424.
7. Masaoka S, Fu T. Malignant lymphoma in skeletal muscle with rhabdomyolysis: a report of two cases. *J Orthop Sci* 2002; 7: 688–693.
8. Valk P. Muscle localization of Tc-99m MDP after exertion. *Clin Nucl Med* 1984; 9: 493–494.
9. Lafforgue P, Siles S, Daumen-Legre V, Acquaviva PC. An unexpected, benign cause of increased muscular uptake at bone scintigraphy. *Clin Exp Rheumatol* 1994; 12: 309–311.
10. du Cret RP, Boudreau RJ, Block JB, Jensen PR, Loken MK. Exercise-induced muscle uptake of technetium-99m MDP. *Clin Nucl Med* 1987; 12: 354–355.