

Alper Kahvecioglu¹, Sezin Yuce Sari¹, Hasan Cagri Yildirim², Zafer Arik², Melis Gultekin¹, Ferah Yildiz¹

¹Department of Radiation Oncology, Faculty of Medicine, Hacettepe University, Ankara, Turkey

²Department of Medical Oncology, Faculty of Medicine, Hacettepe University, Ankara, Turkey

Leptomeningeal metastasis in primary uterine cervical cancer: a rare case and review of the literature

Address for correspondence:

Assoc. Prof. Sezin Yuce Sari, MD
 Department of Radiation Oncology,
 Faculty of Medicine, Hacettepe University
 06100, Sıhhiye, Ankara, Turkey
 phone: +90-312-3052900
 fax: +90-312-3092914
 e-mail: sezin.yuce@hacettepe.edu.tr

ABSTRACT

Objectives. Leptomeningeal metastasis (LM) of primary uterine cervical cancer is rare and treatment options are limited. In this case report and literature review, we aimed to present a patient with cervical cancer with LM and discuss previously reported cases in the literature.

Case presentation. Our case was a 58-year-old patient who was initially diagnosed with metastatic primary uterine cervical cancer and treated with chemotherapy and chemoradiotherapy. During follow-up, she developed neurological symptoms, and LM was detected in the craniospinal regions. Cerebrospinal fluid cytology examination has confirmed metastatic disease. The patient was treated with concurrent intrathecal methotrexate and whole-brain radiotherapy (WBRT). A good clinical and cytological response was obtained. However, while intrathecal methotrexate was being continued after WBRT, she succumbed to hematological toxicity before the radiological response could be evaluated.

Conclusions. LM is an extremely rare and catastrophic distant spread pattern in patients with cervical cancer. In the literature, a total of 26 patients were reported up to date. Median survival after detection of LM was nine weeks, including our case. Multimodal treatment combinations such as systemic and intrathecal chemotherapy and radiotherapy (RT) were used. However, most of these reports did not have detailed information about toxicity. Despite the combined use of aggressive treatment modalities, patients have limited survival and very high risks of hematologic toxicity. Concurrent use of intrathecal chemotherapy and radiotherapy should be avoided due to increased risk of morbidity.

Key words: cervical cancer, leptomeningeal metastasis, radiotherapy

Oncol Clin Pract 2023; 19, 3: 184–189

Oncology in Clinical Practice
 DOI: 10.5603/OCP.2023.0010
 Copyright © 2023 Via Medica
 ISSN 2450-1654
 e-ISSN 2450-6478

Introduction

Cervical cancer is the most common gynecological cancer worldwide and a significant health problem, particularly in underdeveloped countries [1]. At the time of diagnosis, approximately 44% of patients have localized disease, 36% have a regional disease, and 16% have distant metastasis (DM) [2]. In metastatic disease, sys-

temic chemotherapy and palliative radiotherapy (RT) may be beneficial.

Uterine cervical cancer most commonly metastasizes to the lungs [3]. Leptomeningeal metastasis (LM) is an extremely rare site of DM in patients with uterine cervical cancer but is more common in lung cancer, breast cancer, and melanoma [4]. Leptomeningeal metastasis often causes neurological symptoms and is

Received: 10.02.2023 Accepted: 13.02.2023 Early publication date: 28.03.2023

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

usually observed on magnetic resonance imaging (MRI). Cytological examination of the cerebrospinal fluid (CSF) sample taken by lumbar puncture is required for accurate diagnosis unless there is a contraindication. Aggressive treatment modalities, such as a combination of intrathecal chemotherapy (ITC) and RT, are often used for treatment [5]. However, the data on the management of LM in patients with uterine cervical cancer in the literature is scarce.

In this case report, we present a case of a patient with primary uterine cervical cancer who developed LM. The primary aim of this case report and literature review is to share the treatment details for this patient and discuss and compare them with previous reports.

Case presentation

A 58-year-old female was admitted in March 2020 to the Department of Pulmonology with shortness of breath and weight loss lasting two months. Her medical, family, and psycho-social history was unremarkable. The patient underwent computed tomography (CT) of the thorax and abdomen which revealed multiple mediastinal lymph nodes (LN), uterine cervical mass, bilateral parailiac LNs, and suspicious hepatic lesions. The patient was then directed to the Gynecologic Oncology Department. In the gynecological examination under general anesthesia, a necrotic and bleeding tumoral mass of 3 cm was detected in the uterine cervix, and the right parametrium was infiltrated. A pathology study of the cervical biopsy revealed a squamous cell carcinoma (SCC) of the uterine cervix. A positron emission tomography (PET)/CT was performed for staging purposes, and extensive bone and multiple liver metastases were detected in addition to the primary mass (38 × 51 mm) in the uterine cervix and left common iliac, right internal iliac, and left external iliac LNs.

The patient was then evaluated by the Gynecologic Oncology tumor board, and a decision was made to administer systemic therapy with cisplatin (50 mg/m², total dose: 80 mg every three weeks), paclitaxel (180 mg/m², total dose: 288 mg), and bevacizumab (15 mg/kg, total dose: 870 mg). Following 6 cycles, a very good metabolic response, both in the primary and metastatic sites, was observed on the PET/CT, and definitive chemoradiotherapy was planned for the primary site and regional lymphatics after reassessment by the tumor board. A 50.4 Gy volumetric-modulated arc therapy to the uterus, cervix, proximal 1/3 of the vagina, parametrial and paravaginal sites, and bilateral common, internal and external iliac, obturator and presacral lymphatics in a 1.8-Gy fraction dose with concurrent cisplatin of 40 mg/m²/week (total dose: 60 mg) was planned and started in August 2020. However, on the MRI for brachytherapy prepa-

ration during external beam radiotherapy (EBRT), the liver metastases progressed even though the primary cervical mass and the LNs regressed, and the EBRT was stopped at 45 Gy after consideration by the tumor board. The board decided to use additional chemotherapy after completion of intracavitary brachytherapy. A total dose of 28 Gy brachytherapy was administered in 4 fractions. Following this, the patient started to receive the same chemotherapy regimen again in September 2020.

After the third cycle, the PET/CT in December 2020 revealed a complete metabolic response. Chemotherapy was stopped but bevacizumab was continued at the same dose. However, 3 months after the chemotherapy was stopped in March 2021, liver metastases progressed on the abdominal CT and weekly carboplatin (180 mg) and paclitaxel (125 mg) was initiated while bevacizumab was stopped. Following the 10th week of this regimen, the CT revealed a partial regression in the liver metastases. Two weeks after the last chemotherapy in July 2021, the patient reported pain in the right femur, and an X-ray revealed a sclerotic lesion in the distal femur. Because of the risk of pathological fracture, the patient underwent a bone curettage, cementation, and prophylactic fixation with a plaque, and a pathology examination revealed an atypical chondroid tumor of the bone. During the perioperative period, the patient could not receive any systemic therapy.

Approximately 1 month after the operation in August 2021, the patient started to report a diffuse headache and motor weakness in bilateral legs. The craniospinal MRI revealed a total of 5 cortical-meningeal metastases in the supra and infratentorial regions and extensive spinal LM, most prominent in the conus medullaris with diffuse thickening and nodular contrast enhancement in the cauda equine fibers (Fig. 1). Malignant epithelial cells were also detected in CSF cytology, which were reported as SCC metastasis of the uterine cervix. As an initial intervention, high-dose dexamethasone was administered. At this point, 2 different treatment approaches were considered by the radiation oncologists and medical oncologists; either a craniospinal irradiation (CSI) or cranial RT with intrathecal chemotherapy. Considering the possible severe toxicity of CSI, we decided to administer intrathecal methotrexate and cranial RT. Thereupon, 3 cycles of 15 mg intrathecal methotrexate were administered twice a week. Then, 30 Gy whole brain radiotherapy (WBRT) was applied in 10 fractions with concurrent 2 cycles of 15 mg of intrathecal methotrexate once a week between September 16 and 30, 2021.

Whole brain radiotherapy was well tolerated without any severe acute toxicity. A good clinical response was obtained afterward. The patient's headache completely disappeared, and her motor weakness decreased considerably. Intrathecal methotrexate was continued




Figure 1. Cranial (A) and spinal (B) leptomeningeal metastases on magnetic resonance imaging; A. Cortical-meningeal metastasis in the left cerebellar region (red circle); B. Extensive spinal leptomeningeal metastasis in the lumbar and sacral regions




Figure 2. Hematologic parameters of the patient during treatment for leptomeningeal metastases; A. Platelet count/ μL ; B. Neutrophil count/ μL ; C. Hemoglobin levels; WBRT — whole brain radiotherapy; ITC — intrathecal chemotherapy

after WBRT for 6 weeks twice a week until CSF cytology was negative on October 28, 2021. No severe neurological toxicity was observed in the remaining life period of the patient. However, the patient developed severe thrombocytopenia and neutropenia during ITC and had to be supported by platelet suspensions and granulocyte-macrophage colony-stimulating factor. The data on the number of thrombocytes and neutrophils and the level of hemoglobin are shown in Figure 2. At the end of the third month, the patient succumbed to pancytopenia, febrile neutropenia, and septic shock due to bacterial and fungal pneumonia on November 17, 2021 before the radiological response could be evaluated.

Discussion

In patients with primary uterine cervical cancer, the incidence of DM at the time of diagnosis is approximately 16%, and the most common site is the lungs [2, 3]. Metastasis to the central nervous system is unusual. The rate of parenchymal brain metastasis of uterine cervical

cancer was reported as 0.4–2.3% in the literature [6]. Leptomeningeal metastasis is even rarer, and the incidence was reported as 0.03% [7].

To the best of our knowledge, 26 cases have been reported in the literature so far, and we report the 27th patient in this case study [7–27]. We included our patient together with these 26 patients and recalculated the characteristics. Table 1 summarizes the characteristics of these 27 patients. Their median age was 47 years (range: 30–64 years), and 22% of these patients had DM at the time of diagnosis. The most common histological subtype was SCC. The latest diagnosis of LM was reported 17 years after the initial diagnosis; however, in general, LM developed within the first one to five years after initial treatment. Our patient developed LM 17 months following the initial diagnosis while under chemotherapy due to extensive metastatic disease.

Leptomeningeal metastasis of solid tumors often presents with neurologic symptoms and is usually detected by MRI. Most of the aforementioned 27 patients had neurologic symptoms consistent with lesion localization, the most common being a headache. Although

Table 1. Patient, tumor and treatment characteristics of cervical cancer patients with leptomeningeal metastasis (LM)

| Patient no. | Histology | Initial Stage | Primary treatment | Other metastasis | Time to LM diagnosis | Treatment for LM | RT field | C regimen | Survival after LM | Toxicity |
|----------------|-----------|---------------|-------------------|---|----------------------|------------------------------|------------------------|----------------------------|-------------------|--|
| #1 (7) | SCC | Localized | N/A | LN (cervical, pelvic) | 25 weeks | N/A | N/A | N/A | 17 weeks | N/A |
| #2 (7) | SCC | Localized | N/A | Brain, buttock | 190 weeks | N/A | N/A | N/A | 9 weeks | N/A |
| #3 (7) | ASC | Localized | N/A | Lung, brain | 228 weeks | N/A | N/A | N/A | 46 weeks | N/A |
| #4 (7) | AC | Localized | N/A | Cervix, endometrium | 9 weeks | N/A | N/A | N/A | 14 weeks | N/A |
| #5 (8) | NEC | Localized | Surgery SC | Breast, lung, LN (mediastinum, abdominal) | 19 months | RT | Cranial | None | 2 weeks | None |
| #6 (9) | SCC | Localized | Surgery RT | LN (pelvic), Bone | 836 weeks | RT | N/A | None | 12 weeks | N/A |
| #7 (10) | SCC | Localized | RT | LN (pelvic, PA) | 39 months | SC ITC | Craniospinal | IT-MTX, S-MeCCNU | 2 weeks | Facial paralysis, stomatitis, pancytopenia |
| #8 (11) | SCC | Metastatic | SC RT | Lung | 6 weeks | Supportive care (analgesics) | None | None | 2 weeks | None |
| #9 (12) | SCC | Localized | Surgery SC | N/A | 56 weeks | RT | Cranial | None | 4 weeks | N/A |
| #10 (13) | AC | Localized | RT | LN (PA, SCF) | 2 years | ITC RT | Cranial | IT-MTX | 1 week | N/A |
| #11 (14) | ASC | Localized | RT | Bone | 52 weeks | RT SC | Cranial | S-Cisplatin + Topotecan | 8 weeks | Septic shock, DIC |
| #12 (15) | SCC | Localized | CRT | LN (SCF) | 2 years | ITC | None | IT-MTX | 13 weeks | None |
| #13 (16) | ASC | Metastatic | SC | None | At diagnosis | ITC RT SC | Cranial | IT-MTX, S-carboplatin | 35 weeks | N/A |
| #14 (17) | SCC | Localized | Surgery CRT | Brain, lung, LN, vagina | 58 weeks | RT | Cranial | None | 3 weeks | N/A |
| #15 (18) | ASC | Localized | CRT | Liver | 31 months | RT | Craniospinal | None | 8 weeks | N/A |
| #16 (18) | NEC | Localized | CRT | Brain, bone, liver, mediastinum | 19 months | RT SC | Cranial & Focal Spinal | S-Cisplatin + Etoposide | 28 weeks | Infectious toxicity |
| #17 (19) | SCC | Metastatic | N/A | None | At diagnosis | N/A | N/A | N/A | N/A | N/A |
| #18 (20) | SCC | Localized | CRT | Bone, sciatic nerve | 10 months | N/A | N/A | N/A | N/A | N/A |
| #19 (21) | SCC | Localized | CRT | LN (PA, SCF) | 34 months | ITC RT | Cranial | IT-MTX IT-Thiotepa | 26 weeks | Cognitive deterioration |
| #20 (22) | SCC | Localized | CRT | Lung, liver, peritoneum, skin | 35 weeks | RT | Cranial & Focal Spinal | None | N/A | N/A |
| #21 (23) | NEC | N/A | N/A | None | At diagnosis | N/A | N/A | N/A | N/A | N/A |
| #22 (24) | SCC | Localized | Surgery | LN (PA, pelvic) | 13 years | RT ITC | Cranial | IT-Thiotepa | 9 weeks | None |
| #23 (25) | NEC | Metastatic | SC | Bone, LN (pelvic, PA) | 2 weeks | None | None | None | 2 weeks | None |
| #24 (26) | AC | Localized | CRT SC | LN (PA, pelvic) | 10 months | Palliative therapy | None | None | 7 weeks | None |
| #25 (27) | SCC | Metastatic | SC | LN (pelvic, PA) | At diagnosis | RT SC | Cranial | S-Paclitaxel + Carboplatin | 20 weeks | None |
| #26 (27) | SCC | Localized | Surgery RT | Lung, LN (neck, mediastinum, axilla) | 240 weeks | RT | Cranial | None | 3 weeks | None |
| #27 (our case) | SCC | Metastatic | SC CRT | Bone, liver, LN (pelvic, PA) | 68 weeks | ITC RT | Cranial | IT-MTX | 12 weeks | Pancytopenia |

AC — adenocarcinoma; ASC — adenosquamous carcinoma; C — chemotherapy; CRT — chemoradiotherapy; DIC — disseminated intravascular coagulation; IT — intrathecal; ITC — intrathecal chemotherapy; LN — lymph node; MeCCNU — semustin; MTX — methotrexate; N/A — not available; NEC — neuroendocrine carcinoma; PA — paraaortic; RT — radiotherapy; SC — systemic chemotherapy; SCC — squamous cell carcinoma; SCF — supraclavicular fossa

MRI is very valuable in detecting seeding metastases, the gold standard is the cytological examination of CSF via lumbar puncture unless there is a contraindication, such as a skin infection in the puncture site, bleeding diathesis, cardio-respiratory instability, or increased intracranial pressure.

There is no effective or successful standardized treatment in patients with LM that has a poor prognosis. Intrathecal and systemic chemotherapy with various agents and WBRT with or without spinal RT were used in the cases reported so far. In general, the role of WBRT in the treatment of LM is to provide symptom palliation due to cranial involvement and improve neurological functions. Focal spinal RT may also be considered for symptom palliation in cauda equina syndrome or symptomatic gross-nodular spinal LM. Craniospinal irradiation is mostly used in the central nervous system involvement of various hematological malignancies [28]. CSI is typically not appropriate for LM of solid tumors, particularly concurrent with ITC, due to the high risk of toxicity, short life expectancy, and low likelihood of a significant benefit. On the other hand, CSI can also be applied in LM of solid tumors with acceptable toxicity rates [29]. Chemical meningitis and leukoencephalopathy are potentially serious complications of ITC administration. Concurrent use of ITC and RT raises even higher concerns because of the risk of toxicity. In a prospective study involving 59 patients with various cancers, grade 3–5 toxicity was reported in 20% of patients who received concurrent intrathecal methotrexate and RT for LM [30]. Although severe neurological toxicity has not been reported most probably due to low survival rates in patients with cervical cancer and LM, the possibility of severe toxicity due to concurrent treatment should not be underestimated.

Unlike systemic chemotherapy, the hematological toxicity of ITC is not overemphasized. However, intrathecally administered methotrexate can enter the systemic circulation via the choroid plexus and cause systemic effects such as bone marrow suppression [31]. Kose et al. [32] reported that severe hematologic toxicity may develop after intrathecal methotrexate. The majority of case reports on cervical cancer patients with LM did not mention treatment-related toxicity or the reason for death in these patients. In Weed et al. [10] study, pancytopenia was reported in a patient during intrathecal administration of methotrexate. However, the details of this toxicity were not given. In two other studies, an infectious complication was mentioned as septic shock and pneumonia, similarly without any details [14, 18]. We think that this infectious toxicity could have been related to neutropenia. Although no radiologic response could be evaluated, our patient had improved neurologic symptoms but succumbed to treatment toxicity. We believe that in other case reports without toxicity details,

at least several patients could have developed severe hematologic toxicity and even succumbed to this toxicity.

Despite these aggressive treatment combinations, median survival after detection of LM was nine weeks (range: 1–46 weeks) in the 27 patients in the literature. Since patients with LM already have a limited life expectancy, every effort has to be taken in order not to impair their quality of life and cause more neurological toxicity while trying to regress the present neurologic symptoms. Although there are no high-quality data available to allow us to say that concurrent ITC and RT are safe, intrathecal methotrexate and WBRT were concurrently administered in our patient without severe neurological toxicity. It should be kept in mind that the time required for delayed neurological toxicities to occur may not have been reached due to the limited survival rate of the patient. In addition, the patient succumbed to pancytopenia and related sepsis less than 3 months after treatment. Although it is not clear whether the factor causing hematological toxicity was the combination of ITC and RT in our case, it seems highly possible.

Conclusion

There is insufficient high-quality evidence to guide the treatment of LM in patients with uterine cervical cancer. Despite the combined use of aggressive treatment modalities such as RT and ITC, the prognosis is quite poor. Improving the quality of life in this patient group with very low survival rates should be one of the most important goals, and the outcomes of treatment and toxicity should be well-balanced. In order not to cause severe hematologic toxicity, ITC with concurrent RT, even a focal WBRT should be avoided. Furthermore, de-intensifying the number of ITC to weekly doses can minimize hematologic toxicity.

Ethics statement

This retrospective review of patient data did not require ethical approval in accordance with local/national guidelines. Written informed consent was obtained from the patients relative for publication of the details of their medical case and any accompanying images.

Author contributions

All of the authors made significant contributions to this paper regarding the design of the manuscript, writing, and critical review in such a way that they participated sufficiently in the work to assume responsibility for the content.

Acknowledgments

None.

Conflict of interest

Authors declare no conflict of interest.

References

- Sung H, Ferlay J, Siegel RL, et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin.* 2021; 71(3): 209–249, doi: [10.3322/caac.21660](https://doi.org/10.3322/caac.21660), indexed in Pubmed: [33538338](https://pubmed.ncbi.nlm.nih.gov/33538338/).
- Siegel RL, Miller KD, Fuchs HE, et al. Cancer Statistics, 2021. *CA Cancer J Clin.* 2021; 71(1): 7–33, doi: [10.3322/caac.21654](https://doi.org/10.3322/caac.21654), indexed in Pubmed: [33433946](https://pubmed.ncbi.nlm.nih.gov/33433946/).
- Chen X, Chen L, Zhu H, et al. Risk factors and prognostic predictors for Cervical Cancer patients with lung metastasis. *J Cancer.* 2020; 11(20): 5880–5889, doi: [10.7150/jca.46258](https://doi.org/10.7150/jca.46258), indexed in Pubmed: [32922530](https://pubmed.ncbi.nlm.nih.gov/32922530/).
- Leal T, Chang JE, Mehta M, et al. Leptomeningeal Metastasis: Challenges in Diagnosis and Treatment. *Curr Cancer Ther Rev.* 2011; 7(4): 319–327, doi: [10.2174/157339411797642597](https://doi.org/10.2174/157339411797642597), indexed in Pubmed: [23251128](https://pubmed.ncbi.nlm.nih.gov/23251128/).
- Mack F, Baumert BG, Schäfer N, et al. Therapy of leptomeningeal metastasis in solid tumors. *Cancer Treat Rev.* 2016; 43: 83–91, doi: [10.1016/j.ctrv.2015.12.004](https://doi.org/10.1016/j.ctrv.2015.12.004), indexed in Pubmed: [26827696](https://pubmed.ncbi.nlm.nih.gov/26827696/).
- Divine LM, Kizer NT, Hagemann AR, et al. Clinicopathologic characteristics and survival of patients with gynecologic malignancies metastatic to the brain. *Gynecol Oncol.* 2016; 142(1): 76–82, doi: [10.1016/j.ygyno.2016.04.030](https://doi.org/10.1016/j.ygyno.2016.04.030), indexed in Pubmed: [27117923](https://pubmed.ncbi.nlm.nih.gov/27117923/).
- Yust-Katz S, Mathis S, Groves MD. Leptomeningeal metastases from genitourinary cancer: the University of Texas MD Anderson Cancer Center experience. *Med Oncol.* 2013; 30(1): 429, doi: [10.1007/s12032-012-0429-z](https://doi.org/10.1007/s12032-012-0429-z), indexed in Pubmed: [23292836](https://pubmed.ncbi.nlm.nih.gov/23292836/).
- Komiyama S, Nishio E, Torii Y, et al. A case of primary uterine cervical neuroendocrine tumor with meningeal carcinomatosis confirmed by diagnostic imaging and autopsy. *Int J Clin Oncol.* 2011; 16(5): 581–586, doi: [10.1007/s10147-010-0155-5](https://doi.org/10.1007/s10147-010-0155-5), indexed in Pubmed: [21161314](https://pubmed.ncbi.nlm.nih.gov/21161314/).
- Oike T, Ohno T, Noda SE, et al. Leptomeningeal metastasis of uterine cervical cancer 17 years after primary tumor treatment. *Clin Case Rep.* 2016; 4(1): 54–61, doi: [10.1002/ccr3.445](https://doi.org/10.1002/ccr3.445), indexed in Pubmed: [26783437](https://pubmed.ncbi.nlm.nih.gov/26783437/).
- Weed JC, Creasman WT. Meningeal carcinomatosis secondary to advanced squamous cell carcinoma of the cervix: a case report. Meningeal metastasis of advanced cervical cancer. *Gynecol Oncol.* 1975; 3(3): 201–204, doi: [10.1016/s0090-8258\(75\)80003-5](https://doi.org/10.1016/s0090-8258(75)80003-5), indexed in Pubmed: [1193431](https://pubmed.ncbi.nlm.nih.gov/1193431/).
- Weithman AM, Morrison G, Ingram EA. Meningeal metastasis of squamous-cell carcinoma of the uterine cervix: case report and review of the literature. *Diagn Cytopathol.* 1987; 3(2): 170–172, doi: [10.1002/dc.2840030217](https://doi.org/10.1002/dc.2840030217), indexed in Pubmed: [3595416](https://pubmed.ncbi.nlm.nih.gov/3595416/).
- Toyoshima M, Tsuji K, Shigeta S, et al. Leptomeningeal metastasis from gynecologic cancers diagnosed by brain MRI. *Clin Imaging.* 2017; 41: 42–47, doi: [10.1016/j.clinimag.2016.09.013](https://doi.org/10.1016/j.clinimag.2016.09.013), indexed in Pubmed: [27764719](https://pubmed.ncbi.nlm.nih.gov/27764719/).
- Aboulaifa DM, Taylor LP, Crane RD, et al. Carcinomatous meningitis complicating cervical cancer: a clinicopathologic study and literature review. *Gynecol Oncol.* 1996; 60(2): 313–318, doi: [10.1006/gyco.1996.0045](https://doi.org/10.1006/gyco.1996.0045), indexed in Pubmed: [8631558](https://pubmed.ncbi.nlm.nih.gov/8631558/).
- Lu YF, Fong VH, Wu WY, et al. Leptomeningeal metastasis of poorly differentiated uterine cervical adenosquamous carcinoma following reirradiation to metastatic vertebrae: A case report. *Medicine (Baltimore).* 2017; 96(19): e6894, doi: [10.1097/MD.0000000000006894](https://doi.org/10.1097/MD.0000000000006894), indexed in Pubmed: [28489796](https://pubmed.ncbi.nlm.nih.gov/28489796/).
- Rentink MEM, Schrijver HM, Kneppers E, et al. Carcinomatous meningitis in cancer of the uterine cervix. *J Neurooncol.* 2004; 70(1): 87–90, doi: [10.1023/b:neon.0000040836.20884.fc](https://doi.org/10.1023/b:neon.0000040836.20884.fc), indexed in Pubmed: [15527113](https://pubmed.ncbi.nlm.nih.gov/15527113/).
- Wuntkal R, Maheshwari A, Kerkar RA, et al. Carcinoma of uterine cervix primarily presenting as carcinomatous meningitis: a case report. *Aust N Z J Obstet Gynaecol.* 2004; 44(3): 268–269, doi: [10.1111/j.1479-828X.2004.00209.x](https://doi.org/10.1111/j.1479-828X.2004.00209.x), indexed in Pubmed: [15191458](https://pubmed.ncbi.nlm.nih.gov/15191458/).
- Yano H, Nagao S, Yamaguchi S. Leptomeningeal metastases arising from gynecological cancers. *Int J Clin Oncol.* 2020; 25(2): 391–395, doi: [10.1007/s10147-019-01556-1](https://doi.org/10.1007/s10147-019-01556-1), indexed in Pubmed: [31586282](https://pubmed.ncbi.nlm.nih.gov/31586282/).
- Asensio N, Luis A, Costa I, et al. Meningeal carcinomatosis and uterine carcinoma: three different clinical settings and review of the literature. *Int J Gynecol Cancer.* 2009; 19(1): 168–172, doi: [10.1111/IGC.0b013e-31819a1e1a](https://doi.org/10.1111/IGC.0b013e-31819a1e1a), indexed in Pubmed: [19258961](https://pubmed.ncbi.nlm.nih.gov/19258961/).
- Balaji R, Ramachandran K, Kumar A, et al. Pachymeningeal metastasis from squamous cell carcinoma of the uterine cervix with involvement of the optic nerve: case report and review of the literature. *Cancer Imaging.* 2007; 7(1): 138–140, doi: [10.1102/1470-7330.2007.0020](https://doi.org/10.1102/1470-7330.2007.0020), indexed in Pubmed: [17905681](https://pubmed.ncbi.nlm.nih.gov/17905681/).
- Han L, Bhan R, Johnson S, et al. Leptomeningeal metastasis in a patient with squamous cell carcinoma of the uterine cervix: Report of a case and review of the literature. *Diagn Cytopathol.* 2007; 35(10): 660–662, doi: [10.1002/dc.20705](https://doi.org/10.1002/dc.20705), indexed in Pubmed: [17854087](https://pubmed.ncbi.nlm.nih.gov/17854087/).
- Ignatius RT, Wills SM, Nadeau L, et al. Leptomeningeal carcinomatosis due to squamous cell carcinoma of the uterine cervix associated with HPV-45. *J Clin Oncol.* 2008; 26(1): 154–156, doi: [10.1200/JCO.2007.14.3180](https://doi.org/10.1200/JCO.2007.14.3180), indexed in Pubmed: [18165650](https://pubmed.ncbi.nlm.nih.gov/18165650/).
- Kastritis E, Mouloupoulos LA, Politi Ek, et al. Intramedullary spinal cord and leptomeningeal metastases in a patient with carcinoma of the uterine cervix. *Gynecol Oncol.* 2006; 102(1): 124–127, doi: [10.1016/j.ygyno.2006.01.025](https://doi.org/10.1016/j.ygyno.2006.01.025), indexed in Pubmed: [16494929](https://pubmed.ncbi.nlm.nih.gov/16494929/).
- Kumar S, Nair S, Alexander M. Carcinomatous meningitis occurring prior to a diagnosis of large cell neuroendocrine carcinoma of the uterine cervix. *J Postgrad Med.* 2004; 50(4): 311–312, indexed in Pubmed: [15623983](https://pubmed.ncbi.nlm.nih.gov/15623983/).
- Portera CC, Gottesman RF, Srodon M, et al. Optic neuropathy from metastatic squamous cell carcinoma of the cervix: an unusual CNS presentation. *Gynecol Oncol.* 2006; 102(1): 121–123, doi: [10.1016/j.ygyno.2006.01.040](https://doi.org/10.1016/j.ygyno.2006.01.040), indexed in Pubmed: [16507318](https://pubmed.ncbi.nlm.nih.gov/16507318/).
- Watanabe Y, Nakai H, Imaoka I, et al. Carcinomatous meningitis during systematic chemotherapy in a patient with advanced small-cell neuroendocrine carcinoma of the uterine cervix. *J Obstet Gynaecol Res.* 2012; 38(1): 336–339, doi: [10.1111/j.1447-0756.2011.01634.x](https://doi.org/10.1111/j.1447-0756.2011.01634.x), indexed in Pubmed: [21917069](https://pubmed.ncbi.nlm.nih.gov/21917069/).
- Yamauchi N, Sameshima H, Osato K, et al. Carcinomatous meningitis from adenocarcinoma of the uterine cervix: a case report and literature review. *J Obstet Gynaecol Res.* 2010; 36(2): 444–447, doi: [10.1111/j.1447-0756.2009.01163.x](https://doi.org/10.1111/j.1447-0756.2009.01163.x), indexed in Pubmed: [20492405](https://pubmed.ncbi.nlm.nih.gov/20492405/).
- Omori M, Ogawa T, Oyama K, et al. Leptomeningeal metastasis from cervical cancer: Report of two cases and a review of the literature. *J Obstet Gynaecol Res.* 2021; 47(8): 2782–2789, doi: [10.1111/jog.14825](https://doi.org/10.1111/jog.14825), indexed in Pubmed: [34002430](https://pubmed.ncbi.nlm.nih.gov/34002430/).
- Sanders KE, Ha CS, Cortes-Franco JE, et al. The role of craniospinal irradiation in adults with a central nervous system recurrence of leukemia. *Cancer.* 2004; 100(10): 2176–2180, doi: [10.1002/cncr.20280](https://doi.org/10.1002/cncr.20280), indexed in Pubmed: [15139061](https://pubmed.ncbi.nlm.nih.gov/15139061/).
- Devecka M, Duma MN, Wilkens JJ, et al. Craniospinal irradiation(CSI) in patients with leptomeningeal metastases: risk-benefit-profile and development of a prognostic score for decision making in the palliative setting. *BMC Cancer.* 2020; 20(1): 501, doi: [10.1186/s12885-020-06984-1](https://doi.org/10.1186/s12885-020-06984-1), indexed in Pubmed: [32487151](https://pubmed.ncbi.nlm.nih.gov/32487151/).
- Pan Z, Yang G, He H, et al. Concurrent radiotherapy and intrathecal methotrexate for treating leptomeningeal metastasis from solid tumors with adverse prognostic factors: A prospective and single-arm study. *Int J Cancer.* 2016; 139(8): 1864–1872, doi: [10.1002/ijc.30214](https://doi.org/10.1002/ijc.30214), indexed in Pubmed: [27243238](https://pubmed.ncbi.nlm.nih.gov/27243238/).
- Rubin R, Owens E, Rall D. Transport of methotrexate by the choroid plexus. *Cancer Res.* 1968; 28(4): 689–694, indexed in Pubmed: [5649058](https://pubmed.ncbi.nlm.nih.gov/5649058/).
- Kose F, Abali H, Sezer A, et al. Little dose, huge toxicity: profound hematological toxicity of intrathecal methotrexate. *Leuk Lymphoma.* 2009; 50(2): 282–283, doi: [10.1080/10428190802603169](https://doi.org/10.1080/10428190802603169), indexed in Pubmed: [19197737](https://pubmed.ncbi.nlm.nih.gov/19197737/).