

Zuzanna Gurzyńska¹ , Krzysztof Sobczak² , Milena Aneta Lachowicz³ 

¹Student Scientific Circle of Medical Communication, Medical University of Gdansk, Gdańsk, Poland

²Department of Sociology of Medicine and Social Pathology, Faculty of Health Sciences, Medical University of Gdansk, Gdańsk, Poland

³Department of Oncology and Radiotherapy, Medical University of Gdansk, Gdańsk, Poland

Exploring the readiness of hospice and oncology unit staff to offer spiritual support to patients: preliminary findings

Abstract

Introduction: Patients' spiritual needs are often marginalized by medical staff, who identify them with religious practice, faith, and God. Illness and related human suffering are not just physical ailments alone. The loss of health entails several changes in a patient's life and requires reorganization of the family, professional, social, and spiritual aspects of it. The purpose of this article is to analyze the medical personnel's sense of readiness to provide spiritual care to patients of oncology units and hospice facilities.

Methods: The study was carried out with the use of the Spiritual Supporter Scale whose psychometric values were determined on a high level of reliability with Cronbach's $\alpha = 0.88$.

Results: The results of the Spiritual Supporter Scale showed that oncology professionals got significantly lower scores [median (Me) = 5; mean (M) = 5.11; standard deviation (SD) = 1.89] in the overall scale score than those who work in hospices (Me = 7; M = 6.9; SD = 1.5). The analysis also showed that oncology unit employees (54%), declared that providing spiritual care to patients is an integral part of their work ($p < 0.02$) significantly less often than those employed in hospices (88%). They were also less likely to declare (77%) that spiritual support is necessary in their workplace than persons who provide care in hospices (95%; $p < 0.01$).

Conclusions: The study showed the differences in the sense of preparedness as well as competencies to provide spiritual care among medical workers in hospices and oncology units.

Palliat Med Pract

Keywords: spiritual care, hospice care, oncology patients, palliative care, psycho-oncology

Introduction

In the 1990s, Hiatt [1] proposed to extend the psychosomatic model of health and sickness, which was standard at the time, to include a spiritual dimension. This stipulation resulted

from many research observations, which pointed to the spiritual dimension being an important and inseparable element of behavior in health and sickness [2]. Even though spirituality in healthcare has already been defined in multiple ways, there is an overall consensus in the reference sources to usually understand

Address for correspondence:

Zuzanna Gurzyńska

Student Scientific Circle of Medical Communication, Medical University of Gdansk, Tuwima 15, 80–210 Gdańsk, Poland

e-mail: zuz.gurzynska@gumed.edu.pl



Palliative Medicine in Practice

Copyright © 2024 Via Medica, ISSN 2545–0425, e-ISSN 2545–1359

DOI: 10.5603/pmp.98092

Received: 3.11.2023 Accepted: 29.02.2024 Early publication date: 1.03.2024

This article is available in open access under Creative Commons Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

it in a broader context than just religious experience because it is an existential construct with a more fundamental meaning [3]. One of the suggestions for defining spirituality is framing it as “how people understand and live their lives given their ultimate meaning and value” [4]. Current scientific theories highlight, more and more boldly, the categories of spirituality as one of the elements of the holistic concepts of humans. Some ideas include not only mental and emotional intelligence but also a spiritual one as a foundation for all internal processes which allow people to discover things and give them meaning which is so important for understanding and conscious experiencing of existential dilemmas [5].

According to the phenomenological concept of Schütz [6], illness changes one’s perception of the world exponentially. From this perspective, it can be assumed that in the areas connected with receiving bad news, fear of death is a fundamental experience. The need to control “my world” is one of the elementary existential needs. The experience of an illness becomes a prism that, to various extents or sometimes fully, deconstructs previous meanings, experiences, and social relationships (and sometimes even metaphysical experiences) or creates new ones in the light of new values. Therefore, besides the physical, psychological, and sociological aspects, sicknesses involve a spiritual aspect. A plethora of research provides a lot of interesting information to confirm this view. They point to the fact that experiencing an illness can substantially change the existential perception in the area of the meaning of life as well as values [7, 8]. In this context, the spiritual dimension turns out to be an important element. It correlates with strategies of coping which, especially in the perspective of a life-ending, become an area that determines life quality [9]. Spiritual aspects of existence stay strictly connected with the emotional sphere as well as cognitive functioning. In stressful situations, persons with a more coherent spiritual structure have access to a wider variety of coping strategies, which relates to better psychological and physical health [10]. A positive correlation between spiritual sensitivity and a subjective sense of well-being was also pointed out [11]. Research revealed that a higher spiritual sensitivity is associated with a better adaptation to a chronic, somatic disease, optimistic predictions regarding its further course, and a sense of a bigger influence on that course [12]. Moreover, a study conducted among oncology patients also showed a negative correlation between the level of depression or anxiety and the level of spiritual sensitivity [13].

Analyses conducted over the years have revealed that sensitivity to spiritual categories can be a sig-

nificant factor in the realm of therapeutic influence, particularly in contexts related to facing illnesses with pessimistic prognoses [14]. For that reason, Anglo-Saxon countries started including programs in the structure of medical education, which develop competencies of health professionals to increase their sensitivity to spiritual aspects of diseases as well as patients’ needs in this regard [15]. As persons who support oncology patients and their families, like nurses, doctors, psychologists, physiotherapists, or capelans, play a special role, formal guidelines that provide recommendations concerning clinical practice were developed [16].

Though educational interventions to include the subject of spirituality in medical education were successfully implemented in Anglo-Saxon countries, this has not been so in Poland. According to available information, the only spiritual care course with particular emphasis on incurable diseases for medical students in Poland began at Collegium Medicum in Bydgoszcz of the Nicolaus Copernicus University in Toruń in 2018 [17]. It continues to be taught as a mandatory subject to this day. Additionally, the Polish Association for Spiritual Care in Medicine, which was established in 2015, remains active. Despite areas regarding the spiritual care of patients in Poland being noticed in the research space [18–21], there remains a serious deficit of analyses concerning this aspect of intervention. That’s why this research aimed to try to analyze the medical staff’s sense of being prepared to provide spiritual care to patients. It is a significant problem because both medical staff and the patients notice the need for this type of intervention and point to its deficit [22, 23].

This study aimed to analyze the medical staff’s sense of being prepared to provide spiritual care to hospice and oncology patients. The authors wanted to find out if health professionals who provide active and direct care to hospice and oncology patients differ in terms of their readiness to provide spiritual care to their patients and their sense of this readiness as well as the self-assessment of their competencies in this area.

Methods

Study design

The cross-sectional study was conducted using a survey questionnaire technique. The research was executed with the help of a standardized Spiritual Supporter Scale, for which the psychometric values were defined at a high level of reliability with Cronbach’s $\alpha = 0.88$ [24]. The questionnaire contains 31 closed questions based on multiple-choice answers in the

form of the five-point Likert scale. The results in the scale's construction were structured into five subscales: Attitude to prayer; Beliefs regarding spirituality; Spirituality in relation to one's own suffering and the suffering of others; Sensitivity to the suffering of others; Attitude to the community as well as the general score. The raw results were converted to sten scores accordingly. Scores within sten 1–2 were defined as very low, 3–4 as low, 5–6 as medium, 7–8 as high, and 9–10 as very high [24].

The research instrument additionally included six additional statements, which participants addressed using a five-point Likert scale. These statements pertained to respondents' opinions regarding their readiness to provide patients with preparation for spiritual support, the importance of spiritual care in caring for the patient and their family, the role of spiritual support as a professional task, and readiness to develop competencies in spiritual care. Multiple-choice responses of "strongly disagree" and "disagree", as well as "strongly agree" and "agree", were aggregated during the scoring analysis. The categorical variables in this study included: gender, profession, workplace, and religious affiliation.

Setting

The field stage of the study was conducted between the 12th of December 2022 and the 26th of April 2023 at the University Clinical Centre in Gdańsk, Polish Red Cross Marine Hospital in Gdynia, Saint Lawrence Hospice in Gdynia and Father Pio Hospice in Puck. After obtaining the necessary consent of the institutions' administrators and chiefs of the units, the research questionnaires were distributed. Every physician, psychologist, and nurse had an opportunity to take part in the study.

Participant identification

The conditions for inclusion in the study were: employment as a physician, nurse, or psychologist and completing the whole questionnaire. 180 questionnaires were distributed. Saturation for this study was 43%. Ultimately, 75 people were included in the analysis. Four questionnaires were excluded: 3 of them were not filled out properly. One person declared being employed in both a hospice and an oncology ward, which did not allow to assign them to any group.

Ethics approval and consent to participate

The Independent Bioethics Committee issued a decision (KB/35/2024) to exempt this study from the requirement of Bioethics Committee approval. Before beginning the study, formal permissions for conducting the study were obtained from the heads

of the departments and hospices where the research was performed. No sensitive data was being collected or processed during the study. Each respondent gave consent to take part in the study.

Data analysis

The obtained data was subjected to statistical analysis with the use of Statistica v.13.3. The Shapiro–Wilk test was used to analyze the distribution. Student's t-test was also used, as well as the U Mann–Whitney test for non-parametric variables, which enabled defining differences between groups and analyzing their statistical significance. To assess differences between groups in the independent questions, Pearson's chi-square test was used. The r-Pearson's linear correlation coefficient was also calculated to verify the existence of a correlation between age and the scale scores as well as between seniority and the scale scores. During the analysis, a statistical significance of $p < 0.05$ was assumed.

Results

Seventy-nine people took part in the study. Seventy-five people were included in the analysis, of which forty declared being hospice workers and thirty-five — oncology unit workers. There were 70 women and 5 men, including 53 nurses, 13 physicians, and 9 psychologists between the ages of 22 and 63 ($M = 43.64$). Within that, 64 people declared a denomination, 5 stated they were atheists and 6 agnostics. The respondents' seniority varied from 1 year to 42 years ($M = 19.64$).

The score analysis of the Spiritual Supporter Scale [24] revealed that significant differences were observed between the two groups in the overall score ($p < 0.001$) as well as in 4 out of 5 subscales: Spirituality in relation to one's own suffering and the suffering of others ($p < 0.00$); Attitude to prayer ($p < 0.00$); Beliefs regarding spirituality ($p < 0.05$); Sensitivity to the suffering of others ($p < 0.02$). On average, in the overall score, respondents who worked in oncology wards got lower scores [median (Me) = 5; mean (M) = 5.11; standard deviation (SD) = 1.89] compared to those who worked in hospices ($Me = 7$; $M = 6.9$; $SD = 1.50$). Additionally, in each of the 4 subscales in which there were statistically significant differences, oncology unit employees had lower average scores than those who provided care in hospices (Table 1). The score analysis did not show a linear correlation between the respondents' seniority or age and their scores on the Spiritual Supporter Scale (Table 2).

Both groups were given questions concerning their opinion on providing spiritual care with answers

Table 1. Analysis of inter-group differences in total score and individual scales of the Spiritual Supporter Scale

Scale	Researched group										p-value
	Hospice workers (n = 40)					Oncology unit workers (n = 35)					
	Me	M	SD	Min	Max	Me	M	SD	Min	Max	
Spirituality in relation to one's own suffering and the suffering of others	7	7.12	1.65	5	10	5	5.46	2.05	1	10	0.00*
Attitude to prayer	7	7.25	1.53	5	10	5	5.00	1.94	1	9	0.00*
Beliefs regarding spirituality	6	5.27	1.38	3	7	5	4.54	1.58	1	7	0.047*
Attitude to community	6	6.00	2.01	2	9	6	6.06	1.81	3	9	0.91
Sensitivity to the suffering of others	6	6.47	1.38	2	9	6	5.66	1.33	3	9	0.01*
Total score	7	6.9	1.50	5	10	5	5.11	1.89	1	10	0.00*

*Scores significant with $p < 0.05$; M — mean; Max — maximum; Me — median; Min — minimum; SD — standard deviation

Table 2. Results of correlation analysis between seniority and age of respondents and individual scales and total score of the Spiritual Supporter Scale

Scale	Seniority	Age of respondents
	r-value	
Spirituality in relation to one's own suffering and the suffering of others	0.07	0.07
Attitude to prayer	0.29	0.35
Beliefs regarding spirituality	-0.00	0.05
Attitude to community	-0.15	-0.1
Sensitivity to the suffering of others	0.15	0.15
Total score	0.15	0.2

in the form of the five-point Likert scale. The answers strongly disagree and disagree as well as strongly agree and agree were grouped into two categories. Analysis showed that persons employed in oncology units (54%) declared that providing spiritual care to patients is an inseparable part of their job ($p < 0.02$) significantly less often than hospice staff (88%). Compared to the respondents who worked in hospices (95%; $p < 0.01$), they also were less likely (77%) to say that this type of support is necessary in their workplace. Groups of hospice (50%) and oncology unit (71%) healthcare professionals expressed a similar level of disagreement with the claim that studies prepared them for providing spiritual care to patients ($p = 0.21$). However, oncology employees

(44%) declared a lack of desire ($p < 0.03$) to develop competencies in this area (Table 3) significantly more often than hospice staff (13%).

Discussion

In this study, results obtained with the help of the Spiritual Supporter Scale point to a lower sense of being prepared for providing spiritual support among oncology unit healthcare professionals in comparison to persons who work in hospices. It relates to both understanding one's own spirituality as well as attitude towards it to a person in need of support and care [24]. What is more, oncology unit medical staff feel that providing spiritual care to patients and their families lies in their area of responsibility much less often. It might be that this view stems from slightly different foundations underlying the therapeutic impact.

It was decided to compare the results of the study with the ones obtained during the standardization of the Spiritual Supporter Scale [24]. The comparison revealed that the average score of the subscales: Spirituality in relation to one's own suffering and the suffering of others; Attitude to prayer; Sensitivity to the suffering of others as well as the scale's overall score coincided with the average score of the study's oncology staff group, which placed it at the average level on the sten scale. However, in the present study, the hospice healthcare workers group got high scores on the sten scale in all the subscales above. It might result from the fact that students of medicine, other medical fields as well as non-medical ones, and teachers took part in the standardization of the tool. As students were the majority of the study's population, they might not have gained extensive experience regarding

Table 3. Distribution of responses regarding statements about the preparedness and need for spiritual care for patients

Statements	Strongly disagree and disagree		Strongly agree and agree		p-value
	Hospice workers	Oncology unit workers	Hospice workers	Oncology unit workers	
My studies prepared me to provide spiritual care to patients	50%	71%	33%	17%	0.21
Providing spiritual care to patients is essential in my place of work	3%	14%	95%	77%	0.01*
Providing spiritual care to the patient's family is essential in my workplace	3%	17%	93%	63%	0.00*
Providing spiritual care to patients is an integral part of my work	5%	29%	88%	54%	0.02*
Providing spiritual care to the patient's family is an integral part of my work	13%	33%	45%	41%	0.02*
I would like to develop my competencies in spiritual care	13%	44%	45%	37%	0.02*

*Scores significant with $p < 0.05$

spiritual care in the medical field. Those types of competencies are also gained through incidental learning, especially in the area of end-of-life care. In the scale of Beliefs Regarding Spirituality, hospice staff got comparable scores to the study group in the standardization, whereas oncology employees obtained significantly lower ones. In the scale of Attitude to Community, both groups in the study got a similar score to the population analyzed in the questionnaire's standardization. As this portion of the results was summed up, it had to be noticed that the level of spiritual sensitivity of the analyzed healthcare professionals turned out both low and lower than expected.

Most of the hospice (95%) and oncology unit (77%) employees who participated in the study admitted that providing spiritual care to patients they take care of is necessary in their workplace. Balboni et al. [22] got similar results in their study. Over 80% of the oncology unit doctors and nurses who were asked, declared that spiritual care should be provided to patients at least sporadically. There were interesting results in the questions concerning the sense of being prepared for providing spiritual care to patients and one's own need to expand competencies in this area. Firstly, it should be noted that hospice employees recognized that they were better prepared to provide spiritual care within their formal education than oncology unit employees. Despite that, healthcare professionals who worked in oncology units declared the desire to develop their skills in providing spiritual care significantly less often (Table 3). It might be that this

stance, as suggested by Peteet and Balboni [25] in their research, results from the assumption that medical workers do not acknowledge providing spiritual care as an element of their professional duty. The authors think that it can be assumed, with some probability, that this stance results partly from understanding spiritual support as interference in the patient's religious sphere.

Although this study delivers important observations, the authors realize it also has some objective limitations. In the presented preliminary research, there was an overrepresentation of women, who constituted 93% of the respondents. What is more, the overrepresentation of nurses (71%) in comparison to other medical professionals included in the study did not allow us to obtain a correlation between the respondents' profession with the questionnaire scores. The research tool used in the study, in the subscale Attitude to Prayer may, to some extent, lead respondents to identify spirituality with an element of religious practice. Associating spirituality with religiousness also had a significant influence on the saturation obtained in the study. A significant churn rate was observed — in the group of oncology unit employees who withdrew from the study during the phase of questionnaire distribution. Many potential respondents refused to their part in the study due to their atheistic worldview as well as the reluctance to reveal their value system or their attitude towards God and religion. It seems to us that this aspect, although not directly analyzed in the study, is an

interesting direction that outlines a possibility to extend the analyses to the understanding of spirituality by healthcare workers.

Conclusions

Despite the growing amount of research, pointing to spiritual care as an important element of medical intervention, especially in areas concerning pessimistic prognoses and end-of-life, there are significant deficits in the domain of a sense of being prepared and a desire to provide spiritual care to patients among hospice and oncology unit medical workers.

Article information and declarations

The authors declare that they have not used or cooperated with Chat Generative Pre-trained Transformer (ChatGPT) technology or other artificial intelligence (AI) models in conducting the study, analyzing the data, and preparing the manuscript.

Acknowledgments

None.

Data availability statement

The data that support the findings of this study are available on request from the corresponding author.

Ethics statement

The Independent Bioethics Committee issued a decision (KB/35/2024) to exempt this study from the requirement of review by bioethical committees.

Author contributions

ZG was responsible for preparing the study, completion of the field stage, statistical analysis of the results as well as preparing the first manuscript draft. KS was responsible for project oversight and result analysis. He also prepared the last manuscript version. ML took part in the field research and was responsible for proofreading and editing the article.

Conflict of interest

The authors declare that they have no competing interests.

Funding

This research received no specific grant from any funding agency.

Supplementary material

None.

References

- Hiatt JF. Spirituality, medicine, and healing. *South Med J*. 1986; 79(6): 736–743, doi: [10.1097/00007611-198606000-00022](https://doi.org/10.1097/00007611-198606000-00022), indexed in Pubmed: [3715539](https://pubmed.ncbi.nlm.nih.gov/3715539/).
- Balboni TA, VanderWeele TJ, Doan-Soares SD, et al. Spirituality in serious illness and health. *JAMA*. 2022; 328(2): 184–197, doi: [10.1001/jama.2022.11086](https://doi.org/10.1001/jama.2022.11086), indexed in Pubmed: [35819420](https://pubmed.ncbi.nlm.nih.gov/35819420/).
- de Brito Sena MA, Damiano RF, Lucchetti G, et al. Defining spirituality in healthcare: a systematic review and conceptual framework. *Front Psychol*. 2021; 18(12): 756080, doi: [10.3389/fpsyg.2021.756080](https://doi.org/10.3389/fpsyg.2021.756080), indexed in Pubmed: [34867654](https://pubmed.ncbi.nlm.nih.gov/34867654/).
- Muldoon M, King N. Spirituality, health care, and bioethics. *J Relig Health*. 1995; 34(4): 329–349, doi: [10.1007/BF02248742](https://doi.org/10.1007/BF02248742), indexed in Pubmed: [11660133](https://pubmed.ncbi.nlm.nih.gov/11660133/).
- Michałowski S. Spiritual intelligence and its development. Unifying thinking and “psychagogical” meditation. *Prace Naukowe Akademii im. Jana Długosza w Częstochowie. Pedagogika*. 2014; 23: 49–75, doi: [10.16926/p.2014.23.03](https://doi.org/10.16926/p.2014.23.03).
- Schütz A. *Phenomenology and the social sciences*. Polish. PIW, Warszawa 1989.
- Brady MJ, Peterman AH, Fitchett G, et al. A case for including spirituality in quality of life measurement in oncology. *Psychooncology*. 1999; 8(5): 417–428, doi: [10.1002/\(sici\)1099-1611\(199909/10\)8:5<417::aid-pon398>3.0.co;2-4](https://doi.org/10.1002/(sici)1099-1611(199909/10)8:5<417::aid-pon398>3.0.co;2-4), indexed in Pubmed: [10559801](https://pubmed.ncbi.nlm.nih.gov/10559801/).
- Ferrell BR, Dow KH, Leigh S, et al. Quality of life in long-term cancer survivors. *Oncol Nurs Forum*. 1995; 22(6): 915–922, indexed in Pubmed: [7567610](https://pubmed.ncbi.nlm.nih.gov/7567610/).
- McClain CS, Rosenfeld B, Breitbart W. Effect of spiritual well-being on end-of-life despair in terminally-ill cancer patients. *Lancet*. 2003; 361(9361): 1603–1607, doi: [10.1016/S0140-6736\(03\)13310-7](https://doi.org/10.1016/S0140-6736(03)13310-7), indexed in Pubmed: [12747880](https://pubmed.ncbi.nlm.nih.gov/12747880/).
- Pargament KI. *The psychology of religion and coping: theory, research, practice*. The Guilford Press, New York 2001.
- Bożek A, Nowak PF, Blukacz M. The relationship between spirituality, health-related behavior, and psychological well-being. *Front Psychol*. 2020; 14(11): 1997, doi: [10.3389/fpsyg.2020.01997](https://doi.org/10.3389/fpsyg.2020.01997), indexed in Pubmed: [32922340](https://pubmed.ncbi.nlm.nih.gov/32922340/).
- Heszen-Niejodek I, Gruszczyńska E. Spirituality as a human dimension, its importance in health psychology, and its measurement. *Przeegl Psychol*. 2004; 47(1): 15–31.
- Turke KC, Canonaco JS, Artioli T, et al. Depression, anxiety and spirituality in oncology patients. *Rev Assoc Med Bras (1992)*. 2020; 66(7): 960–965, doi: [10.1590/1806-9282.66.7.960](https://doi.org/10.1590/1806-9282.66.7.960), indexed in Pubmed: [32844942](https://pubmed.ncbi.nlm.nih.gov/32844942/).
- Balboni T, Balboni M, Paulk ME, et al. Support of cancer patients’ spiritual needs and associations with medical care costs at the end of life. *Cancer*. 2011; 117(23): 5383–5391, doi: [10.1002/cncr.26221](https://doi.org/10.1002/cncr.26221), indexed in Pubmed: [21563177](https://pubmed.ncbi.nlm.nih.gov/21563177/).
- Osório IH, Gonçalves LM, Pozzobon PM, et al. Effect of an educational intervention in “spirituality and health” on knowledge, attitudes, and skills of students in health-related areas: A controlled randomized trial. *Med Teach*. 2017; 39(10): 1057–1064, doi: [10.1080/0142159X.2017.1337878](https://doi.org/10.1080/0142159X.2017.1337878), indexed in Pubmed: [28641027](https://pubmed.ncbi.nlm.nih.gov/28641027/).
- Ferrell BR, Baird P. Deriving meaning and faith in caregiving. *Semin Oncol Nurs*. 2012; 28(4): 256–261, doi: [10.1016/j.soncn.2012.09.008](https://doi.org/10.1016/j.soncn.2012.09.008), indexed in Pubmed: [23107183](https://pubmed.ncbi.nlm.nih.gov/23107183/).
- Fopka-Kowalczyk M, Groves R, Larkin P, et al. A training programme for medical students in providing spiritual care to people with advanced diseases and their loved ones: A case study from the Collegium Medicum in Bydgoszcz, Nicolaus Copernicus University in Toruń, Poland. *Front Cardiovasc*

- Med. 2022; 29(9): 909959, doi: [10.3389/fcvm.2022.909959](https://doi.org/10.3389/fcvm.2022.909959), indexed in Pubmed: [36247435](https://pubmed.ncbi.nlm.nih.gov/36247435/).
18. Klimasiński M. Spiritual care in medicine: forgiveness. *Sztuka Leczenia*. 2018; 33(2): 33–40, doi: [10.4467/18982026szl](https://doi.org/10.4467/18982026szl).
 19. Klimasiński MW. Spiritual care in the intensive care unit. *Anaesthesiol Intensive Ther*. 2021; 53(4): 350–357, doi: [10.5114/ait.2021.109920](https://doi.org/10.5114/ait.2021.109920), indexed in Pubmed: [34714016](https://pubmed.ncbi.nlm.nih.gov/34714016/).
 20. Klimasiński MW, Theda J, Cofta S. Spiritual care in medicine: spiritual perception of illness, spiritual coping with suffering — a quantitative survey study on the Polish population of chronically ill adults. *Sztuka Leczenia*. 2020; 35(1): 9–18, doi: [10.34938/h1k9-t164](https://doi.org/10.34938/h1k9-t164).
 21. Zurzycka PZ, Czyżowicz K, Radzik T. Spiritual care is a duty in the nursing practice — an outline of the problem. *Problemy Pielęgniarstwa*. 2017; 25(3): 210–214.
 22. Balboni MJ, Sullivan A, Amobi A, et al. Why is spiritual care infrequent at the end of life? Spiritual care perceptions among patients, nurses, and physicians and the role of training. *J Clin Oncol*. 2013; 31(4): 461–467, doi: [10.1200/JCO.2012.44.6443](https://doi.org/10.1200/JCO.2012.44.6443), indexed in Pubmed: [23248245](https://pubmed.ncbi.nlm.nih.gov/23248245/).
 23. Chow HH, Chew QH, Sim K. Spirituality and religion in residents and inter-relationships with clinical practice and residency training: a scoping review. *BMJ Open*. 2021; 11(5): e044321, doi: [10.1136/bmjopen-2020-044321](https://doi.org/10.1136/bmjopen-2020-044321), indexed in Pubmed: [34049909](https://pubmed.ncbi.nlm.nih.gov/34049909/).
 24. Fopka-Kowalczyk M, Best M, Krajnik M. The spiritual supporter scale as a new tool for assessing spiritual care competencies in professionals: design, validation, and psychometric evaluation. *J Relig Health*. 2023; 62(3): 2081–2111, doi: [10.1007/s10943-022-01608-3](https://doi.org/10.1007/s10943-022-01608-3), indexed in Pubmed: [35881265](https://pubmed.ncbi.nlm.nih.gov/35881265/).
 25. Peteet JR, Balboni MJ. Spirituality and religion in oncology. *CA Cancer J Clin*. 2013; 63(4): 280–289, doi: [10.3322/caac.21187](https://doi.org/10.3322/caac.21187), indexed in Pubmed: [23625473](https://pubmed.ncbi.nlm.nih.gov/23625473/).