

Spiritual care competence and caring abilities among Polish nurses: a correlation descriptive study

Abstract

Background: Spiritual care is an obligatory aspect of nursing care for a patient at the end of life and plays an important role in providing quality nursing care. This study aimed to describe the level of spiritual care competence and caring abilities of Polish nurses and to examine the relationship between them.

Participants and methods: In the study based on the Caring Ability Inventory (CAI) and the Spiritual Care Competence Scale (SCCS) questionnaires descriptive, correlational, cross-sectional design was adopted. The study involved 451 clinical nurses.

Results: Respondents' overall score of SCCS was high (median = 101.22; mean = 103.00; SD = 17.14) and low in the overall score of CAI (median = 185.44; mean = 185.00; SD = 21.05). The respondents who believe that nurses should assess the patient's spiritual needs obtained statistically higher scores in all subscales and the overall score of SCCS and CAI. The total score of spiritual care competence was also positively correlated with the level of caring abilities ($p < 0.01$). The caring abilities and competencies to provide spiritual care of Polish nurses correlate, among others, with their age, professional experience, level of education and job satisfaction.

Conclusions: The implementation by academic teachers and nursing managers of strategies promoting patient-centred nursing care and humanistic values will contribute to strengthening the caring abilities of nurses and the skills to provide spiritual care to patients, especially in the most basic human experience of dying.

Palliat Med Pract 2023; 17, 4: 216–224

Keywords: caring, nurses, spiritual care, competence, caring abilities, palliative care

Introduction

For nursing care to be effective and efficient, patients' physiological, mental, socio-cultural, developmental, and spiritual needs must be met.

However, compared to other matters of nursing care, spiritual care has received less attention. The biological model based on the positivist damage paradigm predominated in medicine and science during the nineteenth and twentieth centuries when

Address for correspondence:

Michał Machul

Department of Holistic Care and Nursing Management, Medical University of Lublin, Staszica 4/6, 20–081 Lublin, Poland

e-mail: michal.machul@umlub.pl



Palliative Medicine in Practice 2023; 17, 4, 216–224

Copyright © 2023 Via Medica, ISSN 2545–0425, e-ISSN 2545–1359

DOI: 10.5603/PMPI.a2023.0022

Received: 21.05.2023 Accepted: 26.06.2023 Early publication date: 29.06.2023

This article is available in open access under Creative Commons Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

spiritual care started to be disregarded [1]. Spiritual care is an obligatory aspect of nursing care, which should be provided by personnel with high spiritual competence [2]. Palliative and end-of-life guidelines [3], as well as professional standards and nursing theories, emphasize and recognize the importance of spirituality and spiritual care in patient care [4, 5]. By implementing all aspects of care, nurses provide services at a high level and can meet all the needs of their patients [6]. Nurses must be educated in all aspects of healthcare to provide effective and efficient patient care. Giving patients access to spiritual care aids their recovery, even though most health services pay little attention to patients' spiritual needs [7]. As a result, in order to successfully and securely support patients' health, nurses must be knowledgeable in spiritual care.

The quality of nursing care is another important aspect of healthcare delivery. Healthcare delivery must consider whether the patient is receiving the desired level of care and satisfaction [8]. Previous studies have shown that nurses with high spiritual competencies may provide care of higher quality [9]. Rationing or not providing spiritual care has a negative impact on the recovery of patients, often contributing to prolonged hospitalization and convalescence [10]. In addition, not taking into account spiritual care in patient care by nursing staff also contributes to the occurrence of social isolation and mental suffering among patients especially among terminally ill and those approaching the end of life [11, 12]. In relation to palliative care, Jo has shown that the empathy and caring abilities of nurses are significant predictors of terminal care efficiency [13].

Despite the important role of spiritual care in nursing practice, limited studies have been conducted to investigate the relationship between nurses' spiritual care competence and the quality of care reflected in the nurses' caring abilities, so far this relationship has not been explored in Poland. Identification of elements that affect nurses' caring abilities is of key importance for improving the quality of patient care, especially in the context of previous studies that showed that Polish nurses have low caring abilities. This study also appears useful because a better understanding of the relationship between spiritual care competence and caring abilities will benefit nursing education and practice as well as improve the quality of nursing care and will answer the question of what factors determine nurses' caring abilities in Poland. This study was conducted to determine the level of spiritual care competence and caring abilities of Polish nurses and to examine the relationship between them.

Participants and methods

Study design

This was a descriptive, correlational, cross-sectional study conducted in 2022 in Poland. The research conformed to the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: Guidelines for Reporting Observational Studies.

Participants and setting

The Raosoft Sample Size Calculator was used to determine the sample size [14]. There were approximately 232 387 nurses in Poland at the end of 2021 [15]. For a confidence level of 0.95, a margin of error of 0.05, and a response distribution of 0.50, a sample size of 384 was required. Considering the loss of 10% of the sample, a sample size of 422 was required. The participants were selected based on the following criteria: (a) being an active registered nurse, (b) minimum 2 years of working experience, (c) practising in the clinical settings, (d) access to the Internet, (e) speaking and understanding the Polish language, and (f) consent to participate in the study. Staff nurses who had no direct contact with patients prior to this study were excluded.

Measures

Three instruments were used in the study:

- 1. Spiritual Care Competence Scale.** Van Leeuwen et al. [16] developed the Spiritual Care Competence Scale (SCCS), a self-reporting scale to measure nurses' spiritual care competence. It includes the following six subscales: Assessment and implementation of spiritual care; Professionalisation and improvement of the quality of spiritual care; Personal support and patient counselling; Referral to professionals; Attitude towards the patient's spirituality; and Communication. The questionnaire contains 27 items, all of which are scored on a 5-point scale from completely disagree to fully agree, scoring from 1 to 5. The minimum and maximum possible values are 27 and 135, respectively. A score lower than 64 indicates low spiritual competence, a score of 64–98 indicates average spiritual competence and a score of 99 and above implies high spiritual competence [16]. Machul and Dobrowolska revised the Polish version of the SCCS (referred to as SCCS-PL), composed of 27 items displaying a five-factor structure. The Polish version of the scale received 0.95 Cronbach alpha [17]. The Polish version of the SCCS-PL can be found in Supplementary File 1.
- 2. Caring Ability Inventory.** The Caring Ability Inventory (CAI) is a self-reporting instrument including 37 items with a 5-point Likert scale. It was constructed by Nkongho [18] and is intended to

measure the degree of a person's caring ability. The CAI contains three dimensions: Knowledge (14 items), Courage (13 items), and Patience (10 items). The answers are organized on a Likert-type scale from 1 (I strongly disagree) to 7 (I strongly agree), with scores ranging from 37 to 259. The reliability of the CAI was measured by Cronbach's alpha which ranged from 0.71 to 0.84. The higher the score, the higher the level of care ability [18]. The Polish version of the CAI validated in 2022 by Machul and Dobrowolska (referred to as CAI-PL) consists of 36 items and the possible scores to be obtained range from 36 to 252 with Cronbach's alpha 0.715. The Polish version of the CAI-PL can be found in Supplementary File 2.

3. **A short self-made questionnaire.** A short self-made questionnaire to collect socio-demographic characteristics such as sex, age, marital status, degree of nursing education, postgraduate qualification, place of work, years of nursing experience, faith, job satisfaction, and an individual opinion on the initial assessment of spiritual needs.

Data collection

The study was conducted between January and the end of July 2022. Given that the research was conducted during the SARS-CoV-2 pandemic, an online questionnaire was prepared. Convenience sampling was conducted. The respondents were enrolled through invitations posted on blogs, discussion forums and social networking sites devoted to health sciences and nursing. The questionnaire was delivered to 1197 respondents, with 451 questionnaires (38%) correctly filled in and returned. To collect data, an online questionnaire was used, which was formatted using Survio software. The study was conducted with the international standard for information security requirements and is fully compliant with the processing of personal details pursuant to the German personal data protection act. Each response was protected by an international safety certificate with extended validation (Organization Validation SSL Certificate). The online form contained information about the purpose and significance of the study and informed the prospective participants of the fact that participation was completely voluntary.

Ethical considerations

The standards stipulated in the Declaration of Helsinki were followed during each step of the study. Participation was voluntary, anonymous and confidential. All the participants were informed about the study purpose and procedures on the first page of the online survey and were assured that all data would

be confidential. In addition, they were allowed to withdraw from the study at any time. The data stored and managed in the questionnaire available via the Survio platform was handled by a personal account and a password that only one researcher had access to. Ethical approval was obtained from the Bioethics Commission of the Medical University of Lublin (KE-0254/222/2020).

Statistical analysis

Data was analysed using the IBM SPSS Statistic suite in its 26.0 version. Numbers, percentages, and means (ranges) were used for the analysis. The Pearson's correlation coefficient (r) test was used to measure the association between nurses' age, experience, spiritual care competence and caring abilities. Non-normally distributed data were analysed using the Kruskal-Wallis and Mann-Whitney U tests. The statistical significance level was set at 0.05.

Results

Participants' characteristics

The study comprised 451 professionally active nurses. Among 451 participants, 96% ($n = 433$) were females. Their average age was 40.49 ($SD = 10.92$); 85.8% of them ($n = 387$) were Catholics, and 7.3% ($n = 33$) were non-religious; 85.2% (384) were not single (married or cohabitated). More than half of the participants (52.7%) had a Bachelor's degree. As for their professional experience, their average seniority was 16.26 years and it was found that they did not receive any training related to spiritual care. More than half ($n = 256$) of the participants stated 56.8% that nurses should make an initial assessment of the patient's spiritual needs upon admission to the hospital (Table 1).

Spiritual Care Competence Scale

The scores for the total scale and the subscales of SCCS-PL described nurses' spiritual care competence. The total score of SCCS-PL was 101.22 ($SD = 17.14$), ranging from 33 to 135. The highest score on competence was obtained for the 'Attitude towards the patient's spirituality' subscale, amounting to 4.34 ($SD = 0.67$), and the nurses had the lowest score for the 'Professionalization and improvement of the quality of spiritual care' subscale, amounting to 3.35 ($SD = 0.86$).

The nurses who completed their education at the level of medical high school had a higher mean SCCS score on the Communication, individual support and counselling subscale ($SCCS = 4.19$; $p = 0.025$) than the nurses with Bachelor's ($SCCS = 3.95$) or

Table 1. Characteristics of the study participants

| Variables | | n | % |
|---|--------------------------------------|-----|------|
| Sex | Female | 433 | 96.0 |
| | Male | 18 | 4.0 |
| Civil status | Married | 308 | 68.3 |
| | In cohabitation | 76 | 16.9 |
| | Single | 58 | 12.9 |
| | Widowed | 8 | 1.7 |
| | Religious person | 1 | 0.2 |
| Place of residence | Urban area | 299 | 66.3 |
| | Rural area | 152 | 33.7 |
| Level of education | Medical High School | 34 | 7.5 |
| | Bachelor's degree | 204 | 45.2 |
| | Master's degree | 208 | 46.2 |
| | Doctoral degree | 5 | 1.1 |
| Religious | Catholic | 387 | 85.8 |
| | Protestant | 3 | 0.7 |
| | Jehovah's Witness | 2 | 0.4 |
| | Agnostic | 1 | 0.2 |
| | Catholic in past, now a non-believer | 1 | 0.2 |
| | Non-believer | 33 | 7.3 |
| | I don't want to answer | 24 | 5.4 |
| In your opinion, should a nurse make an initial assessment of the patient's spiritual needs upon admission to the hospital? | Definitely yes | 83 | 18.4 |
| | Rather yes | 173 | 38.3 |
| | Hard to say | 124 | 27.5 |
| | Rather not | 59 | 13.1 |
| | Definitely not | 12 | 2.7 |
| Does working as a nurse give you satisfaction? | Definitely yes | 216 | 47.8 |
| | Rather yes | 189 | 41.9 |
| | Hard to say | 35 | 7.8 |
| | Rather not | 8 | 1.8 |
| | Definitely not | 3 | 0.7 |

M — mean; SD — standard deviation

Master's degrees (SCCS = 3.81) (Table 2). The most statistically significant results concerned the following question: Does working as a nurse give you satisfaction? The nurses who provided an affirmative answer obtained statistically significantly higher scores in the following subscales: Assessment and

implementation of spiritual care (SCCS-PL = 3.64; p = 0.009), Communication, individual support and counselling (SCCS-PL = 3.93; p = 0.03), Professionalisation and improvement of the quality of spiritual care (SCCS-PL = 3.41; p < 0.001), and the overall score (SCCS-PL = 102.20; p = 0.002) (Table 2). The respondents who believe that nurses should assess the patient's spiritual needs upon admission to the hospital obtained statistically higher scores in all subscales and the overall score (Table 3).

Caring Abilities Inventory

The total score of CAI-PL was 185.44. The highest score was observed in the Knowledge subscale, amounting to 71.57 (SD = 10.62), then on the Courage subscale nurses obtained a score of 54.22 (12.37). The lowest score was obtained in the Patience subscale with a score of 65.63 (7.69). When analysing the care abilities and socio-demographic characteristics, Polish nurses with more years of work experience obtained higher overall caring abilities scores (Table 4). In the analysis of the relationship between these variables, it was found that the level of education was associated, in a statistically significant manner, with the Knowledge dimension (p = 0.013). The nurses who were professionally satisfied with their work achieved statistically higher scores on the overall scale (p = 0.007) and in the Knowledge dimension (p < 0.001). In the Knowledge (p = 0.002) and Patience (p = 0.019) dimensions, and the overall score (p < 0.001), a statistically significant difference was observed between the nurses who claimed that an initial spiritual needs assessment should be done and those who claimed otherwise (Table 5).

Correlation between spiritual care competence and caring abilities

Table 6 shows a positive Pearson's correlation between the dimensions of the SCCS and those of the CAI, whereby all dimensions (apart from 'Courage' and 'Assessment and implementation of spiritual care', and 'Courage' and 'Professionalisation and improvement of the quality of spiritual care') were positively correlated, with the correlation coefficient ranging from 0.113 to 0.458. The total score of spiritual care competence was also positively correlated with the level of caring abilities (p < 0.01).

Discussion

Nurses, with whom the patient and the caregivers spend most of their time, should have sufficient knowledge and experience to be able to provide spiritual care. In this study, the total score of Polish nurses' spi-

Table 2. The level of education and work satisfaction in relation to the results of SCCS

| The level of education | | | | | | | |
|--|---------------------|-------|-------------------|-------|-----------------|-------|--------------|
| | Medical high school | | Bachelor's degree | | Master's degree | | p-value* |
| | M | SD | M | SD | M | SD | |
| Attitude towards the patient's spirituality | 4.39 | 0.70 | 4.31 | 0.62 | 4.35 | 0.70 | 0.318 |
| Assessment and implementation of spiritual care | 3.81 | 0.55 | 3.64 | 0.73 | 3.54 | 0.88 | 0.218 |
| Referral, consultation and spiritual care | 4.23 | 0.47 | 4.11 | 0.59 | 4.01 | 0.72 | 0.206 |
| Communication, individual support and counselling | 4.19 | 0.50 | 3.95 | 0.66 | 3.81 | 0.86 | 0.025 |
| Professionalisation and improvement of the quality of spiritual care | 3.61 | 0.59 | 3.36 | 0.80 | 3.30 | 0.94 | 0.13 |
| Does working as a nurse give you satisfaction? | | | | | | | |
| | Yes | | No | | Hard to say | | p-value* |
| | M | SD | M | SD | M | SD | |
| Attitude towards the patient's spirituality | 4.35 | 0.67 | 4.27 | 0.83 | 4.21 | 0.65 | 0.38 |
| Assessment and implementation of spiritual care | 3.64 | 0.79 | 3.10 | 1.02 | 3.33 | 0.79 | 0.009 |
| Referral, consultation and spiritual care | 4.09 | 0.62 | 3.73 | 1.10 | 3.92 | 0.74 | 0.231 |
| Communication, individual support and counselling | 3.93 | 0.74 | 3.20 | 0.99 | 3.83 | 0.85 | 0.03 |
| Professionalisation and improvement of the quality of spiritual care | 3.41 | 0.83 | 2.58 | 1.10 | 2.91 | 0.91 | 0.001 |
| Total score | 102.20 | 16.73 | 87.18 | 23.03 | 94.31 | 16.79 | 0.002 |

*p-value < 0.05, M — mean; SD — standard deviation

Table 3. The relationship between the nurse's opinion according assessment of the spiritual needs of the patient and scores of the SCCS

| | Yes | | No | | Hard to say | | p-value* |
|--|------|------|------|------|-------------|------|-------------------|
| | M | SD | M | SD | M | SD | |
| Attitude toward the patient's spirituality | 4.38 | 0.70 | 4.22 | 0.66 | 4.31 | 0.58 | 0.03 |
| Assessment and implementation of spiritual care | 3.74 | 0.74 | 3.22 | 0.91 | 3.53 | 0.77 | < 0.001 |
| Referral, consultation and spiritual care | 4.15 | 0.63 | 3.86 | 0.72 | 4.03 | 0.61 | 0.002 |
| Communication, individual support and counselling | 4.06 | 0.69 | 3.51 | 0.86 | 3.79 | 0.73 | < 0.001 |
| Professionalisation and improvement of the quality of spiritual care | 3.53 | 0.79 | 2.84 | 0.99 | 3.27 | 0.78 | < 0.001 |

*p-value < 0.05; M — mean; SD — standard deviation

Table 4. Correlations between the respondents' ages and the CAI scores

| | Age | Work experience |
|-------------|----------|-----------------|
| Knowledge | 0.217* | 0.189* |
| Courage | -0.070 | -0.055 |
| Patience | 0.119*** | 0.139** |
| Total score | 0.107*** | 0.109*** |

*p-value < 0.001; **p-value < 0.01; ***p-value < 0.05

ritual care competence was 101.22 (SD = 17.14) and it was at a high level. In line with the results of the present study, the professional competence of nurses was estimated as optimal in earlier investigations [19] and much higher than the result for Taiwan's clinical nurses [20]. In this and recent studies, the majority of nurses were Catholic [21] or Muslim [22]; however, in Hsieh's studies almost half of the nurses (42.7%) had no religious preference [20]. These findings confirm the impact of the nurse's core beliefs and values on the delivery of patient care, and some researchers suggest that similar faith of nurses allows them to provide spiritual care at a higher level [16].

The study findings also show the correlations observed by previous researchers who have found the importance of work experience and age for the level of spiritual competence of nurses [23]. The average age of a Polish nurse in 2022 was 53.7 years [15]. The study found that older nurses, with more years of experience, got higher scores, both in total and in individual domains. The age of nurses and their work experience trigger an increased awareness of the spiritual needs of patients, which is due to, among others, greater life and professional experience, which also allows for the growth of personal spirituality [24]. The results of the research are confirmed by Kim's research, who stated that, with the increase in the seniority of nurses, their perception of spiritual care increased, as younger nurses and nurses with less professional experience were characterized by low competencies in the field of spiritual care.

In previous studies, the nurses who were enthusiastic about providing spiritual care to patients showed a high sensitivity to spiritual care [25]. The results of the present study revealed that most of the Polish nurses who believed that patients' spiritual needs should be assessed upon admission to the hospital obtained statistically significantly higher scores in all the subscales and a higher overall score by over 13 points. These results are supported by a study conducted by Aldaz which revealed that the nurses who believed that spiritual care activities were effective with patients had high spiritual competence. The

Table 5. The relationships between the level of education, postgraduate education, opinion about work, spiritual assessment and CAI scores

| | Knowledge | | | Courage | | | Patience | | | Total score | | |
|--|---------------------|-------|---|------------------|-------|---|------------------|-------|---|---------------------|-------|---|
| | M | SD | p | M | SD | p | M | SD | p | M | SD | p |
| Level of education | | | | | | | | | | | | |
| Medical high school | 76.09 | 11.34 | | 55.24 | 11.82 | | 66.65 | 8.47 | | 191.74 | 22.26 | |
| Bachelor's degree | 71.23 | 9.92 | | 53.38 | 12.61 | | 66.10 | 6.79 | | 184.64 | 18.86 | |
| Master's degree | 71.19 | 11.04 | | 54.87 | 12.24 | | 65.01 | 8.33 | | 185.21 | 22.70 | |
| Statistic | p = 0.013 | | | p = 0.455 | | | p = 0.442 | | | p = 0.086 | | |
| Does working as a nurse give you satisfaction? | | | | | | | | | | | | |
| Yes | 72.28 | 10.08 | | 54.37 | 12.44 | | 65.90 | 7.32 | | 186.55 | 20.50 | |
| No | 63.73 | 17.88 | | 56.27 | 11.23 | | 60.45 | 15.53 | | 174.73 | 31.38 | |
| Hard to say | 65.83 | 11.48 | | 51.89 | 12.02 | | 64.14 | 7.92 | | 176.00 | 20.83 | |
| Statistic | p < 0.001 | | | p = 0.519 | | | p = 0.258 | | | p = 0.007 | | |
| Should a nurse make an initial assessment of the patient's spiritual needs upon admission to the hospital? | | | | | | | | | | | | |
| Yes | 72.92 | 10.29 | | 55.00 | 12.99 | | 66.38 | 7.29 | | 188.25 | 20.28 | |
| No | 68.70 | 11.72 | | 51.87 | 10.85 | | 63.77 | 8.63 | | 178.59 | 20.18 | |
| Hard to say | 70.44 | 10.26 | | 53.98 | 11.78 | | 65.13 | 7.75 | | 183.58 | 22.12 | |
| Statistic | p = 0.002 | | | p = 0.107 | | | p = 0.019 | | | p < 0.001 | | |

p-value < 0.05; M — mean; SD — standard deviation

Table 6. Correlations between the SCCS and the CAI

| | Knowledge | Courage | Patience | Total score |
|--|-----------|----------|----------|-------------|
| Attitude towards the patient's spirituality | 0.244* | 0.258* | 0.292* | 0.368* |
| Assessment and implementation of spiritual care | 0.327* | 0.064 | 0.200* | 0.265* |
| Referral, consultation and spiritual care | 0.419* | 0.142** | 0.347* | 0.404* |
| Communication, individual support and counselling | 0.382* | 0.113*** | 0.279* | 0.346* |
| Professionalisation and improvement of the quality of spiritual care | 0.454* | 0.078 | 0.244* | 0.350* |
| Total score | 0.458* | 0.134** | 0.310* | 0.406* |

*p-value <0.001; **p-value <0.01; ***p-value <0.05

high awareness of nurses about the effectiveness of spiritual care is associated with a higher level of their spiritual sensitivity and ability to provide spiritual care [26].

The caring abilities of Polish nurses have rarely been studied at the national level. The analysis of the competence to provide spiritual care and the caring abilities of Polish nurses constitutes pioneering research in Poland. The identification of the basic values in nursing practice, which is caring and its level analysis, contributes to the development of educational processes at various levels of education, through which nurses can strengthen or shape their caring abilities. In this study, the overall score of the CAI indicates that Polish nurses display low caring abilities, which is in line with previous studies [27]. The analysis showed that the nurses' care ability varies for each dimension, with 'knowledge' being the highest, followed by 'patience', and 'courage' being the lowest, and all of the three dimensions were lower than the Nkongho norms. The research results seem to be related to the trend of medicalisation of nursing education described by Jeffers [28]. The education of Polish nurses is dominated by medical and scientific education, and students and nurses are more interested in technical knowledge and in gaining new competencies and qualifications [29]. This is in line with previous research conducted by Wang, which showed that nursing students focus more on the technical dimension of care than on the emotional dimension, which results in a low level of caring abilities measured by the CAI [30].

The study findings indicate that the nurses who were professionally satisfied with their work achieved statistically higher scores on the overall scale ($p = 0.007$). The results of previous studies show that job satisfaction is a very important variable and is related to professional burnout [31]. The studies conducted by Mohammadi and Moseley indicated that the caring abilities of nurses are directly cor-

related with their professional quality of life and job satisfaction, and inversely correlated with the Burnout Inventory. Previous studies have shown that a lower level of professional burnout is associated with an increase in nurses' caring abilities [32]. Professional burnout among Polish is at an average level [33]. The average levels of emotional exhaustion, depersonalisation and job dissatisfaction are closely related to a low level of caring abilities. Undoubtedly, special attention should be paid to the problem of professional burnout and appropriate preventive programmes should be implemented in order to reduce the incidence of this phenomenon, which may contribute to raising the level of caring abilities among Polish nurses.

In the current study, the results indicate that clinical experience has an impact on caring abilities. The nurses with higher professional experience obtained higher results in the Knowledge ($p < 0.001$) and Patience subscales ($p < 0.01$), and in the total score ($p < 0.05$). This is consistent with other research findings claiming that clinical practice is an important contributor to learning about caring [34]. One of the possible explanations for this phenomenon can be related to the fact that nurses with longer work experience have contact with more patients and caregivers, which enables them to establish a greater amount of therapeutic relationships. Work experience can also allow for a deeper understanding of professional priorities and attention to humanistic care, other than just the technical dimension of care [30].

The results also show a statistically significant correlation between the CAI and the SCCS scores ($p < 0.001$). The confirmation of the present research can be found in the results obtained by Hoover, who showed that spiritual care has a major impact on the caring abilities of nursing students [35]. In his research, Hoover confirmed the present thesis that the higher the level of spirituality and spiritual care, the better the nursing behaviour. In addition, the nurses who

had a positive attitude towards spiritual care obtained higher scores for caring abilities [35].

Study limitations

The study was conducted online and the data were collected only from 38% of the respondents who had opened the online form. The study covered only people who had access to the Internet and obtained information from one of the distribution channels. Moreover, the associations between nurses' spiritual care competence and caring abilities were investigated. Other nursing factors that may also correlate with spiritual health, should be addressed in future studies. It would be also interesting to examine whether the level of caring abilities and competencies to provide spiritual care changes over time, and if so, why, and to investigate the dynamic transformations of Polish nursing. Further longitudinal studies need to be conducted to provide evidence of the potential associations between nurses' caring abilities, spiritual care competence, and other variables.

Conclusions

The SCCS and the CAI are holistic methods of assessing spiritual care competencies in nursing practice and testing nurses' caring abilities. Nurses in Poland have poor humanistic caring abilities while spiritual care competence is at a high level. The value of humanistic care should be promoted and put into practice through undergraduate and postgraduate education of nurses. In order to strengthen the caring abilities and competence to provide spiritual care, it should be based on the clinical experience of students and nurses. The present study indicated that age, level of education, job satisfaction and work experience were the main influencing factors of the overall caring ability and spiritual care competence. Nurse management systems should support nurses, especially young ones without substantial work experience, instructing them on how to practise patient-centred care based on Watson's caring theory. Improving nursing skills in the field of spiritual care is a key factor in maintaining the quality of care provided to patients, especially in the most basic human experience of dying.

Article information and declarations

Data availability statement

The data that support the findings of this study are available from the corresponding author MM upon reasonable request.

Ethics statement

The standards stipulated in the Declaration of Helsinki were followed during each step of the study. Ethical approval was obtained from the Bioethics Commission of the Medical University of Lublin (KE-0254/222/2020).

Author contributions

Conception: MM, BD. Design: MM, BD. Execution: MM. Interpretation of the data: MM, BD. Writing-original draft preparation: MM. Supervision: BD. Writing, reviewing and editing: MM, BD.

Funding

This research received no grant from any founding agency in public, commercial or not-for-profit sectors.

Acknowledgements

We are thankful to all participants in the study.

Conflict of interest

No potential conflict of interest was reported by the authors. The authors received no financial support for the research, authorship, and/or publication of this article.

Supplementary materials

The Supplementary materials for this article can be found online at https://journals.viamedica.pl/palliative_medicine_in_practice/article/view/PMPLa2023.0022#supplementaryFiles.

References

1. Zamanzadeh V, Jasemi M, Valizadeh L, et al. Effective factors in providing holistic care: a qualitative study. *Indian J Palliat Care*. 2015; 21(2): 214–224, doi: 10.4103/0973-1075.156506, indexed in Pubmed: 26009677.
2. Lee SE, Lee MH, Peters AB, et al. Assessment of patient safety and cultural competencies among senior baccalaureate nursing students. *Int J Environ Res Public Health*. 2020; 17(12), doi: 10.3390/ijerph17124225, indexed in Pubmed: 32545746.
3. Leadership alliance for the care of dying people. Commitment Statements [Internet]. 2014. <https://www.bl.uk/collection-items/leadership-alliance-for-the-care-of-dying-people-lacdp-commitment-statements> (18.05.2023).
4. Nursing and Midwifery Council. Future nurse: Standards of proficiency for registered nurses. [Internet]. 2018. <https://www.nmc.org.uk/globalassets/sitedocuments/education-standards/future-nurse-proficiencies.pdf> (18.05.2023).
5. Biro AL. Creating conditions for good nursing by attending to the spiritual. *J Nurs Manag*. 2012; 20(8): 1002–1011, doi: 10.1111/j.1365-2834.2012.01444.x, indexed in Pubmed: 23151102.
6. Zehtab S, Adib-Hajbaghery M. The importance of spiritual care in nursing. *Nurs Midwifery Stud*. 2014; 3(3): e22261, doi: 10.17795/nmsjournal22261, indexed in Pubmed: 25699285.

7. Koenig HG. Religion, spirituality, and health: the research and clinical implications. *ISRN Psychiatry*. 2012; 1–33, doi: [10.5402/2012/278730](https://doi.org/10.5402/2012/278730), indexed in Pubmed: [23762764](https://pubmed.ncbi.nlm.nih.gov/23762764/).
8. Kowalska W, Leśnierowska A, Mazur E, et al. Analysis of sociodemographic and clinical factors determining experience and satisfaction with nursing care among cardio-surgical unit patients. *Medical Stud*. 2022; 38(1): 31–43, doi: [10.5114/ms.2022.115145](https://doi.org/10.5114/ms.2022.115145).
9. Hökkä M, Martins Pereira S, Pölkki T, et al. Nursing competencies across different levels of palliative care provision: A systematic integrative review with thematic synthesis. *Palliat Med*. 2020; 34(7): 851–870, doi: [10.1177/0269216320918798](https://doi.org/10.1177/0269216320918798), indexed in Pubmed: [32452294](https://pubmed.ncbi.nlm.nih.gov/32452294/).
10. Chahrour WH, Hvidt NC, Hvidt EA, et al. Learning to care for the spirit of dying patients: the impact of spiritual care training in a hospice-setting. *BMC Palliat Care*. 2021; 20(1): 115, doi: [10.1186/s12904-021-00804-4](https://doi.org/10.1186/s12904-021-00804-4), indexed in Pubmed: [34273974](https://pubmed.ncbi.nlm.nih.gov/34273974/).
11. Mehdi A, Hossaini SM. The relationship of spiritual health with quality of life, mental health, and burnout: the mediating role of emotional regulation. *Iran J Psychiatry*. 2018; 13(1): 22–31, indexed in Pubmed: [29892314](https://pubmed.ncbi.nlm.nih.gov/29892314/).
12. Batstone E, Bailey C, Hallett N. Spiritual care provision to end-of-life patients: a systematic literature review. *J Clin Nurs*. 2020; 29(19-20): 3609–3624, doi: [10.1111/jocn.15411](https://doi.org/10.1111/jocn.15411), indexed in Pubmed: [32645236](https://pubmed.ncbi.nlm.nih.gov/32645236/).
13. Jo K, Park A, Lee J, et al. The effect of suffering experience, empathy ability, caring behaviors on terminal care performance of clinical nurses. *Korean J Hosp Palliat Care*. 2015; 18(4): 276–284, doi: [10.14475/kjhpc.2015.18.4.276](https://doi.org/10.14475/kjhpc.2015.18.4.276).
14. Raosoft I. Raosoft sample size calculator. <http://www.raosoft.com/samplesize.html> [Internet]. 2004 <http://www.raosoft.com/samplesize.html> (18.05.2023).
15. Katastrofa kadrowa pielęgniarek i położnych. [Internet]. 2021. <https://nipip.pl/raport2021/> (18.05.2023).
16. van Leeuwen R, Tiesinga LJ, Middel B, et al. The validity and reliability of an instrument to assess nursing competencies in spiritual care. *J Clin Nurs*. 2009; 18(20): 2857–2869, doi: [10.1111/j.1365-2702.2008.02594.x](https://doi.org/10.1111/j.1365-2702.2008.02594.x), indexed in Pubmed: [19220618](https://pubmed.ncbi.nlm.nih.gov/19220618/).
17. Machul M, van Leeuwen R, Ozga D, et al. The level of spiritual care competence of Polish nurses and the psychometric properties of the spiritual care competence scale (SCCS). *BMC Nurs*. 2022; 21(1): 106, doi: [10.1186/s12912-022-00889-z](https://doi.org/10.1186/s12912-022-00889-z), indexed in Pubmed: [35524308](https://pubmed.ncbi.nlm.nih.gov/35524308/).
18. Nkongho N. *Assessing and measuring caring in nursing and health sciences*. Springer, New York. 2009.
19. Hellman A, Williams W, Hurley S. Meeting spiritual needs. *J Christ Nurs*. 2015; 32(4): 236–241, doi: [10.1097/cnj.0000000000000207](https://doi.org/10.1097/cnj.0000000000000207).
20. Hsieh SI, Hsu LL, Kao CY, et al. Factors associated with spiritual care competencies in Taiwan's clinical nurses: A descriptive correlational study. *J Clin Nurs*. 2020; 29(9-10): 1599–1613, doi: [10.1111/jocn.15141](https://doi.org/10.1111/jocn.15141), indexed in Pubmed: [31855298](https://pubmed.ncbi.nlm.nih.gov/31855298/).
21. Attard J, Baldacchino DR, Camilleri L. Nurses' and midwives' acquisition of competency in spiritual care: a focus on education. *Nurse Educ Today*. 2014; 34(12): 1460–1466, doi: [10.1016/j.nedt.2014.04.015](https://doi.org/10.1016/j.nedt.2014.04.015), indexed in Pubmed: [24814103](https://pubmed.ncbi.nlm.nih.gov/24814103/).
22. Azarsa T, Davoodi A, Khorami Markani A, et al. Spiritual wellbeing, attitude toward spiritual care and its relationship with spiritual care competence among critical care nurses. *J Caring Sci*. 2015; 4(4): 309–320, doi: [10.15171/jcs.2015.031](https://doi.org/10.15171/jcs.2015.031), indexed in Pubmed: [26744730](https://pubmed.ncbi.nlm.nih.gov/26744730/).
23. Boutell KA, Bozett FW. Nurses' assessment of patients' spirituality: continuing education implications. *J Contin Educ Nurs*. 1990; 21(4): 172–176, doi: [10.3928/0022-0124-19900701-10](https://doi.org/10.3928/0022-0124-19900701-10), indexed in Pubmed: [2116464](https://pubmed.ncbi.nlm.nih.gov/2116464/).
24. Oswald K. *Nurses' perceptions of spirituality and spiritual care*. Drake University, Des Moines 2004: USA.
25. Ozbasaran F, Ergul S, Temel AB, et al. Turkish nurses' perceptions of spirituality and spiritual care. *J Clin Nurs*. 2011; 20(21-22): 3102–3110, doi: [10.1111/j.1365-2702.2011.03778.x](https://doi.org/10.1111/j.1365-2702.2011.03778.x), indexed in Pubmed: [21797943](https://pubmed.ncbi.nlm.nih.gov/21797943/).
26. Aldaz BE, Treharne GJ, Knight RG, et al. Oncology healthcare professionals' perspectives on the psychosocial support needs of cancer patients during oncology treatment. *J Health Psychol*. 2017; 22(10): 1332–1344, doi: [10.1177/1359105315626999](https://doi.org/10.1177/1359105315626999), indexed in Pubmed: [26837692](https://pubmed.ncbi.nlm.nih.gov/26837692/).
27. Xu T, Wang Y, Wang R, et al. Predictors of caring ability and its dimensions among nurses in China: a cross-sectional study. *Scand J Caring Sci*. 2021; 35(4): 1226–1239, doi: [10.1111/scs.12941](https://doi.org/10.1111/scs.12941), indexed in Pubmed: [33615516](https://pubmed.ncbi.nlm.nih.gov/33615516/).
28. Jeffers S. Nurse faculty perceptions of end-of-life education in the clinical setting: a phenomenological perspective. *Nurse Educ Pract*. 2014; 14(5): 455–460, doi: [10.1016/j.nepr.2014.03.009](https://doi.org/10.1016/j.nepr.2014.03.009), indexed in Pubmed: [24745790](https://pubmed.ncbi.nlm.nih.gov/24745790/).
29. Dobrowolska B, Mazur E, Pilewska-Kozak A, et al. Predicted difficulties, educational needs, and interest in working in end of life care among nursing and medical students. *Nurse Educ Today*. 2019; 83: 104194, doi: [10.1016/j.nedt.2019.08.012](https://doi.org/10.1016/j.nedt.2019.08.012), indexed in Pubmed: [31493620](https://pubmed.ncbi.nlm.nih.gov/31493620/).
30. Wang Y, Zhang Xi, Xie Q, et al. Humanistic caring ability of midwifery students in China and its associated factors: A multi-centre cross-sectional study. *Nurse Educ Today*. 2022; 111: 105276, doi: [10.1016/j.nedt.2022.105276](https://doi.org/10.1016/j.nedt.2022.105276), indexed in Pubmed: [35131563](https://pubmed.ncbi.nlm.nih.gov/35131563/).
31. Dos Santos Alves DF, da Silva D, de Brito Guirardello E. Nursing practice environment, job outcomes and safety climate: a structural equation modelling analysis. *J Nurs Manag*. 2017; 25(1): 46–55, doi: [10.1111/jonm.12427](https://doi.org/10.1111/jonm.12427), indexed in Pubmed: [27730709](https://pubmed.ncbi.nlm.nih.gov/27730709/).
32. Mohammadi M, Peyrovi H, Mahmoodi M. The relationship between professional quality of life and caring ability in critical care nurses. *Dimens Crit Care Nurs*. 2017; 36(5): 273–277, doi: [10.1097/DCC.0000000000000263](https://doi.org/10.1097/DCC.0000000000000263), indexed in Pubmed: [28777112](https://pubmed.ncbi.nlm.nih.gov/28777112/).
33. Gotlib J, Panczyk M, Zarzeka A, et al. Wypalenie zawodowe w grupie pielęgniarek — przegląd aktualnego polskiego piśmiennictwa naukowego. *Pielęgniarstwo Polskie*. 2017; 63(1): 125–130, doi: [10.20883/pielpol.2017.17](https://doi.org/10.20883/pielpol.2017.17).
34. Nelms TP, Jones JM, Gray DP. Role modeling: a method for teaching caring in nursing education. *J Nurs Educ*. 1993; 32(1): 18–23, doi: [10.3928/0148-4834-19930101-06](https://doi.org/10.3928/0148-4834-19930101-06), indexed in Pubmed: [8380201](https://pubmed.ncbi.nlm.nih.gov/8380201/).
35. Hoover J. The personal and professional impact of undertaking an educational module on human caring. *J Adv Nurs*. 2002; 37(1): 79–86, doi: [10.1046/j.1365-2648.2002.02051.x](https://doi.org/10.1046/j.1365-2648.2002.02051.x), indexed in Pubmed: [11784401](https://pubmed.ncbi.nlm.nih.gov/11784401/).