



POLISH HEART JOURNAL

Kardiologia Polska

The Official Peer-reviewed Journal
of the Polish Cardiac Society
since 1957

Online first

This is a provisional PDF only. Copyedited and fully
formatted version will be made available soon

ISSN 0022-9032

e-ISSN 1897-4279

Malnutrition as a marker of carotid in-stent restenosis

Authors: Elena García-Campos Losada, Sergio Raposeiras Roubin

Article type: Editorial

Received: October 10, 2024

Accepted: October 11, 2024

Early publication date: October 24, 2024

This article is available in open access under Creative Commons Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

Malnutrition as a marker of carotid in-stent restenosis

Elena García-Campos Losada¹, Sergio Raposeiras Roubín^{1, 2, 3}

¹Cardiology Department. University Hospital Álvaro Cunqueiro, Vigo, Spain

²Instituto de Investigación Sanitaria Galicia Sur, Vigo, Spain

³Medicine Department, Vigo University. Vigo, Spain

Related article

by Karaduman et al.

Correspondence to:

Sergio Raposeiras Roubín, MD, PhD,

Cardiology Department,

Hospital Álvaro Cunqueiro,

Estrada de Clara Campoamor,

341, 36213 Vigo, Pontevedra, Spain,

phone: +34 62 017 03 87,

e-mail: raposeiras26@hotmail.com

In the latest issue of the *Polish Heart Journal*, the article by Karaduman et al. [1], titled “Predictive Significance of the Prognostic Nutritional Index (PNI) for In-Stent Restenosis Following Carotid Artery Stenting”, explores an interesting hypothesis: PNI — a simple and traditionally tool used to assess the nutritional and immune status — can serve as a predictor for in-stent restenosis (ISR) in patients who have undergone carotid artery stenting (CAS). This hypothesis is based on the premise that nutritional and inflammatory states may influence restenosis, as do other factors such as age, female sex, diabetes or dyslipidemia, as reported in the International Carotid Stenting Study study [2].

CAS has emerged as an intervention for patients with carotid artery disease, as an alternative to carotid endarterectomy. CAS helps to prevent ischemic strokes by revascularizing symptomatic or asymptomatic carotid stenosis. In the CREST trial, no significant difference in the estimated 4-year rates of the primary end point (stroke, myocardial infarction, or death from any cause during the periprocedural period or any

ipsilateral stroke within 4 years after randomization) was found between the stenting group and the endarterectomy group [3]. Despite those benefits, ISR remains a significant concern, often leading to repeat interventions and compromised outcomes.

ISR is due to a simultaneous process of neointimal hyperplasia and smooth muscle cell proliferation within the stent, leading to reforming the atherosclerotic plaque [4]. Traditionally, risk assessment for ISR has focused on factors like lesion characteristics, procedural details, and patient comorbidities. However, Karaduman et al. introduce an innovative approach by examining the predictive value of PNI, reflecting patients' nutritional and inflammatory status, which can be calculated using serum albumin and blood lymphocyte count. Recently, the PNI has been validated to identify the negative impact of malnutrition on the prognosis of cardiovascular diseases (heart failure, valvular heart disease, atrial fibrillation, and acute coronary syndrome) [5–7]. Compared to other clinical variables, malnutrition has the advantage of being a modifiable risk factor on which physicians could act.

Karaduman et al. report a retrospective single-center study of 819 patients who were subdivided into groups in line with the presence or absence of ISR. During a follow-up of 24 months, ISR was observed in 35 patients (4.3%). Independent predictors of ISR were total cholesterol, stent overlap, residual stenosis, and lower PNI. The best cut-off for PNI to predict IST was 39.5 points, with a sensitivity of 83% and a specificity of 77%. This result reinforces the prior findings of the study of Öcal et al. [8] who found that PNI was independently associated with long-term mortality and major stroke in CAS patients.

The strength of the study by Karaduman et al. lies in the methodological rigor, using a well-defined cohort, and performing an appropriate multivariate analysis helping to control confounding factors. Authors identify a simple, cost-effective biomarker that could have significant clinical implications. Nevertheless, there are certain considerations to address. The study's retrospective nature introduces potential biases, also the reliance on single-center data might limit the generalizability of the findings. Given that ISR can occur over varying time frames, short-term follow-up could not capture late events. Future prospective, multi-center studies would be valuable to validate these results across diverse populations and settings. Additionally, while PNI offers promise, it should be considered alongside other established risk factors rather than as a standalone predictor.

The incorporation of PNI into risk stratification models could refine our ability to identify patients at higher risk for adverse events following CAS, thus guiding more personalized and preventive care strategies. Identifying patients at higher risk could lead

to more targeted surveillance and intervention strategies, such as optimizing nutritional status or managing inflammation, and it could be employed to mitigate the risk of IST. Therefore, future research understanding the underlying mechanisms and clinical trials are needed to validate these findings.

Article information

Conflict of interest: None declared.

Funding: None.

Open access: This article is available in open access under Creative Commons Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, which allows downloading and sharing articles with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially. For commercial use, please contact the journal office at polishheartjournal@ptkardio.pl

REFERENCES

1. Karaduman Z, Yilmaz C, Tiryaki MM, et al. Predictive significance of the nutritional index for in-stent restenosis following carotid artery stenting. *PHJ*. 2024, doi: 10.33963/v.phj.102443.
2. Bonati LH, Gregson J, McCabe DJ, et al. Restenosis and risk of stroke after stenting or endarterectomy for symptomatic carotid stenosis in the International Carotid Stenting Study (ICSS): secondary analysis of a randomized trial. *Lancet Neurol*. 2018; 17: 587–596, doi: 10.1016/S1474-4422(18)30195-9, indexed in Pubmed: 29861139.
3. Brott TG, Hobson RW, Howard G. Stenting versus endarterectomy for treatment of carotid-artery stenosis. *N Engl J Med*. 2010; 363(1): 11–23, doi: 10.1056/NEJMoa0912321, indexed in Pubmed: 20505173.
4. Megaly M, Alani F, Cheng CI, et al. Risk factors for the development of carotid artery in-stent restenosis: multivariable analysis. *Cardiovasc Revasc Med*. 2021; 24: 65–69, doi: 10.1016/j.carrev.2020.09.005, indexed in Pubmed: 32928694.
5. Raposeiras Roubín S, Abu Assi E, Cespón Fernandez M, et al. Prevalence and prognostic significance of malnutrition in patients with acute coronary syndrome. *J Am Coll Cardiol*. 2020; 76(7): 828–840, doi: 10.1016/j.jacc.2020.06.058, indexed in Pubmed: 32792081.

6. Pezzola E, Tomasoni D, Caneiro-Queija B, et al. Prevalence and prognostic significance of malnutrition in patients with secondary mitral regurgitation undergoing transcatheter edge-to-edge repair. *Catheter Cardiovasc Interv.* 2024; 104(2): 390–400, doi: 10.1002/ccd.31079, indexed in Pubmed: 38736245.
7. Stephenson SS, Kravchenko G, Korycka-Błoch R, et al. How immunonutritional markers are associated with age, sex, body mass index and the most common chronic diseases in the hospitalized geriatric population-a cross sectional study. *Nutrients.* 2024; 16(15): 2464, doi: 10.3390/nu16152464, indexed in Pubmed: 39125344.
8. Öcal L, Küp A, Keskin M, et al. Prognostic significance of pre-procedural prognostic nutritional index in patients with carotid artery stenting. *J Stroke Cerebrovasc Dis.* 2020; 29(8): 104932, doi: 10.1016/j.jstrokecerebrovasdis.2020.104932, indexed in Pubmed: 32689619.