

Atrial tachycardia during pregnancy: Postnatal treatment

Julia Woźniak^{1*}, Agnieszka Wojdyła-Hordyńska^{1,2*}, Anastazja Murawska¹, Grzegorz Hordyński^{1,2},
Marek Gierlotka^{1,2}

¹Medical Sciences Institute, University of Opole, Opole, Poland

²Department of Cardiology, University Hospital Opole, Opole, Poland

*Both authors equally contributed to the study.

Correspondence to:

Agnieszka Wojdyła-Hordyńska,
MD, PhD,
Department of Cardiology,
University of Opole,
Witosa 26, 45–040 Opole, Poland,
phone: +48 77 452 06 92,
e-mail: agnieszka.wojdyła-
hordyńska@uni.opole.pl

Copyright by the Author(s), 2025

DOI: 10.33963/v.phj.104048

Received:

October 16, 2024

Accepted:

December 13, 2024

Early publication date:

December 18, 2024

A 32-year-old woman presented to the Emergency Department with heart palpitations and increased fatigue. She was at 26 weeks' gestation. An electrocardiogram showed a narrow complex tachycardia, at an approximate rate of 150 beats per minute, with visible P waves (Figure 1A). The patient's preliminary laboratory investigations showed no abnormalities, and no structural changes were seen on echocardiography. There were no signs of fetal distress. An attempt was made to terminate the tachycardia with vagal maneuvers, but it was ineffective, and so intravenous adenosine was administered. The arrhythmia failed to convert to sinus rhythm after the first dose of 6 mg adenosine; however, the second dose of 12 mg was successful (Figure 1B). She was discharged home in good condition with a recommendation for 25 mg of metoprolol daily. In later weeks of pregnancy and the perinatal period, the patient did not report any similar symptoms, and no further tachycardia was documented. The patient's child was born at term with no complications. Unfortunately, the arrhythmia returned in the postpartum period. An electrophysiological study with low fluoroscopy technique was performed to determine the substrate of the arrhythmia, which induced atrial tachycardia at 352 ms cycle length. Using the CARTO electroanatomical mapping system, potential and activation maps were made of the right atrium, which revealed the earliest activation site in the anterolateral tricuspid annulus (Figure 1C). A series of radiofrequency ablation applications were performed, resulting in termination of the tachycardia. The patient made a full recovery and was discharged home. During a 10-month follow-up period 2 arrhythmia episodes were

reported, each lasting approximately a couple of seconds.

Hospitalizations due to arrhythmia are not common and it is estimated that out of 100 000 hospitalizations in pregnant women, 68–166 are caused by arrhythmia. The most common cases reported are common arrhythmias, such as atrial fibrillation and paroxysmal supraventricular tachycardia, which account for approximately 14% [1]. Gestational arrhythmias usually affect mothers giving birth at the age from 41 to 50 years, and the incidence of atrial tachycardia and AVNRT increases with age. The patient's history, i.e., previous episodes of SVT or structural heart disease, can also indicate a higher risk of SVT during pregnancy. Dangerous cardiac arrhythmias during pregnancy include supraventricular and ventricular tachycardia with hemodynamic symptoms. Patient care requires a multidisciplinary approach by cardiology, pediatric electrophysiology, anesthesiology, and neonatology specialists [2–4].

If a patient is experiencing episodes of supraventricular tachycardia before pregnancy and is planning to get pregnant, the best option is to perform an ablation before the planned pregnancy. During pregnancy, standard catheter ablation using fluoroscopy is not recommended. The first treatment option in hemodynamically stable patients with symptomatic or prolonged SVT confirmed by electrocardiography should be vagal maneuvers and further administration of adenosine if these prove ineffective [5].

In the case of the patient described above, the source of the arrhythmia was a focus at the lateral wall of the tricuspid annulus without low amplitude potentials in the bipolar map.




Figure 1. A. Electrocardiogram of atrial tachycardia — visible P waves in lead V1, II and aVR (blue arrows). B. Electrocardiogram of sinus rhythm after the administration of adenosine. C. High density activation map (LAT) and bipolar map of the tricuspid annulus in left anterior oblique (LAO) and right (RAO) projection. The red arrows point to the earliest activation and the location of tachycardia cessation. The mapping catheter electrocardiogram under the maps presents two last beats (pointed with red arrows) of the tachycardia and its cessation during application with the following sinus rhythm

The symptoms first appeared in the second trimester of pregnancy, which led to the suspicion of gestational etiology of the arrhythmogenic focus.

Article information

Conflict of interest: None declared.

Funding: None.

Open access: This article is available in open access under Creative Commons Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, which allows downloading and sharing articles with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially. For commercial use, please contact the journal office at polishheartjournal@ptkardio.pl

REFERENCES

1. Wojdyła-Hordyńska A. Leczenie zaburzeń rytmu serca u kobiet w ciąży na podstawie stanowiska ekspertów HRS 2023. *WDR*. 2023; 67(2): 10–18.
2. Joglar JA, Kapa S, Saarel EV, et al. 2023 HRS expert consensus statement on the management of arrhythmias during pregnancy. *Heart Rhythm*. 2023; 20(10): e175–e264, doi: [10.1016/j.hrthm.2023.05.017](https://doi.org/10.1016/j.hrthm.2023.05.017), indexed in Pubmed: [37211147](https://pubmed.ncbi.nlm.nih.gov/37211147/).
3. Li JM, Nguyen C, Joglar JA, et al. Frequency and outcome of arrhythmias complicating admission during pregnancy: Experience from a high-volume and ethnically-diverse obstetric service. *Clin Cardiol*. 2008; 31(11): 538–541, doi: [10.1002/clc.20326](https://doi.org/10.1002/clc.20326), indexed in Pubmed: [19006111](https://pubmed.ncbi.nlm.nih.gov/19006111/).
4. Vaidya VR, Arora S, Patel N, et al. Burden of arrhythmia in pregnancy. *Circulation*. 2017; 135(6): 619–621, doi: [10.1161/CIRCULATIONAHA.116.026681](https://doi.org/10.1161/CIRCULATIONAHA.116.026681), indexed in Pubmed: [28154000](https://pubmed.ncbi.nlm.nih.gov/28154000/).
5. Jiménez-Quevedo P, Alonso-Martin C, Campuzano Ruiz R, et al. Cardiovascular disease in women: Do we need new diagnostic and therapeutic strategies? *Kardiol Pol*. 2023; 81(4): 338–349, doi: [10.33963/KP.a2023.0051](https://doi.org/10.33963/KP.a2023.0051), indexed in Pubmed: [36871309](https://pubmed.ncbi.nlm.nih.gov/36871309/).