

Association between 22 cytokine gene polymorphisms and dilated cardiomyopathy in Macedonian patients

Vera Spiroska¹, Sashko Kedev¹, Slobodan Antov¹, Dejan Trajkov², Aleksandar Petlichkovski², Slavica Hristomanova², Eli Djulejic², Meri Paneva², Aleksandar Senev², Mirko Spiroski²

¹ Cardiology Clinic, Faculty of Medicine, Ss. Cyril and Methodius University, Skopje, Republic of Macedonia

² Institute of Immunobiology and Human Genetics, Faculty of Medicine, Ss. Cyril and Methodius University, Skopje, Republic of Macedonia

Abstract

Background: Inflammation is an important component in the pathogenesis of many cardiovascular diseases and one of the commonest mechanisms in cardiomyopathy. There have been several studies on the cytokine polymorphism and dilated cardiomyopathy (DCM), but the results obtained were contradictory.

Aim: To examine a possible role of 22 cytokine gene polymorphisms in host susceptibility to or protection against DCM in Macedonians.

Methods: In this study 301 healthy unrelated individuals and 52 patients with DCM were studied. Cytokine genotyping was performed by PCR with sequence-specific priming (PCR-SSP) (Heidelberg kit).

Results: After the Bonferroni adjustment, the *IL-4 -1098/T*, *IL-4 -1098/T:T*, *IL-4/TCC*, and *IL-4/TCC:TTC* cytokine genes were positively associated with DCM, while a negative association was identified for *IL-4 -1098/G*, *IL-4 -1098/G:T*, *IL-1B +3962/C:C*, *IL-4/GCC*, and *IL-4/GCC:TTC*.

Conclusions: These results suggest that some cytokine gene polymorphisms are significantly associated and affect host susceptibility/resistance to DCM in Macedonians.

Key words: dilated cardiomyopathy, cytokine polymorphism, Republic of Macedonia

Kardiol Pol 2009; 67: 1237-1247

Introduction

Dilated cardiomyopathy (DCM), a disorder characterised by cardiac dilation and reduced systolic function, represents an outcome of a heterogeneous group of inherited and acquired disorders. Myocarditis, coronary artery disease, systemic diseases and myocardial toxins have been identified as causative factors. Idiopathic DCM, in which these causes are excluded, represents approximately a half of all DCM cases [1]. This condition was diagnosed in 1426 children younger than 18 years from the United States and Canada [2]. Unlike other parts of the world, in which cardiomyopathy is rare, DCM is a major cause of heart failure in Africa [3].

Among cases of idiopathic DCM, familial form accounts for 20-25% of cases, with an exception of rare cases resulting from mutations in dystrophin (e.g. OMIM 300377.0021) [4]. Familial DCM is characterised by an autosomal dominant pattern of inheritance with age-

-related penetrance. Ventricular dilatation and systolic dysfunction usually develop in the second or third decade of life. Mutations in many other genes have been found to cause various forms of DCM (<http://www.ncbi.nlm.nih.gov/entrez/dispomim.cgi?id=115200>) [4].

Inflammation is an important component in the pathogenesis of many cardiovascular diseases and one of the commonest mechanisms in cardiomyopathy. In most cases, the role of inflammation is a natural response to injury, and an important mechanism for healing and tissue repair. However, the inflammatory response can be either inadequate or exaggerated, leading to direct injury or a severe host disease. Accumulating data have revealed an important inflammatory component in the pathogenesis of DCM, and there is growing evidence that myocarditis and DCM are closely related. The term 'cardiomyopathy' is no longer reserved for the idiopathic forms but can be used interchangeably

Address for correspondence:

Mirko Spiroski MD, PhD, Institute of Immunobiology and Human Genetics, Faculty of Medicine, Ss. Cyril and Methodius University, 1109 Skopje, PO Box 60, Republic of Macedonia, tel.: +389 2 311 05 56, fax: +389 2 311 05 58, e-mail: mspiroski@yahoo.com

Received: 29 November 2008 Accepted: 15 July 2009

with the term 'heart muscle diseases' including specific secondary forms [5].

Several studies on the cytokine polymorphism and DCM have been conducted. The association between *TNF- α* and/or *IL-10* gene polymorphism and DCM was studied [6–9]. However, the results were contradictory. Studies performed in patients with different ethnic origins yielded contrasting results [8, 9].

We have previously published data for the cytokine polymorphisms in a healthy Macedonian population [10–12]. The aim of this study was to investigate the existence of possible associations between 22 cytokine gene polymorphisms and DCM in Macedonians.

Methods

Patients

The total studied sample consisted of 353 subjects, divided into two groups: healthy individuals and patients with DCM.

Healthy individuals

There were 301 unrelated individuals, born in different parts of Macedonia who attended the Institute of Immunobiology and Human Genetics for DNA donation between 1 May 2001 and 25 April 2002 and agreed to take part in this study as a control group. Individuals with a family history of DCM were excluded from the investigation.

Dilated cardiomyopathy

Fifty two patients with a diagnosis of DCM of different aetiologies were included in this study. They were 20–59-year-old consecutive patients who attended the Clinic of Cardiology, Faculty of Medicine, Ss. Cyril and Methodius University, Skopje, Republic of Macedonia for treatment between 18 September 2000 and 21 July 2002. The diagnosis of ischemic ($n = 25$), valvular ($n = 17$), and idiopathic cardiomyopathy ($n = 10$) was confirmed according to the report of the 1995 World Health Organization [13]. All patients underwent transthoracic echocardiography (Hewlett Packard, Sonos 5500), had left ventricular (LV) dilatation (end diastolic diameter ≥ 5.5 cm), and showed an impaired LV function (LV ejection fraction $< 40\%$, or fractional shortening $< 25\%$).

All individuals were of Macedonian origin and nationality, and residents of different regions of the Republic of Macedonia. Each individual was interviewed on a one-to-one basis, his/her genealogy was recorded for the last three generations, and his/her signed consent was obtained. Admixture, if any, was recorded for each individual. Subjects with only one Macedonian parent were excluded from the study.

All of the patients and healthy individuals included in this study signed written consent to participate in the

study which was approved by the Committee of the Macedonian Ministry of Education and Science (No.13-874/3-05) and the Ethical Committee of the Medical Faculty in Skopje.

Genomic DNA isolation and storage

The DNA was isolated from peripheral blood leukocytes using the phenol-chlorophorm extraction method or BioRobot EZ1 workstation (QIAGEN) [14]. The quality and quantity of DNA were analysed by GeneQuant (Pharmacia Biotech, Uppsala, Sweden). Isolated DNA samples were stored in the Macedonian Human DNA Bank [15].

Typing methods

Cytokine genotyping was performed by PCR-SSP (Heidelberg kit). Fourteen cytokine genes with 22 single nucleotide polymorphisms (SNP) were typed: *IL-1 α* -889, *IL-1 β* -511, *IL-1 β* +3962, *IL-1R psti1970*, *IL-1RA mspa11100*, *IL-4R α* +1902, *IL-12* -1188, *IFN γ utr5644*, *TGF- β 1 cdn10*, *TGF- β 1 cdn25*, *TNF- α* -308, *TNF- α* -238, *IL-2* -330, *IL-2* +166, *IL-4* -1098, *IL-4* -590, *IL-4* -33, *IL-6* -174, *IL-6* 565, *IL-10* -1082, *IL-10* -819, and *IL-10* -592. Briefly, PCR-SSP typing Heidelberg kit consists of 48 PCR primer mixes aliquoted in 96-well PCR trays (two typings per tray). Master mix, which was supplied along with the reagents and consisted of $MgCl_2$, buffer, dNTP's, and glycerol was mixed with 1.2–3.0 μ g DNA and 20 U Taq polymerase and dispensed in 48 wells [16]. Agarose gel electrophoresis on a 2% gel revealed a positive or negative signal for specific amplification in each well. Subsequently, the results were analysed according to the interpretation scheme provided with the kit.

Statistical analysis

The population genetics analysis package, PyPop, developed by the Biostatistics Core for the Workshop [17–19], was used for analysis of the cytokine data in this study. Allele frequencies and expected Hardy Weinberg proportions (HWP) for each SNP were determined [20]. The exact test for genotype frequency deviation from HWP was calculated using the Arlequin implementation accessed via PyPop [21]. Those SNPs that did not fit HWP were evaluated to determine whether there was an excess of homozygotes or heterozygotes, or if any particular genotype frequencies were significantly different from the expected frequencies. Comparisons of different genotypes for two groups were tested by the χ^2 test with Bonferroni corrected p-value [22]. Crude odds ratios (OR), as estimates of the relative risk, were calculated within 95% CI.

Results

Cytokine alleles

Table I shows frequencies of polymorphic cytokine alleles, Fisher exact p-value, odds ratio (OR) and Wald's 95% CI in patients with DCM and healthy Macedonians.

Table I. Cytokine allele frequency, Fisher exact p-value, odds ratio and Wald's 95% confidence interval in patients with dilated cardiomyopathy and healthy Macedonian population

Cytokine polymorphism	Allele	DCM (n = 52)		Control (n = 301)		Fisher exact p-value	Odds ratio	Wald' 95% CI
		N	F	N	F			
<i>IL-1α -889</i>	C	85	0.817	482	0.814	0.940	1.021	0.596-1.750
	T	19	0.183	110	0.186	0.940	0.980	0.572-1.679
<i>IL-1β -511</i>	C	66	0.635	404	0.671	0.466	0.851	0.552-1.314
	T	38	0.365	198	0.329	0.466	1.175	0.761-1.813
<i>IL-1β +3962</i>	C	68	0.654	439	0.729	0.115	0.701	0.451-1.091
	T	36	0.346	163	0.270	0.115	1.426	0.916-2.219
<i>IL-1R psti1970</i>	C	74	0.712	399	0.662	0.329	1.255	0.795-1.981
	T	30	0.288	203	0.337	0.329	0.797	0.505-1.258
<i>IL-1RA mspa11100</i>	T	67	0.644	420	0.698	0.277	0.785	0.507-1.215
	C	37	0.356	182	0.302	0.277	1.274	0.823-1.974
<i>IL-4Rα +1902</i>	A	89	0.856	502	0.834	0.577	1.182	0.657-2.127
	G	15	0.144	100	0.166	0.577	0.846	0.470-1.523
<i>IL-12 -1188</i>	A	73	0.702	433	0.744	0.369	0.810	0.512-1.283
	C	31	0.298	149	0.256	0.369	1.234	0.779-1.954
<i>IFNγ utr5644</i>	T	52	0.500	259	0.520	0.709	0.923	0.605-1.408
	A	52	0.500	239	0.480	0.709	1.084	0.710-1.654
<i>TGF-β1 cdn10</i>	T	59	0.567	282	0.502	0.219	1.302	0.854-1.985
	C	45	0.433	280	0.498	0.219	0.768	0.504-1.171
<i>TGF-β1 cdn25</i>	G	101	0.971	532	0.947	0.290	1.899	0.569-6.340
	C	3	0.029	30	0.053	0.290	0.527	0.158-1.759
<i>TNF-α -308</i>	A	8	0.078	74	0.123	0.195	0.607	0.284-1.301
	G	94	0.922	528	0.877	0.195	1.647	0.769-3.527
<i>TNF-α -238</i>	A	3	0.029	27	0.045	0.475	0.645	0.192-2.168
	G	99	0.971	575	0.955	0.475	1.550	0.461-5.206
<i>IL-2 -330</i>	G	33	0.317	191	0.332	0.758	0.932	0.596-1.459
	T	71	0.683	383	0.667	0.758	1.073	0.686-1.679
<i>IL-2 +166</i>	G	66	0.635	422	0.735	0.036	0.626	0.403-0.971
	T	38	0.365	152	0.264	0.036	1.599	1.029-2.482
<i>IL-4 -1098</i>	G	15	0.144	176	0.308	< 0.001*	0.379	0.213-0.674
	T	89	0.856	396	0.692	< 0.001*	2.637	1.484-4.687
<i>IL-4 -590</i>	C	74	0.712	377	0.659	0.296	1.276	0.807-2.017
	T	30	0.288	195	0.341	0.296	0.784	0.496-1.239
<i>IL-4 -33</i>	C	85	0.817	479	0.837	0.612	0.869	0.504-1.498
	T	19	0.183	93	0.163	0.612	1.151	0.668-1.985
<i>IL-6 -174</i>	C	36	0.346	182	0.302	0.372	1.222	0.787-1.897
	G	68	0.654	420	0.698	0.371	0.819	0.527-1.271
<i>IL-6 nt565</i>	A	36	0.346	173	0.287	0.225	1.313	0.845-2.040
	G	68	0.654	429	0.713	0.225	0.762	0.490-1.184
<i>IL-10 -1082</i>	A	60	0.577	352	0.589	0.823	0.953	0.625-1.453
	G	44	0.423	246	0.411	0.823	1.049	0.688-1.600
<i>IL-10 -819</i>	C	79	0.760	435	0.727	0.494	1.184	0.729-1.922
	T	25	0.240	163	0.272	0.494	0.845	0.520-1.371
<i>IL-10 -592</i>	A	25	0.240	173	0.289	0.306	0.777	0.480-1.261
	C	79	0.760	425	0.710	0.306	1.286	0.793-2.086

Abbreviations: N – absolute number, F – frequency, CI – confidence interval, DCM – dilated cardiomyopathy

* statistically significant after Bonferroni adjustment (p-value × number of alleles) < 0.05

We found a negative association for *IL-4 -1998/G* allele (OR = 0.379), and *IL-2 +166/G* allele (OR = 0.626). An inverse positive association was found for the alternative alleles for the same cytokine genes: *IL-4 -1998/T* allele (OR = 2.673), and *IL-2 +166/T* allele (OR = 1.599). After the Bonferroni adjustment, only *IL-4 -1998* alleles were significantly associated with DCM (Table I).

Cytokine genotypes

Table II contains results for different cytokine genotypes found in our study.

Analysis of all cytokine genotypes showed that six genotypes were positively associated with DCM. The highest OR was found for *IL-4 -1098/T:T* (OR = 4.279), and the lowest OR for *IL-1RA mspa11100/C:T* (OR = 1.850). Three cytokine genotypes only were negatively associated with DCM: *IL-4 -1098/G:T* (OR = 0.215), *IL-1 α +3962/C:C* (OR = 0.376), and *IFN γ utr5644/A:T* (OR = 0.458). After the Bonferroni adjustment, only *IL-4 -1098/T:T*, *IL-4 -1098/G:T*, *IL-1 α +3962/C:C*, and *IL-1 β +3962/T:T* genotypes were significantly associated with DCM (Table II).

Cytokine haplotypes

Using the Heidelberg PCR-SSP kit we were able to detect true haplotypes for several genes with multiple SNPs per gene (*TGF- β 1*, *TNF- α* , *IL-2*, *IL-4*, *IL-6*, *IL-10*). Cytokine haplotype frequency in patients with DCM and healthy Macedonians, together with the Fisher exact p-value, OR and Wald's 95% CI are shown in Table III.

Significant associations with DCM were observed in two *IL-4* haplotypes and one *IL-2* haplotype. Positive associations were shown (according the level of susceptibility) for *IL-4/TCC* ($p < 0.001$), OR 2.221 (1.456-3.390); and *IL-2/TT* ($p = 0.013$), OR 1.745 (1.118-2.722). A negative association was found only for *IL-4/GCC* haplotype ($p = 0.003$), OR 0.423 (0.238-0.753). Haplotypes *IL-4/GCT*, *IL-4/GTC*, *IL-4/GTT*, *IL-6/CG*, *IL-6/GA*, *IL-10/ACA*, and *IL-10/ATC* were present only in a healthy Macedonian population, while only patients with DCM had *TGF- β 1/GG* and *TNF- α /AA* haplotypes. After the Bonferroni adjustment, only two haplotypes (*IL-4/TCC* and *IL-4/GCC*) were significantly associated with DCM (Table III).

Cytokine diplotypes (haplotype zygosity)

Cytokine diplotypes (or haplotype zygosity) are combinations of haplotypes from both parents. Table IV contains results of the cytokine diplotype analysis.

Four haplotype combinations had a positive association [*IL-4/TCC:TTC* (OR = 10.693), *IL4/TCC:TTT* (OR = 2.764), *IL-2/TT:TT* (OR = 2.495), *IL-4/GCC:TCC* (OR = 2.380), *TNF- α /GG:GG* (OR = 2.152), and *IL-6/CA:GG* (OR = 2.001)] with DCM. We found a negative association of two haplotype combinations [*IL-4/GCC:TTC* (OR = 0.071), and *IL-4/GCC:TTT* (OR = 0.156)] with DCM. We found that

TGF- β 1/CC:CG, *TNF- α /AG:AG*, *TNF- α /GA:GA*, *IL-2/GT:GG*, *IL-2/GT:TT*, *IL-4/GCT:TTT*, *IL-4/GTC:TTC*, *IL-4/GTT:TTC*, *IL-6/CG:GG*, *IL-6/GA:GG*, *IL-10/GCC:GCC*, *IL-10/ACA:GCC*, *IL-10/ACA:ATA*, and *IL-10/ATC:GCC* haplotype combinations were present only in a healthy Macedonian population. On the other hand, *TGF- β 1/CC:GG*, *TNF- α /GG:AA*, and *TNF- α /AG:GA* diplotypes were found only in patients with DCM. Only two diplotypes (*IL-4/TCC:TTC* and *IL-4/GCC:TTC*) were significantly associated with DCM, after Bonferroni correction of p-value (Table IV).

Table V presents a summary of all positive (susceptible) and negative (protective) cytokine polymorphisms. The majority of cytokine genotypes (six of them) and cytokine diplotypes (six of them) were positively associated with DCM, while only three cytokine genotypes, and two cytokine diplotypes showed a negative association. Two cytokine alleles and two cytokine haplotypes also showed a positive association with DCM. A negative association with DCM was documented for two cytokine alleles, and only one cytokine haplotype. However, only one allele, one genotype, one haplotype, and one diplotype were positively associated with DCM after Bonferroni correction of p-value. Similarly, only one allele, two genotypes, one haplotype and one diplotype were negatively associated with DCM after Bonferroni correction of p-value.

Discussion

In this study we examined 52 Macedonian patients with DCM, and found several positive and negative associations between cytokine alleles, genotypes, haplotypes, diplotypes and DCM.

The most frequent association was found between *IL-4* gene and DCM. We investigated *IL-4* polymorphism at three positions: *-1098*, *-590*, and *-33*, and we were able to investigate haplotypes and haplotype combinations (diplotypes) of this cytokine gene. We found that *IL-4 -1098/T* allele, *IL-4 -1098/T:T* genotype, *IL-4/TCC* haplotype as well as *IL-4/TCC:TTC*, *IL-4/TCC:TTT*, and *IL-4/GCC:TCC* haplotype combinations (or diplotypes) were positively associated with DCM. The highest OR was found for *IL-4/TCC:TTC* (10.693), five belongs to *IL-4* polymorphisms (*IL-4 -1098/G* allele, *IL-4 -1098/G:T* genotype, *IL-4/GCC* haplotype, as well as *IL-4/GCC:TTC* and *IL-4/GCC:TTT* haplotype combinations). The cumulative effects of *IL-4 -1098/T* alleles: OR = 2.634 in *IL-4 -1098/T* allele, doubled in *IL-4 -1098/T:T* genotype (OR = 4.729), and four times bigger in *IL-4/TCC:TTC* haplotype combination (OR = 10.693) were documented (Table V). These results are the first ones in the literature. Several SNPs (*IL-1 α -889*, *IL-1 β +3962*, *IL-2 +166*, *IL-4 -1098*, *IL-4 -590*, *IL-4 -33*, and *IL-10 -592*) in the control group were not in HWP ($p < 0.005$) (12) and we should be very careful about their associations with cardiomyopathy.

After Bonferroni correction of p-values, we found a lower number of alleles, genotypes, haplotypes, and

Table II. Cytokine genotype frequency, Fisher exact p-value, odds ratio and Wald's 95% confidence interval in dilated cardiomyopathy patients and healthy Macedonian population

Polymorphism	Genotype	DCM (n = 52)		Controls (n = 301)		Fisher exact p-value	Odds ratio	Wald' 95% CI
		N	F	N	F			
<i>IL-1α</i> -889	C:C	36	0.692	204	0.689	0.003*	0.376	0.218-0.647
	C:T	13	0.250	74	0.250	0	1	0.506-1.975
	T:T	3	0.058	18	0.061	0.931	0.946	0.268-3.332
<i>IL-1β</i> -511	C:C	21	0.404	143	0.475	0.342	0.749	0.411-1.362
	C:T	24	0.461	118	0.392	0.345	1.329	0.735-2.404
	T:T	7	0.135	40	0.133	0.973	1.015	0.428-2.406
<i>IL-1β</i> +3962	C:C	29	0.558	174	0.578	0.784	0.920	0.509-1.665
	C:T	10	0.192	91	0.302	0.105	0.550	0.264-1.143
	T:T	13	0.250	36	0.120	0.012*	2.454	1.197-5.030
<i>IL-1Rα</i> psti1970	C:C	26	0.500	133	0.442	0.437	1.263	0.701-2.277
	C:T	22	0.423	133	0.442	0.810	0.926	0.511-1.680
	T:T	4	0.077	35	0.116	0.403	0.633	0.215-1.863
<i>IL-1RA</i> mspa11100	C:C	4	0.077	30	0.100	0.608	0.753	0.254-2.233
	C:T	29	0.558	122	0.405	0.040	1.850	1.022-3.350
	T:T	19	0.365	149	0.495	0.084	0.587	0.320-1.079
<i>IL-4Rα</i> +1902	A:A	38	0.731	212	0.704	0.698	1.140	0.588-2.207
	A:G	13	0.250	78	0.259	0.889	0.953	0.484-1.878
	G:G	1	0.019	11	0.037	0.525	0.517	0.065-4.091
<i>IL-12</i> -1188	A:A	26	0.500	160	0.550	0.506	0.819	0.454-1.478
	A:C	21	0.404	113	0.388	0.833	1.067	0.584-1.948
	C:C	5	0.096	18	0.062	0.362	1.614	0.571-4.556
<i>IFNγ</i> utr5644	A:A	19	0.365	64	0.257	0.112	1.664	0.885-3.131
	A:T	14	0.270	111	0.446	0.019	0.458	0.236-0.888
	T:T	19	0.365	74	0.297	0.333	1.362	0.728-2.548
<i>TGF-β1</i> cdn10	C:C	6	0.115	65	0.231	0.061	0.433	0.177-1.061
	C:T	33	0.635	150	0.534	0.180	1.517	0.823-2.795
	T:T	13	0.250	66	0.235	0.814	1.086	0.547-2.155
<i>TGF-β1</i> cdn25	C:G	3	0.058	30	0.107	0.276	0.512	0.150-1.745
	G:G	49	0.942	251	0.893	0.277	1.952	0.573-6.650
	C:C	0	0	0	0	0	&	&
<i>TNF-α</i> -308	A:G	8	0.157	66	0.219	0.312	0.662	0.297-1.478
	G:G	43	0.843	231	0.768	0.229	1.629	0.731-3.627
	A:A	0	0	4	0.013	&	&	&
<i>TNF-α</i> -238	A:G	3	0.059	23	0.076	0.657	0.755	0.218-2.615
	G:G	48	0.941	276	0.917	0.554	1.449	0.421-4.989
	A:A	0	0	2	0.007	&	&	&
<i>IL-2</i> -330	G:G	5	0.096	27	0.094	0.962	1.024	0.376-2.795
	G:T	23	0.442	137	0.477	0.641	0.868	0.479-1.573
	T:T	24	0.462	123	0.429	0.659	1.143	0.632-2.068
<i>IL-2</i> +166	G:G	24	0.462	162	0.565	0.170	0.661	0.366-1.197
	G:T	18	0.346	98	0.341	0.948	1.021	0.549-1.900
	T:T	10	0.192	27	0.094	0.037	2.293	1.035-5.079
<i>IL-4</i> -1098	G:T	13	0.250	174	0.608	<0.001*	0.215	0.110-0.420
	T:T	38	0.731	111	0.388	<0.001*	4.279	2.218-8.257
	G:G	1	0.019	1	0.004	0.174	5.588	0.344-90.785

Abbreviations: & – cannot be calculated because expected < 5, χ^2 test, * statistically significant after Bonferroni adjustment (p -value \times number of genotypes) < 0.05. Rest of abbreviations as in Table I

Table II. continued

Polymorphism	Genotype	DCM (n = 52)		Controls (n = 301)		Fisher exact p-value	Odds ratio	Wald' 95% CI
		N	F	N	F			
IL-4 -590	C:C	23	0.442	95	0.332	0.125	1.595	0.875-2.906
	C:T	28	0.539	187	0.654	0.112	0.618	0.340-1.122
	T:T	1	0.019	4	0.014	0.773	1.382	0.151-12.621
IL-4 -33	C:C	36	0.692	209	0.731	0.568	0.829	0.435-1.579
	C:T	13	0.250	61	0.213	0.556	1.230	0.618-2.448
	T:T	3	0.058	16	0.056	0.960	1.033	0.290-3.680
IL-6 -174	C:C	3	0.058	25	0.083	0.532	0.676	0.197-2.325
	C:G	30	0.577	132	0.439	0.064	1.746	0.963-3.167
	G:G	19	0.365	144	0.478	0.131	0.628	0.342-1.153
IL-6 nt565	A:A	3	0.058	25	0.083	0.532	0.676	0.197-2.325
	A:G	30	0.577	123	0.409	0.024	1.973	1.087-3.582
	G:G	19	0.365	153	0.508	0.057	0.557	0.303-1.023
IL-10 -1082	A:A	8	0.154	70	0.234	0.199	0.595	0.267-1.323
	A:G	44	0.846	212	0.709	0.040	2.257	1.021-4.991
	G:G	0	0	17	0.057	&	&	&
IL-10 -819	C:C	29	0.558	155	0.518	0.600	1.171	0.648-2.118
	C:T	21	0.404	125	0.418	0.848	0.943	0.518-1.718
	T:T	2	0.038	19	0.064	0.481	0.590	0.133-2.610
IL-10 -592	A:A	2	0.038	28	0.094	0.189	0.387	0.089-1.677
	A:C	21	0.404	117	0.391	0.864	1.054	0.578-1.921
	C:C	29	0.558	154	0.515	0.570	1.187	0.657-2.147

Abbreviations: & – cannot be calculated because expected < 5, χ^2 test, * statistically significant after Bonferroni adjustment (p-value \times number of genotypes) < 0.05. Rest of abbreviations as in Table I

diplotypes associated with DCM in Macedonians. Several types of multiple testing corrections are used: i) Bonferroni; ii) Bonferroni Step-down (Holm); iii) Westfall and Young Permutation; and iv) Benjamini and Hochberg False Discovery Rate [22, 23]. The methods are listed in the order of their stringency, with the Bonferroni being the most stringent, and the Benjamini and Hochberg FDR being the least stringent. The more stringent the multiple testing correction, the less false positive genes are allowed. The trade-off of a stringent multiple testing corrections is that the rate of false negatives (alleles, genotypes, haplotypes, and diplotypes that are called non-significant when they are) is very high. Inclusion of Bonferroni correction of p-value in our paper means more false negatives with subsequently less significant associations with DCM.

In a paper of Bijlsma et al., 2002 [24] an incidence of rejection was significantly lower in patients who received a donor heart with the IL-4 -590/T-positive genotype compared with patients who received a heart from a /T-negative donor. Patients who had the /T-negative genotype and received a heart from a /T-positive donor, suffered significantly less from rejection than /T-negative patients who received a T-negative donor heart. This indicates that IL-4 production within the donor heart and by cells from the donor is important for reducing the incidence of

episodes of rejection [24]. Liu et al., 2006 [25] showed that sCTLA-4 levels of patients with idiopathic DCM were associated with the haplotype and genotype. Patients with -1772/TC genotype or -1772/TC -1661/AA, -1772/TC -1661/AG haplotypes had higher sCTLA-4 levels than patients with other haplotypes. The frequency of -1772/TC genotype was significantly higher in patients with low LV ejection fraction values, whereas the frequency of -1661/G allele and -1661/GG genotype was lower in idiopathic DCM patients. Levels of IL-4 were increased in this group. It was concluded that patients with idiopathic DCM have an aberrant expression of the CTLA-4 products, and the -1772 C/T and -1661 A/G polymorphisms. The two SNPs may function as genetic markers for disease susceptibility [25]. We can add that other IL-4 polymorphisms (especially IL-4 -1098) could be also responsible for the aberrant expression of CTLA-4 products in patients with DCM.

We investigated IL-2 gene polymorphism at two positions: -330 and +166, and were able to investigate haplotypes and haplotype combinations (diplotypes). We found a positive association of four IL-2 polymorphisms and one negative association of IL-2 with DCM. However, Bonferroni-corrected p-values were not significant. Huang et al., 2008 [26] investigated whether -384T/G, -475A/T, and -631G/A polymorphisms in the IL-2 gene promoter region

were associated with idiopathic DCM in a South-western Chinese Han population. They found that the *T* allele of the *IL-2* gene promoter at position -384 might increase the risk of developing idiopathic DCM [21]. Our results are the first ones that show the positive association of *IL-2 +166/T* allele, and the opposite – a negative association of *IL-2 +166/G* allele with DCM, if not corrected with Bonferroni. Homozygous combinations of *IL-2* polymorphisms (*IL-2 +166/T:T* genotype, *IL-2/TT* haplotype, and *IL-2/TT:TT* haplotype combinations) are positively associated with DCM.

We have also published for the first time that *IL-1 β +3962/T:T* homozygous genotype is positively associated with DCM, while *IL-1 β +3962/C:C* homozygous genotype is negatively associated with DCM. The *IL-1RA mspa11100/C:T* heterozygous genotype is positively associated with DCM, while *IFN γ utr5644/A:T* heterozygous genotype is negatively associated with DCM. These results suggest that *IL-1* cluster genes are included in the development of DCM.

Our results showing a positive association of *TNF- α /GG:GG* homozygous combination of haplotypes (diplotypes) with DCM are similar to the results of Ito et al., 2000 [8] in Japanese patients. Several other studies were unable to find any association between the *TNF- α* and/or *IL-10* polymorphisms and DCM [5-7, 22].

The presence and absence of certain haplotypes and haplotype combinations (diplotypes) in a healthy Macedonian population and in patients with DCM, could be a result of small frequencies of those cytokine polymorphisms.

The number of patients in our study, as well as in other published papers [6-8, 20], is very small. In the association studies, some positive results might be spurious and some negative findings might be a consequence of low statistical power. It could be due to a small sample size or methodological shortcomings, such as the selection of an inappropriate control group. It is necessary to investigate cytokine gene polymorphisms in our population in well-

Table III. Haplotype frequency of cytokine polymorphism, Fisher exact p-value, odds ratio and Wald's 95% confidence interval in dilated cardiomyopathy patients and healthy Macedonian population

Polymorphism	Haplotype	DCM (n = 52)		Control (n = 301)		Fisher exact p-value	Odds ratio	Wald' 95% CI
		N	F	N	F			
<i>TGF-β1</i>	CC	3	0.029	30	0.053	0.290	0.527	0.158-1.759
	CG	42	0.404	250	0.445	0.439	0.845	0.552-1.294
	TG	58	0.558	282	0.502	0.295	1.252	0.822-1.907
	GG	1	0.009	0	0	&	&	&
<i>TNF-α</i>	AG	7	0.068	74	0.123	0.112	0.526	0.235-1.176
	GA	2	0.020	26	0.043	0.260	0.443	0.104-1.896
	GG	92	0.902	502	0.834	0.080	1.833	0.922-3.643
	AA	1	0.010	0	0	&	&	&
<i>IL-2</i>	GG	32	0.308	178	0.310	0.723	0.921	0.586-1.450
	GT	1	0.009	14	0.024	0.346	0.388	0.051-2.986
	TG	34	0.327	244	0.425	0.061	0.657	0.422-1.022
	TT	37	0.356	138	0.240	0.013	1.745	1.118-2.722
<i>IL-4</i>	GCC	15	0.144	163	0.285	0.003*	0.423	0.238-0.753
	GCT	0	0	8	0.014	&	&	&
	GTC	0	0	4	0.007	&	&	&
	GTT	0	0	1	0.002	&	&	&
	TCC	57	0.548	202	0.353	<0.001*	2.221	1.456-3.390
	TCT	2	0.019	4	0.007	0.221	2.784	0.503-15.40
	TTC	13	0.125	110	0.192	0.102	0.600	0.324-1.112
	TTT	17	0.164	80	0.140	0.528	1.202	0.679-2.127
<i>IL-6</i>	CA	36	0.346	172	0.286	0.212	1.324	0.852-2.057
	CG	0	0	9	0.150	&	&	&
	GG	68	0.654	420	0.698	0.372	0.819	0.527-1.271
	GA	0	0	1	0.002	&	&	&
<i>IL-10</i>	ACA	0	0	12	0.020	&	&	&
	ACC	35	0.337	177	0.296	0.406	1.207	0.775-1.879
	ATA	25	0.240	161	0.269	0.538	0.859	0.529-1.395
	ATC	0	0	2	0.003	&	&	&
	GCC	44	0.423	246	0.411	0.823	1.049	0.688-1.600

Abbreviations: as in Tables I and II

Table IV. Cytokine diplotypes (haplotype zygotes), Fisher exact p-value, odds ratio and Wald's 95% confidence interval in dilated cardiomyopathy patients and healthy Macedonian population

Polymorphism	Genotype	DCM (n = 52)		Control (n = 301)		Fisher exact p-value	Odds ratio	Wald' 95% CI
		N	F	N	F			
<i>TGF-β1</i>	<i>CC:CG</i>	0	0	16	0.057	&	&	&
	<i>CC:TG</i>	3	0.058	14	0.050	0.813	1.168	0.324-4.215
	<i>CG:CG</i>	6	0.115	49	0.174	0.293	0.618	0.250-1.526
	<i>CG:TG</i>	29	0.558	136	0.484	0.329	1.344	0.741-2.438
	<i>TG:TG</i>	13	0.250	66	0.235	0.814	1.086	0.547-2.155
	<i>CC:GG</i>	1	0.019	0	0	&	&	&
<i>TNF-α</i>	<i>AG:GG</i>	6	0.117	66	0.219	0.096	0.475	0.194-1.161
	<i>GA:GG</i>	1	0.020	24	0.080	0.122	0.231	0.031-1.745
	<i>GG:GG</i>	42	0.823	206	0.684	0.044	2.152	1.007-4.601
	<i>AG:AG</i>	0	0	4	0.013	&	&	&
	<i>GG:AA</i>	1	0.020	0	0	&	&	&
	<i>GA:GA</i>	0	0	1	0.004	&	&	&
	<i>AG:GA</i>	1	0.020	0	0	&	&	&
<i>IL-2</i>	<i>GG:GG</i>	5	0.096	27	0.094	0.962	1.024	0.376-2.795
	<i>GG:TG</i>	10	0.192	85	0.296	0.125	0.566	0.271-1.180
	<i>GG:TT</i>	12	0.231	38	0.133	0.066	1.966	0.948-4.079
	<i>GT:TG</i>	1	0.019	11	0.058	0.493	0.492	0.062-3.894
	<i>TG:TG</i>	9	0.174	50	0.174	0.984	0.992	0.455-2.165
	<i>TG:TT</i>	5	0.096	48	0.168	0.194	0.530	0.200-1.401
	<i>TT:TT</i>	10	0.192	25	0.087	0.022	2.495	1.118-5.567
	<i>GT:GG</i>	0	0	1	0.003	&	&	&
	<i>GT:TT</i>	0	0	2	0.007	&	&	&
<i>IL-4</i>	<i>GCC:GCC</i>	1	0.019	1	0.003	0.174	5.588	0.344-90.785
	<i>GCC:TCC</i>	10	0.192	26	0.091	0.029	2.380	1.071-5.293
	<i>GCC:TTC</i>	2	0.039	103	0.360	< 0.001*	0.071	0.017-0.298
	<i>GCC:TTT</i>	1	0.019	32	0.112	0.038	0.156	0.021-1.165
	<i>TCC:TCC</i>	12	0.231	68	0.238	0.913	0.962	0.478-1.937
	<i>TCC:TTC</i>	11	0.211	7	0.025	< 0.001*	10.693	3.923-29.148
	<i>TCC:TTT</i>	12	0.231	28	0.098	0.006	2.764	1.301-5.875
	<i>TTT:TTT</i>	1	0.019	4	0.014	0.773	1.382	0.151-12.621
	<i>GCT:TTT</i>	0	0	8	0.028	&	&	&
	<i>GTC:TTC</i>	0	0	4	0.014	&	&	&
	<i>TCT:TTT</i>	2	0.039	4	0.014	0.219	2.82	0.503-15.809
	<i>GTT:TTC</i>	0	0	1	0.003	&	&	&
<i>IL-6</i>	<i>CA:CA</i>	3	0.058	25	0.083	0.532	0.676	0.197-2.325
	<i>CA:GG</i>	30	0.577	122	0.405	0.021	2.001	1.102-3.632
	<i>CG:GG</i>	0	0	9	0.030	&	&	&
	<i>GG:GG</i>	19	0.365	144	0.479	0.131	0.628	0.342-1.153
	<i>GA:GG</i>	0	0	1	0.003	&	&	&
<i>IL-10</i>	<i>ACC:ACC</i>	4	0.077	21	0.070	0.863	1.103	0.363-3.355
	<i>ACC:ATA</i>	2	0.038	21	0.070	0.393	0.530	0.120-2.329
	<i>ACC:GCC</i>	25	0.481	114	0.381	0.176	1.503	0.831-2.716
	<i>ATA:ATA</i>	2	0.038	19	0.064	0.481	0.590	0.133-2.610
	<i>ATA:GCC</i>	19	0.366	93	0.311	0.438	1.275	0.689-2.360
	<i>GCC:GCC</i>	0	0	17	0.057	&	&	&
	<i>ACA :GCC</i>	0	0	3	0.010	&	&	&
	<i>ACA :ATA</i>	0	0	9	0.030	&	&	&
	<i>ATC :GCC</i>	0	0	2	0.007	&	&	&

Abbreviations: as in Tables I and II

Table V. Summary of all susceptible and protective cytokine polymorphisms for dilated cardiomyopathy in Macedonian population

Cytokine	Susceptible (positive) association			Protective (negative) association		
	Polymorphism	p	Odds ratio	Polymorphism	p	Odds ratio
Alleles	<i>IL-4 -1098/T</i>	< 0.001*	2.637	<i>IL-4 -1098/G</i>	< 0.001*	0.379
	<i>IL-2 +166/T</i>	0.036	1.599	<i>IL-2 +166/G</i>	0.036	0.626
Genotypes	<i>IL-4 -1098/ T:T</i>	< 0.001*	4.279	<i>IL-4 -1098/G:T</i>	< 0.001*	0.215
	<i>IL-1? +3962/T:T</i>	0.012*	2.454	<i>IL-1β +3962/C:C</i>	0.003*	0.376
	<i>IL-2 +166/ T:T</i>	0.037	2.293	<i>IFNγ utr5644/A:T</i>	0.019	0.458
	<i>IL-10 -1082/A:G</i>	0.040	2.257			
	<i>IL-6 nt565/A:G</i>	0.024	1.973			
	<i>IL-1RA mspa11100/C:T</i>	0.040	1.850			
Haplotypes	<i>IL-4/TCC</i>	< 0.001*	2.221	<i>IL-4/GCC</i>	0.003*	0.423
	<i>IL-2/TT</i>	0.013	1.745			
Diploypes (haplotype combinations)	<i>IL-4/TCC:TTC</i>	< 0.001*	10.693	<i>IL-4/GCC:TTC</i>	< 0.001*	0.071
	<i>IL-4/TCC:TTT</i>	0.006	2.764	<i>IL-4/GCC:TTT</i>	0.038	0.156
	<i>IL-2/TT:TT</i>	0.022	2.495			
	<i>IL-4/GCC:TCC</i>	0.029	2.380			
	<i>TNF-α/GG:GG</i>	0.044	2.152			
	<i>IL-6/CA:GG</i>	0.021	2.001			

* statistically significant after Bonferroni adjustment ($p\text{-value} \times \text{number of alleles, genotypes, haplotypes or diploypes}$) < 0.05

-defined subgroups of phenotypes with more participants in order to have more precise conclusions for genetic background of DCM development in Macedonians. Multicentric studies and/or meta-analysis of the patients with DCM and an association with cytokine polymorphisms should be very useful.

It can be concluded that, at least in Macedonian patients with DCM, some cytokine polymorphisms contribute to susceptibility/protection to the disease. Ethnic factors might play a role in the variability of results in different populations. Therefore, additional studies are needed to clarify this issue.

Acknowledgements

This research is a part of the project 'Molecular analysis of cytokine gene polymorphisms in the Republic of Macedonia' supported by the Macedonian Ministry of Education and Science (Project no. 13-874/3-05). We would like to gratefully acknowledge Prof. G. Opelz and Dr. J. Mytilineos from the Institute of Immunology, Department of Transplantation Immunology, University of Heidelberg, Heidelberg, Germany, for kindly supplying the Heidelberg PCR-SSP kit reagents in this project. We thank Elena Zaharieva for sample collection, technical support, and laboratory direction.

References

- Olson TM, Thibodeau SN, Lundquist PA, et al. Exclusion of a primary gene defect at the HLA locus in familial idiopathic dilated cardiomyopathy. *J Med Genet* 1995; 32: 876-80.
- Towbin JA, Lowe AM, Colan SD, et al. Incidence, causes, and outcomes of dilated cardiomyopathy in children. *JAMA* 2006; 296: 1867-76.
- Sliwa K, Damasceno A, Mayosi BM. Epidemiology and etiology of cardiomyopathy in Africa. *Circulation* 2005; 112: 3577-83.
- Online Mendelian Inheritance in Man, OMIM (TM). Johns Hopkins University, Baltimore, MD. MIM Number: *607093:6/15/2007. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>.
- Pankuweit S, Ruppert V, Maisch B. Inflammation in dilated cardiomyopathy. *Herz* 2004; 29: 788-93.
- Kubota T, McNamara DM, Wang JJ, et al. Effects of tumor necrosis factor gene polymorphisms on patients with congestive heart failure. VEST investigators for TNF genotype analysis. Vesnarinone survival trial. *Circulation* 1998; 97: 2499-501.
- Tiret L, Mallet C, Poirier O, et al. Lack of association between polymorphisms of eight candidate genes and idiopathic dilated cardiomyopathy: the CARDIGENE study. *J Am Coll Cardiol* 2000; 35: 29-35.
- Alikasifoglu M, Tokgözoğlu L, Acil T, et al. Tumor necrosis factor- α polymorphism in Turkish patients with dilated cardiomyopathy. *Eur J Heart Fail* 2003; 5: 161-3.
- Ito M, Takahashi H, Fuse K, et al. Polymorphisms of tumor necrosis factor- α and interleukin-10 genes in Japanese patients with idiopathic dilated cardiomyopathy. *Jpn Heart J* 2000; 41: 183-91.
- Trajkov D, Arsov T, Petlichkovski A, et al. Cytokine gene polymorphisms in population of ethnic Macedonians. *Croat Med J* 2005; 46: 685-92.
- Trajkov D, Atanasovska-Stojanovska A, Petlichkovski A, et al. IL-1 Gene Cluster Polymorphisms in the Macedonian Population. *Maced J Med Sci* 2008; 1: 26-33.
- Trajkov D, Arsov T, Petlichkovski A, et al. Distribution of the 22 cytokine gene polymorphisms in healthy Macedonian population. *Bratisl Lek Listy* 2009; 110: 7-17.
- Richardson P, McKenna W, Bristow M, et al. Report of the 1995 World Health Organization/International Society and Federation of Cardiology Task Force on the definition and classification of cardiomyopathies. *Circulation* 1996; 93: 841-2.
- Towner P. Purification of DNA. In: Brown TA (ed.). *Essential molecular biology*. Oxford University Press, Oxford 1995; 47-54.

15. Spiroski M, Arsov T, Petlichkovski A, et al. Case Study: Macedonian Human DNA Bank (hDNAMKD) as a source for public health Genetics. In: Georgieva L, Burazeri G (eds.). Health determinants in the scope of new public health. *Hans Jacobs Company, Sofia* 2005; 33-44.
16. Tseng LH, Chen PJ, Lin MT, et al. Simultaneous genotyping of single nucleotide polymorphisms in the IL-1 gene complex by multiplex polymerase chain reaction-restriction fragment length polymorphism. *J Immunol Methods* 2002; 267: 151-6.
17. Lancaster A, Nelson MP, Meyer D, et al. PyPop: a software framework for population genomics: analyzing large-scale multi-locus genotype data. *Pac Symp Biocomput* 2003; 514-25.
18. Lancaster AK, Single RM, Solberg OD, et al. PyPop update-a software pipeline for large-scale multilocus population genomics. *Tissue Antigens* 2007; 69: 192-7.
19. Single RM, Meyer D, Mack SJ, et al. 14th International HLA and Immunogenetics Workshop: report of progress in methodology, data collection, and analyses. *Tissue Antigens* 2007; 69: 185-7.
20. Guo S, Thomson E. Performing the exact test of Hardy Weinberg proportion for multiple alleles. *Biometrics* 1992; 48: 361.
21. Schneider S, Roessli D, Excoffier L. Arlequin version 2.000: a software for population genetics data analysis. Geneva (Switzerland): Genetics and Biometry Laboratory, University of Geneva; 2000.
22. Rice TK, Schork NJ, Rao DC. Methods for handling multiple testing. *Adv Genet* 2008; 60: 293-308.
23. Holm S. A simple sequentially rejective Bonferroni test procedure. *Scand J Statist* 1979; 6: 65-70.
24. Bijlsma FJ, vanKuik J, Tilanus MG, et al. Donor interleukin-4 promoter gene polymorphism influences allograft rejection after heart transplantation. *J Heart Lung Transplant* 2002; 21: 340-6.
25. Liu W, Li WM, Gao C, et al. The research on associating the single nucleotide polymorphism of CTLA-4 gene promoter region with idiopathic dilated cardiomyopathy. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi* 2006; 23: 198-201.
26. Webber SA, Boyle GJ, Gribar S, et al. Polymorphisms in cytokine genes do not predict progression to end-stage heart failure in children. *Cardiol Young* 2002; 12: 461-4.
27. Bijlsma FJ, Bruggink AH, Hartman M, et al. No association between IL-10 promoter gene polymorphism and heart failure or rejection following cardiac transplantation. *Tissue Antigens* 2001; 57: 151-3.

Związek pomiędzy występowaniem polimorfizmu genu 22 cytokiny a kardiomiopatią rozstrzeniową w populacji chorych z Macedonii

Vera Spiroska¹, Sashko Kedev¹, Slobodan Antov¹, Dejan Trajkov², Aleksandar Petlichkovski², Slavica Hristomanova², Eli Djulejic², Meri Paneva², Aleksandar Senev², Mirko Spiroski²

¹ Klinika Kardiologii, Uniwersytet Świętych Cyryla i Metodego, Skopje, Macedonia

² Instytut Immunobiologii i Genetyki, Uniwersytet Świętych Cyryla i Metodego, Skopje, Macedonia

Streszczenie

Wstęp: Zapalenie jest jednym z istotnych czynników patogenetycznych wielu chorób układu sercowo-naczyniowego i częstym mechanizmem powstawania kardiomiopatii. W wielu badaniach analizowano związki pomiędzy polimorfizmem genu cytokiny a kardiomiopatią rozstrzeniową (DCM), ale wyniki były rozbieżne.

Cel: Zbadanie związku pomiędzy występowaniem polimorfizmu genu 22 cytokiny a DCM w populacji macedońskiej.

Metody: Grupa badana składała się z 301 zdrowych ochotników i 52 chorych z DCM. Badania genetyczne wykonano techniką łańcuchowej reakcji polimerazy (ang. *polymerase chain reaction*, PCR).

Wyniki: Po zastosowaniu poprawki Bonferroniego okazało się, że występowanie *IL-4 -1098/T*, *IL-4 -1098/T:T*, *IL-4/TCC*, i *IL-4/TCC:TTC* było pozytywnie związane z obecnością DCM, podczas gdy występowanie *IL-4 -1098/G*, *IL-4 -1098/G:T*, *IL-1β +3962/C:C*, *IL-4/GCC*, i *IL-4/GCC:TTC* było negatywnie związane z obecnością tej choroby.

Wnioski: Powyższe wyniki wskazują, że niektóre polimorfizmy genu cytokiny 22 są związane z występowaniem DCM w populacji macedońskiej.

Słowa kluczowe: kardiomiopatia rozstrzeniowa, polimorfizm genu cytokiny, Macedonia

Kardiol Pol 2009; 67: 1237-1247

Adres do korespondencji:

dr n. med. Mirko Spiroski, Institute of Immunobiology and Human Genetics, Faculty of Medicine, Ss. Cyril and Methodius University, 1109 Skopje, PO Box 60, Republic of Macedonia, tel.: +38 92 311 05 56, faks: +38 92 311 05 58, e-mail: mspiroski@yahoo.com

Praca wpłynęła: 29.11.2008. Zaakceptowana do druku: 15.07.2009.