

Advancing understanding of depression and cardiovascular disease risk within the social milieu

Phillip J. Tully

School of Medicine, The University of Adelaide, Adelaide, Australia

RELATED ARTICLE page 535

In the past 30 years, depression disorders and symptoms have emerged as widely accepted risk factors for the development of cardiovascular disease (CVD).^{1,2} Depression is associated with an approximately 90% increase in risk for acute myocardial infarction or coronary revascularization.³ The association between depression and incident CVD is independent of traditional CVD risk factors such as hypertension, dyslipidemia, diabetes, and obesity. However, depression is merely one of several interrelated psychosocial risk factors implicated in the development of CVD. Aside from depression, further psychosocial risk factors include other negative emotional states such as anxiety and hostility, while others fall within the broader social milieu, such as low social position, access to and attained education, stress at work or in the family, and perceived social support.^{1,2}

An important step towards improved CVD prevention strategies is recognizing that psychosocial risk factors do not function independently or occur in isolation, but rather, they cluster together within individuals or groups.⁴ This line of thought has traditionally been sorely lacking in empirical support specific to CVD, with the recognition of psychosocial risk-factor clustering largely based on narrative review papers. The study by Kozela et al⁵ in this issue of *Kardiologia Polska (Kardiol Pol)* provides an important longitudinal perspective on the cumulative effects of clustered risk factors such as low level of education, material deprivation, low perceived control, and depressive symptoms in relation to incident CVD.

Using a representative sample of 10 012 permanent residents of Kraków, Poland, included in the HAPIEE study (Health, Alcohol and Psychosocial Factors in Eastern Europe), Kozela et

al⁵ were among the first to prospectively study the cumulative effect of psychosocial factors on CVD incidence in Central and Eastern Europe. Specifically, the authors took the approach of tallying whether a person has 0, 1, 2, 3, or all 4 psychosocial risk factors. Upon examining incident CVD risks over 11 years of follow-up, they found a linear trend for an increasing number of psychosocial risk factors and CVD risk. However, as the authors acknowledge, this analytical approach assumes equal weighting of psychosocial risk factors, and precludes identification of risk factor combinations that are most cardiotoxic. This could be important given that low perceived control and depressive symptoms shared more variance with each other (ρ , -0.47 and -0.49 in men and women, respectively), while deprivation was associated with low perceived control and depressive symptoms, but to a lesser extent (ρ range, -0.30 to -0.33).⁵ Accordingly, examining linear and categorical interaction terms between depression and other psychosocial risk factors could be a fruitful avenue of epidemiological investigation to inform concerted efforts to improve CVD prevention strategies and depression treatment access within specific population strata or underresourced communities.⁶

Importantly, Kozela et al⁵ stratified all analyses by sex and confirmed a discrepancy in CVD outcomes, with population attributable risks (PARs) for women ranging from 4.1% for 4 risk factors up to 12.1% for 3 risk factors. The associated PARs in men ranged from 0% for 1 risk factor up to 3.7% for 3 risk factors. The divergence in PARs by sex lies in the fact that women, compared with men, were more susceptible to all 4 psychosocial risk factors studied. These findings⁵ provide an important replication of other studies. The Million Women Study in the United Kingdom

Correspondence to:

Dr. Phillip J. Tully,
Centre for Men's Health,
School of Medicine,
level 6 AHMS Building,
The University of Adelaide,
Adelaide SA 5005, Australia,
phone: +61 8 8313 0514,
email: phillip.tully@adelaide.edu.au

Received: May 16, 2019.

Accepted: May 17, 2019.

Published online: May 24, 2019.

Kardiol Pol. 2019; 77 (5): 507-508

doi:10.33963/KP.14844

Copyright by Polskie Towarzystwo
Kardiologiczne, Warszawa 2019

found not only that lower levels of education and greater geographical deprivation were associated with higher risks of incident CVD in women, but also that associations for education were found within every level of deprivation, and associations for deprivation were found within every level of education.⁷ Previous research also identified an impact of interactions between sex, depression, and lower level of education on CVD risk factors,⁴ as well as of interactions between sex and depression on CVD outcomes.^{8,9} More generally, Kozela et al⁵ also replicate the INTERHEART findings, which demonstrated that severe financial stress was associated with 33% increased odds for acute myocardial infarction.¹⁰

The study by Kozela et al⁵ has certain limitations. Surprisingly, although the HAPIEE study bears alcohol in its title, alcohol use was not investigated.⁵ This is an important limitation to reconcile in subsequent analyses for several reasons. Firstly, the socioeconomic status is negatively associated with alcohol misuse because low status reduces people's sense of personal control,¹¹ and perceived control was 1 of the 4 psychosocial risk factors in HAPIEE. Secondly, the Million Women Study concluded that the association of education and deprivation with incident CVD was mostly attributable to health-related behaviors including alcohol consumption.⁷ Also, depression disorders are frequently comorbid with alcohol abuse or dependence, both concurrently and longitudinally.¹² Moreover, depression and alcohol use disorders were the strongest psychiatric risk factors for earlier-onset CVD in the World Mental Health Surveys.¹³ The HAPIEE study may have included earlier-onset CVD cases, given that participants were enrolled from the age of 45 years with only 11 years of follow-up.

Nonetheless, investigation of the nexus between psychological symptoms such as depression, our vulnerability to them, and the broader social context within which we live affords novel insights into CVD mechanisms and prevention strategies. Indeed, intervention with social risk factors lies largely outside the purview of health-care clinicians, with policymakers most likely to shift broader social, political, and economic factors pertinent to CVD risk.¹⁴ Such strategies can be costly and time consuming, although the declining rates of percutaneous coronary revascularization in Poland might indirectly signify recent advances in CVD prevention.¹⁵

ARTICLE INFORMATION

DISCLAIMER The opinions expressed by the author are not necessarily those of the journal editors, Polish Cardiac Society, or publisher.

CONFLICT OF INTEREST None declared.

HOW TO CITE Tully PJ. Advancing understanding of depression and cardiovascular disease risk within the social milieu. *Kardiol Pol.* 2019; 77: 507-508. doi: 10.33963/KP.14844

REFERENCES

- 1 Ladwig KH, Lederbogen F, Albus C, et al. Position paper on the importance of psychosocial factors in cardiology: update 2013. *Ger Med Sci.* 2014; 12: Doc09.
- 2 Piepoli MF, Hoes AW, Agewall S, et al. 2016 European Guidelines on cardiovascular disease prevention in clinical practice: the sixth joint task force of the European Society of Cardiology and other societies on cardiovascular disease prevention in clinical practice (constituted by representatives of 10 societies and by invited experts). Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR). *Eur Heart J.* 2016; 37: 2315-2381.
- 3 Nicholson A, Kuper H, Hemingway H. Depression as an aetiological and prognostic factor in coronary heart disease: a meta-analysis of 146 538 participants. *Eur Heart J.* 2006; 27: 2763-2774.
- 4 Pogosova N, Kotseva K, De Bacquer D, et al. Psychosocial risk factors in relation to other cardiovascular risk factors in coronary heart disease: results from the EUROASPIRE IV survey. A registry from the European Society of Cardiology. *Eur J Prev Cardiol.* 2017; 24: 1371-1380.
- 5 Kozela M, Doryńska A, Bobak M, Pajak A. Accumulation of psychosocial risk factors and cardiovascular disease incidence. Prospective observation of the Polish HAPIEE cohort. *Kardiol Pol.* 2019; 77: 535-540.
- 6 Wells KB, Jones L, Chung B, et al. Community-partnered cluster-randomized comparative effectiveness trial of community engagement and planning or resources for services to address depression disparities. *J Gen Intern Med.* 2013; 28: 1268-1278.
- 7 Floud S, Balkwill A, Moser K, et al. The role of health-related behavioural factors in accounting for inequalities in coronary heart disease risk by education and area deprivation: prospective study of 1.2 million UK women. *BMC Med.* 2016; 14: 145.
- 8 Guimaraes PO, Granger CB, Stebbins A, et al. Sex differences in clinical characteristics, psychosocial factors, and outcomes among patients with stable coronary heart disease: insights from the STABILITY (Stabilization of Atherosclerotic Plaque by Initiation of Darapladib Therapy) trial. *J Am Heart Assoc.* 2017; 6: e006 695.
- 9 Doyle F, McGee H, Conroy R, et al. Systematic review and individual patient data meta-analysis of sex differences in depression and prognosis in persons with myocardial infarction: a MINDMAPS study. *Psychosom Med.* 2015; 77: 419-428.
- 10 Rosengren A, Hawken S, Ounpuu S, et al. Association of psychosocial risk factors with risk of acute myocardial infarction in 11 119 cases and 13 648 controls from 52 countries (the INTERHEART study): case-control study. *Lancet.* 2004; 364: 953-962.
- 11 Elliott M, Lowman J. Education, income and alcohol misuse: a stress process model. *Soc Psychiatry Psychiatr Epidemiol.* 2015; 50: 19-26.
- 12 Grant BF, Goldstein RB, Chou SP, et al. Sociodemographic and psychopathologic predictors of first incidence of DSM-IV substance use, mood and anxiety disorders: results from the Wave 2 National Epidemiologic Survey on Alcohol and Related Conditions. *Mol Psychiatry.* 2009; 14: 1051-1066.
- 13 Scott KM, de Jonge P, Alonso J, et al. Associations between DSM-IV mental disorders and subsequent heart disease onset: beyond depression. *Int J Cardiol.* 2013; 168: 5293-5299.
- 14 Hemingway H, Marmot M. Evidence based cardiology: psychosocial factors in the aetiology and prognosis of coronary heart disease. Systematic review of prospective cohort studies. *BMJ.* 1999; 318: 1460-1467.
- 15 Dudek D, Siudak Z, Legutko J, et al. Percutaneous interventions in cardiology in Poland in the year 2017. Summary report of the Association of Cardiovascular Interventions of the Polish Cardiac Society AISN PTK and Jagiellonian University Medical College. *Postepy Kardiol Interwencyjne.* 2018; 14: 422-424.