

# Acute coronary syndromes in patients with cancer: Recent advances

Gemina Doolub<sup>1,2</sup>, Aditya S Bharadwaj<sup>3</sup>, Mamas Mamas<sup>1</sup>

<sup>1</sup>Keele Cardiovascular Research Group, Keele University, Keele, United Kingdom

<sup>2</sup>University of Bristol, Bristol, United Kingdom

<sup>3</sup>Loma Linda University, California, United States

## Correspondence to:

Prof. Mamas A Mamas, MD, PhD,  
Keele Cardiovascular  
Research Group,  
Centre for Prognosis Research,  
Institute for Primary Care  
and Health Sciences,  
Keele University, United Kingdom,  
phone: +44 17 82 732 933,  
e-mail:  
masmamas1@yahoo.co.uk

Copyright by the Author(s), 2024

DOI: 10.33963/v.phj.99845

## Received:

February 26, 2024

## Accepted:

March 13, 2024

## Early publication date:

March 18, 2024

## ABSTRACT

Coronary artery disease is presently one of the leading causes of death among cancer survivors. Due to the number of cancer survivors projected to reach 26 million by 2040 managing coronary disease in this population presents a unique challenge. Cancer patients face an elevated risk of atherosclerotic disease due to shared cardiovascular risk factors and the cardiotoxic effects of cancer therapies, predisposing them to acute coronary syndromes. Challenges in treating cancer patients presenting with acute coronary syndromes include atypical presentations, obscured symptoms, and the impact of cancer-related processes on traditional biomarkers. This review explores the complexities of acute coronary syndrome management in cancer patients, addressing challenges involved, recent advances in percutaneous strategies, pharmacology, and considerations for these high-risk individuals. This review discusses a balance between invasive vs. medical strategy, technical advances in multimodal imaging, intravascular physiology, intracoronary imaging, and evolving stent options, highlighting the need for tailored approaches in this complex patient population.

**Key words:** acute coronary syndrome, cancer, cardiovascular disease

## INTRODUCTION


Coronary artery disease (CAD) is presently one of the principal causes of death among cancer survivors [1]. Cardiovascular disease often overtakes cancer as the leading cause of mortality in cancer survivors [2] and certain cancers such as breast, thyroid, endometrial, and prostate cancer are associated with cardiovascular mortality as high as 50% [3].

With rapidly evolving screening and therapeutic innovations, the number of cancer survivors is stabilizing with data from 2022 suggesting that 69% of patients survived their cancer by  $\geq 5$  years from the time of their diagnosis while 47% survived their cancer by  $\geq 10$  years [4]. It is projected that, by 2040, the number of cancer survivors will rise to 26 million, with 74% of that population aged 65 years or more [4].

Cancer patients are at heightened risk of CAD due to a combination of factors (Figure 1). First, cancer populations share overlapping cardiovascular risk factors, such as smoking, obesity, diabetes, and hyperten-

sion [4–7]. Second, certain cancer therapies have been shown to display cardiotoxicity, with some specific treatments predisposing to acute coronary syndromes (ACS) [7]. These specific drugs, and the pathophysiological mechanisms contributing to CAD including ACS, are highlighted in Table 1 [8]. Precipitating factors include accelerated atherosclerosis, plaque rupture (radiation therapy and vascular endothelial growth factor inhibitors); vasospasm (e.g., taxanes and vinca alkaloids); and coronary thrombosis (e.g., alkylating agents such as cisplatin, cyclophosphamide, and platinum-based treatments) [9, 10].

Most data about ACS in cancer patients relies on observational or registry studies, as cancer patients have been excluded from most major randomized CAD trials. These observational studies suggest that ACS patients with underlying cancer are at increased risk of major cardiovascular events and cardiac as well as non-cardiac mortality [9, 11]. In this review, we discuss the particular challenges associated with treating ACS in cancer patients



**Figure 1.** Pathophysiology of cardiovascular disease in cancer

Abbreviation: CVD, cardiovascular disease

**Table 1.** Cancer drugs and their cardiovascular effects [8]

Cardiovascular pathophysiological effect	Cancer drug
Acute thrombosis	Alkylating agents, e.g., cisplatin, cyclophosphamide VEGF inhibitors e.g., bevacizumab
Atherosclerosis acceleration	Tyrosine kinase inhibitors, e.g., nilotinib
Vasospasm	Vinca alkaloids, e.g., vincristine, vinblastine Anti-microtubule agents, e.g., paclitaxel Antimetabolites, e.g., gemcitabine, 5-FU
Endothelial dysfunction	Interferon- $\alpha$

Abbreviations: VEGF, vascular endothelial growth factor; 5FU, 5-fluorouracil

and the recent advances made in percutaneous strategy, and pharmacology, as well as future considerations for treating these high-risk populations.

### CHALLENGES OF TREATING ACS IN CANCER PATIENTS

Treating ACS patients with underlying cancer comes with its own set of challenges. For instance, presentation with ACS can be atypical, with symptoms concealed by cancer itself or treatment-related side effects, and traditional biomarkers sometimes skewed by cancer-related processes [9, 12]. Patients with cancer are also at higher risk of bleeding as well as thrombotic events, such as stroke. Through the stimulation of cytokines, dysregulated platelet activity, endothelial dysfunction, oxidative stress, as well as disorders in coagulation, cancer can lead to pro-inflammatory as well as prothrombotic states [10, 13]. Bleeding can also pose

a significant challenge in this group of patients, and this can be related to local tumor invasion, tumor angiogenesis, oncology therapies, or the systemic effects of the malignancy itself [12, 13]. Thrombocytopenia, which is commonly encountered in certain types of cancer, is associated with worse clinical outcomes, as reported in a study by Yadav et al. [14]. The authors pooled data from two large randomized trials and examined outcomes in 10 603 patients undergoing percutaneous coronary intervention (PCI) for ST-segment elevation myocardial infarction (STEMI) and NSTEMI patients. They found that thrombocytopenia was an independent predictor of 12-month mortality (hazard ratio [HR], 1.74), ischemic target lesion revascularization (HR, 1.37), and major adverse cardiac events (HR, 1.39) [14]. Thus, this is a major consideration in deciding whether to offer interventional options to this particular class of patients, and it requires carefully assessing the associated benefits and risks.

Active malignancy, which is defined as a diagnosis within the previous 12 months or ongoing active cancer therapy including surgery, chemotherapy, or radiotherapy, is considered one of the major criteria for high bleeding risk, as outlined by the Academic Research Consortium for High Bleeding Risk [15, 16]. In a study by Raposeiras-Roubin et al. [17], 1 in 13 post-discharge bleeding events noted in ACS patients was associated with new cancer (positive predictive value for cancer diagnosis of post-discharge bleeding = 7.7%), affecting mainly the gastrointestinal, genitourinary, and bronchopulmonary systems [18].

Moreover, radiation-induced CAD (RICAD), which results from both direct and indirect effects of radiation exposure, is the second most frequent cause of morbidity and mortality among patients exposed to radiotherapy for breast cancer and Hodgkin lymphoma [16]. RICAD has been shown to have a predilection for ostial epicardial coronary lesions, typically involving the left main stem or proximal left anterior descending coronary artery possibly because these vessels lie more anterior/central to the mediastinum in a distribution that is more exposed to radiation [18]. In this group of patients, the relative risk of mortality from myocardial infarction (MI) is roughly double that found in the general population. Treating patients with RICAD can sometimes prove challenging, as the lesions involved can be resistant to treatment, given their fibrotic nature, heavy calcification, and negative remodeling, as demonstrated on intravascular ultrasound [19]. These patients are at high risk for surgical revascularization despite having an indication for it because of concern for bleeding, poor sternal wound healing, and increased morbidity associated with prior chest radiation [20]. PCI in RICAD patients has been shown to be associated with worse outcomes, compared to propensity-match patients, with radiation exposure noted to correlate with higher all-cause mortality [21, 22].

### REVASCULARIZATION FOR ACS IN CANCER PATIENTS

The last few years have seen a shift in the revascularisation strategy adopted in ACS patients with concomitant cancer, moving from a traditionally conservative to a more invasive approach. With the introduction of third-generation drug-eluting stents (DES) and data favoring shorter antiplatelet therapy duration, the new European Society of Cardiology (ESC) guidelines recommend an invasive strategy with coronary angiography and PCI in “patients with cancer presenting with ST-elevation MI (STEMI) or high-risk non-ST elevation ACS (NSTEMI-ACS), with life expectancy  $\geq 6$  months”, or if their ACS is exacerbated by acute complications, such as cardiogenic shock malignant arrhythmias or pulmonary edema [23]. This framework comes as retrospective data [23] suggest better outcomes in cancer patients treated invasively for ACS, compared to the conservative approach. A recent propensity-matched study [24] in STEMI patients with cancer suggested that, despite lower PCI use, the treatment effect of primary PCI was similar to that observed in the no-cancer group. Another study by Balanescu et al. [25], concluded that cancer patients undergoing PCI for acute MI (AMI) had better overall survival rates, compared to patients treated medically, with the most benefit seen when angiography was undertaken within 3 days of admission. However, it is important to note that cancer patients with AMI are a heterogeneous group of patients with varying risk-benefit profiles and clinical outcomes. We have previously demonstrated that AMI patients with lung cancer were associated with the highest in-hospital mortality and MACE rates while those with colon cancer were associated with

the highest risk of bleeding [26]. Additionally, patients with known metastatic disease, who are admitted with an ACS, have been shown to fare worse following PCI, as opposed to having a more conservative strategy [27].

## TECHNICAL ADVANCES

### Multimodality imaging

Since the term MINOCA (myocardial infarction with non-obstructive coronary artery disease syndrome) was first coined in 2013 [28], it has witnessed a growing research interest. MINOCA [29] is defined as the triad of acute MI (positive cardiac biomarker and corroborative clinical evidence of infarction), non-obstructive coronaries on angiography (i.e., no coronary stenosis 50%), and the absence of clear specific cause for the acute presentation. A study by Stepien et al. [30] demonstrated that patients with MINOCA were found to have higher rates of concurrent cancer, compared to patients with MI and obstructive CAD (MI-CAD). Cancer was also noted to correlate with less a favorable survival rate in both groups of patients. Furthermore, takotsubo cardiomyopathy (TC) is common among cancer patients, with a reported prevalence of cancer in TC patients ranging between 6%–28% [31–34]. Multimodal imaging is key in these instances to distinguish a potential presentation of TC from other conditions. Transthoracic echocardiography, for instance, can detect typical appearances of TC, (which include apical ballooning with severe hypokinesia/akinesia of the apical and mid-ventricular segments) although a coronary angiogram or a computed tomography scan is necessary to differentiate it from anterior MI associated with atherosclerotic obstructive CAD. The last decade has seen a dramatic expansion of the use of cardiac magnetic resonance imaging (CMR) in patients with MINOCA, which is a key tool [35] for detecting late gadolinium enhancement, thus localizing the site and pattern of myocardial injury and helping to differentiate between ischemia, myocarditis, and infiltrative processes [36]. CMR can also help stratify risk in patients admitted with MINOCA, with the strongest reported predictors [36] of mortality having been shown to be a CMR diagnosis of cardiomyopathy and ST-segment elevation at the time of presentation. In patients with cancer who present with ambiguous symptoms, non-invasive imaging, such as those described above, is being increasingly used in modern practice to inform diagnosis and management and also avoid invasive procedures or anticoagulants in high-risk patients where imaging can clinch the diagnosis.

### Vascular access

The last decade has seen radial vascular access emerge as the access of choice in both ACS and elective patients undergoing PCI. Radial access [37] has been shown to be associated with lower all-cause mortality and major bleeding, compared with femoral access in numerous randomized controlled trials [38, 39]. In cancer patients

at higher risk of bleeding and vascular complications, this is particularly important, as the radial approach favors prompt ambulation while reducing bleeding risks. Where the radial artery is small in caliber or susceptible to spasm, ultrasound guidance [39], the use of hydrophilic sheaths, and anticoagulation can increase the success rate and reduce complications of radial artery cannulation [40].

### **Intravascular physiology and imaging**

In the last decade, physiological assessment of coronary lesions has emerged as the gold-standard adjunctive tool to conventional coronary angiography in guiding PCI decisions. In cancer patients presenting with ACS and found to have intermediate bystander coronary lesions, fractional flow reserve can be very valuable in identifying hemodynamically significant stenoses in non-culprit vessels [41, 42], thus assisting operators in decision-making on whether those stenotic lesions should be treated with PCI. In recent times, intracoronary imaging has emerged as a critical resource in PCI although there remains considerable geographical and hospital/physician-level variability [43, 44]. In cancer patients, the addition of intravascular imaging can prove extremely valuable in identifying patients with intermediate lesions where acceptable minimum lumen areas can allow for the safe postponement of revascularization [45, 46]. For example, a minimum lumen area of  $\geq 6$  mm was deemed a safe cut-off for deferring revascularization in left-main lesions [47]. This approach can be of particular value in cancer patients with left-main stem disease where the benefits of percutaneous intervention must be balanced with risks conferred by the cancer burden and associated bleeding.

In patients undergoing PCI, intracoronary imaging, such as intravascular ultrasound or optical coherence tomography (OCT), plays a crucial role in defining vessel architecture [48] by detecting and quantifying coronary atheroma, thrombus, and calcium burden [49]. Moreover, intracoronary imaging aids in assessing stent expansion and malapposition, while minimizing periprocedural complications including stent edge dissection and stent thrombosis. Recent consensus position statements from the European Association of Percutaneous Coronary Interventions underscore the pivotal role of imaging in guiding and optimizing stent implantation [49–51]. The PROTECT-OCT Registry [52] showcased the utility of OCT in cancer patients undergoing PCI and allowed operators to identify high-risk patients based on criteria such as uncovered stent struts, stent underexpansion, malapposition, and in-stent restenosis. Consequently, OCT imaging facilitates the identification of cancer patients at low-thrombotic risk, who may safely discontinue dual antiplatelet therapy (DAPT) prematurely to undergo cancer-related surgery [52].


### **Stent options**

The preferred stent strategy when treating cancer patients in the past involved bare metal stents (BMS), to enable

a shorter duration of DAPT. Recent randomized controlled trials have highlighted the superiority of new, third-generation DES [53, 54] over BMS in patient groups at high bleeding risk, especially when long-term DAPT therapy is not a viable option. The new stent platforms, including the polymer-free and carrier-free, umirolimus-coated BioFreedom stent, have been reported, in the LEADERS FREE trial, to outperform BMS with a shortened duration of DAPT (1 month) [55]. Moreover, the ONYX-one study [56] demonstrated that, among patients at significant risk of bleeding, a 1-month DAPT regimen following PCI with zotarolimus-eluting stents was comparable to the use of polymer-free drug-coated stents in terms of safety and adverse outcomes. In addition, the TWILIGHT study [57] demonstrated that among high-risk patients treated with PCI who had completed 3 months of DAPT therapy, ticagrelor monotherapy was associated with a lower incidence of significant bleeding compared to ticagrelor plus aspirin, without an increased risk of death, stroke, or myocardial infarction. Thus, major progress made in treating cancer patients is the feasibility of shorter DAPT therapy with the third-generation stent technology. Finally, drug-eluting balloons (DEB) represent a relatively new technology that enables treatment of in-stent restenotic lesions, but also *de novo* lesions, in small ( $\leq 2.75$  mm) vessels [58, 59] as well as bifurcation lesions [60]. Such a strategy can be particularly useful in cancer patients at exceedingly high bleeding risk, as DAPT following DEB can be stopped after 4 weeks. Studies such as PEPCAD NSTEMI [61] have demonstrated that DCBs for the treatment of *de novo* lesions were non-inferior to BMS or DES although larger trials with DES as a comparator are needed.

### **Antiplatelet therapy and secondary prevention**

In line with the previous discussion, several trials have demonstrated the safety and feasibility of shorter-duration DAPT regimes in cancer patients at high bleeding risk. In addition to the TWILIGHT study [57], the MASTERDAPT subanalysis [62] looked at the effect of 1- or  $\geq 3$ -month DAPT in high bleeding risk (HBR) patients treated with sirolimus stents for complex PCI (defined as one of multivessel PCI;  $\geq 3$  stents/lesions; long stent length, bifurcation disease, etc.). It was found that abbreviated DAPT was associated with lower bleeding complications compared with standard DAPT (HR, 0.64; 95% CI, 0.42–0.9), while not being associated with significantly increased risks of ischemic events [62]. The STOPDAPT-2 trial [63], which compared 1 month of DAPT followed by clopidogrel monotherapy to standard 12-month DAPT, showed the shortened DAPT regime to meet criteria for both non-inferiority and superiority, with significantly reduced rates of cardiovascular and bleeding events [58]. In light of these findings, current ESC guidelines [64] recommend considering an abbreviated DAPT regime (1-month DAPT) in patients at high bleeding risk (class IIb) and de-escalation of P2Y<sub>12</sub> in-



**Central figure.** Acute coronary syndrome and cancer advances

Abbreviations: ACS, acute coronary syndrome; BMS, bare metal stents; CMR, cardiac magnetic resonance; CT, computed tomography; DAPT, dual antiplatelet therapy; FFR, fractional flow reserve; ISR, in-stent restenotic lesions; IVUS, intravascular ultrasound; MINOCA, myocardial infarction with nonobstructive coronary arteries; OCT, optical coherence tomography; SCAD, spontaneous coronary artery dissection

hibitors (e.g., from ticagrelor to clopidogrel) to reduce bleeding propensity. In high-risk cancer patients, these recommendations, together with a careful assessment of thrombotic vs. bleeding risks, are very useful when deciding about the length of the DAPT regime. While undertaking such risk assessments, it will be essential to first define HBR patients, as per the Academic Consortium Consensus document [16]. According to this document, HBR is defined as Bleeding Academic Research Consortium, BARC 3 or 5 bleeding risk of  $\geq 4\%$  at 1 year or a risk of intracranial hemorrhage of  $\geq 1\%$  at 1 year. The consensus document also defines 20 clinical criteria, which are further divided into major criteria (e.g., long-term oral anticoagulation, severe renal failure with  $\text{eGFR} < 30 \text{ ml/min/1.73 m}^2$ , anemia with  $\text{Hb} < 11 \text{ g/dl}$ , active malignancy, liver cirrhosis with portal hypertension); and minor criteria (e.g., age  $\geq 75$  years, long-term use of non-steroidal anti-inflammatory drugs) [16].

Furthermore, secondary prevention has as important a role in cancer patients as in non-cancer patients. Thus, unless there are specific contraindications, consideration should be made regarding starting patients on drugs with an established impact on survival post-ACS, such as angiotensin-converting enzyme inhibitors, beta-blockers, and statins. In addition, lifestyle modification, wherever possible, should also be encouraged, in the form of a healthy diet, smoking cessation, and increased physical activity.

## CONCLUSION

In conclusion, the intricate interplay between CAD and cancer poses substantial challenges in the management of cancer survivors. With the increasing prevalence of both conditions, there is a pressing need for nuanced approaches to address the unique considerations of ACS in this population (Central Figure). As emphasized by a previous expert opinion by Leszek et al. [65], the interplay between CAD and oncology remains very nuanced and extremely important when considering therapeutic options in this category of patients. For this reason, a multidisciplinary approach, including cardiology and oncology, or a specialized cardio-oncologist if available, together with radiologists, surgeons, and, if required, gastroenterologists, should help not only in selecting treatment based on comorbidity and risk evaluation but also in monitoring for any potential complications related to treatment. The development of new stent technology, as well as innovations in intravascular imaging and pharmacology, have shown promise in improving outcomes for cancer patients presenting with ACS. Post-ACS care of these patients must involve collaborative efforts between cardiologists and oncologists to develop patient-specific strategies that optimize cardiovascular outcomes without compromising cancer treatment.

### Article information

**Conflict of interest:** None declared.

**Funding:** None.

**Open access:** This article is available in open access under Creative Commons Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, which allows downloading and sharing articles with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially. For commercial use, please contact the journal office at [polishheartjournal@ptkardio.pl](mailto:polishheartjournal@ptkardio.pl)

## REFERENCES

- Mamas MA, Brown SA, Sun LY. Coronary artery disease in patients with cancer: It's always the small pieces that make the bigger picture. *Mayo Clin Proc.* 2020; 95(9): 1819–1821, doi: [10.1016/j.mayocp.2020.07.006](https://doi.org/10.1016/j.mayocp.2020.07.006), indexed in Pubmed: 32861320.
- Wang Z, Fan Z, Yang L, et al. Higher risk of cardiovascular mortality than cancer mortality among long-term cancer survivors. *Front Cardiovasc Med.* 2023; 10: 1014400, doi: [10.3389/fcvm.2023.1014400](https://doi.org/10.3389/fcvm.2023.1014400), indexed in Pubmed: 36760569.
- Cancer society. Cancer Treatment & Survivorship Facts & Figures 2019–2021; 2019.
- Statistics and graphs. Office of Cancer survivorship. National Cancer institute: Division of cancer control & population sciences. Nov 2022. <https://cancercontrol.cancer.gov/ocs/statistics#stats> (accessed: January 11, 2024).
- Koene RJ, Prizment AE, Blaes A, et al. Shared risk factors in cardiovascular disease and cancer. *Circulation.* 2016; 133(11): 1104–1114, doi: [10.1161/CIRCULATIONAHA.115.020406](https://doi.org/10.1161/CIRCULATIONAHA.115.020406), indexed in Pubmed: 26976915.
- Wolin KY, Carson K, Colditz GA. Obesity and cancer. *Oncologist.* 2010; 15(6): 556–565, doi: [10.1634/theoncologist.2009-0285](https://doi.org/10.1634/theoncologist.2009-0285), indexed in Pubmed: 20507889.
- Giovannucci E, Harlan DM, Archer MC, et al. Diabetes and cancer: a consensus report. *Diabetes Care.* 2010; 33(7): 1674–1685, doi: [10.2337/dc10-0666](https://doi.org/10.2337/dc10-0666), indexed in Pubmed: 20587728.
- Costa IB, Andrade FT, Carter D, et al. Challenges and management of acute coronary syndrome in cancer patients. *Front Cardiovasc Med.* 2021; 8: 590016, doi: [10.3389/fcvm.2021.590016](https://doi.org/10.3389/fcvm.2021.590016), indexed in Pubmed: 34179121.
- Grossman E, Messerli FH, Boyko V, et al. Is there an association between hypertension and cancer mortality? *Am J Med.* 2002; 112(6): 479–486, doi: [10.1016/s0002-9343\(02\)01049-5](https://doi.org/10.1016/s0002-9343(02)01049-5), indexed in Pubmed: 11959059.
- Lyon AR, Dent S, Stanway S, et al. Baseline cardiovascular risk assessment in cancer patients scheduled to receive cardiotoxic cancer therapies: a position statement and new risk assessment tools from the Cardio-Oncology Study Group of the Heart Failure Association of the European Society of Cardiology in collaboration with the International Cardio-Oncology Society. *Eur J Heart Fail.* 2020; 22(11): 1945–1960, doi: [10.1002/ehfj.1920](https://doi.org/10.1002/ehfj.1920), indexed in Pubmed: 32463967.
- Caine GJ, Stonelake PS, Lip GH, et al. The hypercoagulable state of malignancy: pathogenesis and current debate. *Neoplasia.* 2002; 4(6): 465–473, doi: [10.1038/sj.neo.7900263](https://doi.org/10.1038/sj.neo.7900263), indexed in Pubmed: 12407439.
- Potts JE, Iliescu CA, Lopez Mattei JC, et al. Percutaneous coronary intervention in cancer patients: a report of the prevalence and outcomes in the United States. *Eur Heart J.* 2019; 40(22): 1790–1800, doi: [10.1093/eurheartj/ehy769](https://doi.org/10.1093/eurheartj/ehy769), indexed in Pubmed: 30500952.
- Doolub G, Mamas MA. Percutaneous coronary angioplasty in patients with cancer: Clinical challenges and management strategies. *J Pers Med.* 2022; 12(9): 1372, doi: [10.3390/jpm12091372](https://doi.org/10.3390/jpm12091372), indexed in Pubmed: 36143156.
- Yadav M, Génèreux P, Giustino G, et al. Effect of baseline thrombocytopenia on ischemic outcomes in patients with acute coronary syndromes who undergo percutaneous coronary intervention. *Can J Cardiol.* 2016; 32(2): 226–233, doi: [10.1016/j.cjca.2015.05.020](https://doi.org/10.1016/j.cjca.2015.05.020), indexed in Pubmed: 26341303.
- Johnstone C, Rich SE. Bleeding in cancer patients and its treatment: A review. *Ann Palliat Med.* 2018; 7(2): 265–273, doi: [10.21037/apm.2017.11.01](https://doi.org/10.21037/apm.2017.11.01), indexed in Pubmed: 29307210.
- Urban P, Mehran R, Collieran R, et al. Defining high bleeding risk in patients undergoing percutaneous coronary intervention: a consensus document from the Academic Research Consortium for High Bleeding Risk. *Eur Heart J.* 2019; 40(31): 2632–2653, doi: [10.1093/eurheartj/ehz372](https://doi.org/10.1093/eurheartj/ehz372), indexed in Pubmed: 31116395.
- Raposeiras-Roubin S, Abu-Assi E, Muñoz-Pousa I, et al. Usefulness of bleeding after acute coronary syndromes for unmasking silent cancer. *Am J Cardiol.* 2020; 125(12): 1801–1808, doi: [10.1016/j.amjcard.2020.03.023](https://doi.org/10.1016/j.amjcard.2020.03.023), indexed in Pubmed: 32307091.
- Kirresh A, White L, Mitchell A, et al. Radiation-induced coronary artery disease: a difficult clinical conundrum. *Clin Med (Lond).* 2022; 22(3): 251–256, doi: [10.7861/clinmed.2021-0600](https://doi.org/10.7861/clinmed.2021-0600), indexed in Pubmed: 35584837.
- Borges N, Kapadia S. Radiation-induced CAD: incidence, diagnosis, and management outcomes. *American College of Cardiology.* 2018. <https://www.acc.org/Latest-in-Cardiology/Articles/2018/05/24/01/44/Radiation-Induced-CAD> (accessed: January 11, 2024).
- Cuomo JR, Javaheri SP, Sharma GK, et al. How to prevent and manage radiation-induced coronary artery disease. *Heart.* 2018; 104(20): 1647–1653, doi: [10.1136/heartjnl-2017-312123](https://doi.org/10.1136/heartjnl-2017-312123), indexed in Pubmed: 29764968.
- Wu W, Masri A, Popovic ZB, et al. Long-term survival of patients with radiation heart disease undergoing cardiac surgery: a cohort study. *Circulation.* 2013; 127(14): 1476–1485, doi: [10.1161/CIRCULATIONAHA.113.001435](https://doi.org/10.1161/CIRCULATIONAHA.113.001435), indexed in Pubmed: 23569119.
- Reed GW, Rossi JE, Masri A, et al. Long-Term mortality in patients with radiation-associated coronary artery disease treated with percutaneous coronary intervention. *Circ Cardiovasc Interv.* 2016; 9(6): e003483, doi: [10.1161/CIRCINTERVENTIONS.115.003483](https://doi.org/10.1161/CIRCINTERVENTIONS.115.003483), indexed in Pubmed: 27313281.
- Lyon AR, López-Fernández T, Couch LS, et al. 2022 ESC Guidelines on cardio-oncology developed in collaboration with the European Hematology Association (EHA), the European Society for Therapeutic Radiology and Oncology (ESTRO) and the International Cardio-Oncology Society (IC-OS). *Eur Heart J.* 2022; 43(41): 4229–4361, doi: [10.1093/eurheartj/ehac244](https://doi.org/10.1093/eurheartj/ehac244), indexed in Pubmed: 36017568.
- Mohamed MO, Van Spall HGC, Kontopantelis E, et al. Effect of primary percutaneous coronary intervention on in-hospital outcomes among active cancer patients presenting with ST-elevation myocardial infarction: a propensity score matching analysis. *Eur Heart J Acute Cardiovasc Care.* 2021; 10(8): 829–839, doi: [10.1093/ehjacc/zuaa032](https://doi.org/10.1093/ehjacc/zuaa032), indexed in Pubmed: 33587752.
- Balanescu DV, Donisan T, Deswal A, et al. Acute myocardial infarction in a high-risk cancer population: Outcomes following conservative versus invasive management. *Int J Cardiol.* 2020; 313: 1–8, doi: [10.1016/j.ijcard.2020.04.050](https://doi.org/10.1016/j.ijcard.2020.04.050), indexed in Pubmed: 32320781.
- Bharadwaj A, Potts J, Mohamed MO, et al. Acute myocardial infarction treatments and outcomes in 6.5 million patients with a current or historical diagnosis of cancer in the USA. *Eur Heart J.* 2020; 41(23): 2183–2193, doi: [10.1093/eurheartj/ehz851](https://doi.org/10.1093/eurheartj/ehz851), indexed in Pubmed: 31800032.
- Guddati AK, Joy PS, Kumar G. Analysis of outcomes of percutaneous coronary intervention in metastatic cancer patients with acute coronary syndrome over a 10-year period. *J Cancer Res Clin Oncol.* 2016; 142(2): 471–479, doi: [10.1007/s00432-015-2056-5](https://doi.org/10.1007/s00432-015-2056-5), indexed in Pubmed: 26498773.
- Beltrame JF. Assessing patients with myocardial infarction and nonobstructed coronary arteries (MINOCA). *J Intern Med.* 2013; 273(2): 182–185, doi: [10.1111/j.1365-2796.2012.02591.x](https://doi.org/10.1111/j.1365-2796.2012.02591.x), indexed in Pubmed: 22998397.
- Agewall S, Beltrame JF, Reynolds HR, et al. ESC working group position paper on myocardial infarction with non-obstructive coronary arteries. *Eur Heart J.* 2017; 38(3): 143–153, doi: [10.1093/eurheartj/ehw149](https://doi.org/10.1093/eurheartj/ehw149), indexed in Pubmed: 28158518.
- Stepien K, Nowak K, Szlosarczyk B, et al. Clinical characteristics and long-term outcomes of MINOCA accompanied by active cancer: A retrospective insight into a Cardio-Oncology Center Registry. *Front Cardiovasc Med.* 2022; 9: 785246, doi: [10.3389/fcvm.2022.785246](https://doi.org/10.3389/fcvm.2022.785246), indexed in Pubmed: 35669480.
- Zaghlol R, Dey A, Barac A. Takotsubo and cancer: Takotsubo cardiomyopathy in the era of emerging cancer therapies. *Eur Heart J.* 2020; 41(48): 4547–4549, doi: [10.1093/eurheartj/ehaa175](https://doi.org/10.1093/eurheartj/ehaa175).
- Sattler K, El-Battrawy I, Gietzen T, et al. Prevalence of cancer in Takotsubo cardiomyopathy: Short and long-term outcome. *Int J Cardiol.* 2017; 238(7): 159–165, doi: [10.1016/j.ijcard.2017.02.093](https://doi.org/10.1016/j.ijcard.2017.02.093), indexed in Pubmed: 28318661.
- Zaghlol R, Kashyap K, Al-Shbool G, et al. Usefulness of malignancy as a predictor of worse in-hospital outcomes in patients with takotsubo cardiomyopathy. *Am J Cardiol.* 2019; 123(6): 995–1001, doi: [10.1016/j.amjcard.2018.11.054](https://doi.org/10.1016/j.amjcard.2018.11.054), indexed in Pubmed: 30595393.
- El-Battrawy I, Santoro F, Stiermaier T, et al. Prevalence and long-term prognostic impact of malignancy in patients with Takotsubo syndrome. *Eur J Heart Fail.* 2018; 20(4): 816–818, doi: [10.1002/ehfj.868](https://doi.org/10.1002/ehfj.868), indexed in Pubmed: 28849607.

35. Agewall S, Beltrame JF, Reynolds HR, et al. ESC working group position paper on myocardial infarction with non-obstructive coronary arteries. *Eur Heart J*. 2017; 38(3): 143–153, doi: [10.1093/eurheartj/ehw149](https://doi.org/10.1093/eurheartj/ehw149), indexed in Pubmed: 28158518.
36. Dastidar AG, Baritussio A, De Garate E, et al. Prognostic role of CMR and conventional risk factors in myocardial infarction with nonobstructed coronary arteries. *JACC Cardiovasc Imaging*. 2019; 12(10): 1973–1982, doi: [10.1016/j.jcmg.2018.12.023](https://doi.org/10.1016/j.jcmg.2018.12.023), indexed in Pubmed: 30772224.
37. Ferrante G, Rao SV, Jüni P, et al. Radial versus femoral access for coronary interventions across the entire spectrum of patients with coronary artery disease: a meta-analysis of randomized trials. *JACC Cardiovasc Interv*. 2016; 9(14): 1419–1434, doi: [10.1016/j.jcin.2016.04.014](https://doi.org/10.1016/j.jcin.2016.04.014), indexed in Pubmed: 27372195.
38. Gargiulo G, Giacoppo D, Jolly SS, et al. Effects on mortality and major bleeding of radial versus femoral artery access for coronary angiography or percutaneous coronary intervention: meta-analysis of individual patient data from 7 multicenter randomized clinical trials. *Circulation*. 2022; 146(18): 1329–1343, doi: [10.1161/CIRCULATIONAHA.122.061527](https://doi.org/10.1161/CIRCULATIONAHA.122.061527), indexed in Pubmed: 36036610.
39. Valgimigli M, Saia F, Guastaroba P, et al. Transradial versus transfemoral intervention for acute myocardial infarction: a propensity score-adjusted and -matched analysis from the REAL (Registro regionale Angioplastiche dell'Emilia-Romagna) multicenter registry. *JACC Cardiovasc Interv*. 2012; 5(1): 23–35, doi: [10.1016/j.jcin.2011.08.018](https://doi.org/10.1016/j.jcin.2011.08.018), indexed in Pubmed: 22230147.
40. Seto AH, Roberts JS, Abu-Fadel MS, et al. Real-time ultrasound guidance facilitates transradial access: RAUST (Radial Artery access with Ultrasound Trial). *JACC Cardiovasc Interv*. 2015; 8(2): 283–291, doi: [10.1016/j.jcin.2014.05.036](https://doi.org/10.1016/j.jcin.2014.05.036), indexed in Pubmed: 25596790.
41. Munoz-Gonzalez, E. et al. PCI in patients with cancer. *Cardiac Interventions Today*. <https://citoday.com/articles/2019-jan-feb/pci-in-patient-s-with-cancer> (accessed: February 25, 2024).
42. Mehta SR, Wood DA, Storey RF, et al. Complete revascularization with multivessel PCI for myocardial infarction. *N Engl J Med*. 2019; 381(15): 1411–1421, doi: [10.1056/nejmoa1907775](https://doi.org/10.1056/nejmoa1907775), indexed in Pubmed: 31475795.
43. Kim JW, Dayah TJ, Javadi A, et al. Reclassification of treatment strategy with fractional flow reserve in cancer patients with coronary artery disease. *Medicina (Kaunas)*. 2022; 58(7): 884, doi: [10.3390/medicina58070884](https://doi.org/10.3390/medicina58070884), indexed in Pubmed: 35888603.
44. Donisan T, Dayah T, Balanescu D, et al. Clinical outcomes after fractional flow reserve-guided treatment of oncology patients. *J Clin Oncol*. 2018; 36(Suppl 15): e22106, doi: [10.1200/jco.2018.36.15\\_suppl.e22106](https://doi.org/10.1200/jco.2018.36.15_suppl.e22106).
45. Madder RD, Seth M, Sukul D, et al. Rates of intracoronary imaging optimization in contemporary percutaneous coronary intervention: a report from the BMC2 registry. *Circ Cardiovasc Interv*. 2022; 15(10): e012182, doi: [10.1161/CIRCINTERVENTIONS.122.012182](https://doi.org/10.1161/CIRCINTERVENTIONS.122.012182), indexed in Pubmed: 36256694.
46. Munoz-Gonzalez E, Poulin MF, Goel M, et al. PCI in patients with cancer: Approaches for successfully treating this challenging patient population when invasive coronary procedures are needed. *Cardiac Interventions Today*. 2019.
47. de la Torre Hernandez JM, Hernandez FH, Alfonso F, et al. Prospective application of pre-defined intravascular ultrasound criteria for assessment of intermediate left main coronary artery lesions. *J Am Coll Cardiol*. 2011; 58(4): 351–358, doi: [10.1016/j.jacc.2011.02.064](https://doi.org/10.1016/j.jacc.2011.02.064), indexed in Pubmed: 21757111.
48. Richards G, Johnson T. A vision of percutaneous coronary revascularisation in 2021: How to take advantage of intra-coronary imaging to perform more effective PCI. *JRSM Cardiovasc Dis*. 2021; 10: 20480040211049978, doi: [10.1177/20480040211049978](https://doi.org/10.1177/20480040211049978), indexed in Pubmed: 35186282.
49. Räber L, Mintz GS, Koskinas KC, et al. Clinical use of intracoronary imaging. Part 1: guidance and optimization of coronary interventions. An expert consensus document of the European Association of Percutaneous Cardiovascular Interventions. *Eur Heart J*. 2018; 39(35): 3281–3300, doi: [10.1093/eurheartj/ehy285](https://doi.org/10.1093/eurheartj/ehy285), indexed in Pubmed: 29790954.
50. Johnson TW, Räber L, Di Mario C, et al. Clinical use of intracoronary imaging. Part 2: acute coronary syndromes, ambiguous coronary angiography findings, and guiding interventional decision-making: an expert consensus document of the European Association of Percutaneous Cardiovascular Interventions. *Eur Heart J*. 2019; 40(31): 2566–2584, doi: [10.1093/eurheartj/ehz332](https://doi.org/10.1093/eurheartj/ehz332), indexed in Pubmed: 31112213.
51. Lotfi A, Jeremias A, Fearon WF, et al. Expert consensus statement on the use of fractional flow reserve, intravascular ultrasound, and optical coherence tomography: a consensus statement of the Society of Cardiovascular Angiography and Interventions. *Catheter Cardiovasc Interv*. 2014; 83(4): 509–518, doi: [10.1002/ccd.25222](https://doi.org/10.1002/ccd.25222), indexed in Pubmed: 24227282.
52. Iliescu CA, Cilingiroglu M, Giza DE, et al. „Bringing on the light” in a complex clinical scenario: Optical coherence tomography-guided discontinuation of antiplatelet therapy in cancer patients with coronary artery disease (PROTECT-OCT registry). *Am Heart J*. 2017; 194: 83–91, doi: [10.1016/j.ahj.2017.08.015](https://doi.org/10.1016/j.ahj.2017.08.015), indexed in Pubmed: 29223438.
53. Valgimigli M, Patialiakas A, Thury A, et al. Zotarolimus-eluting versus bare-metal stents in uncertain drug-eluting stent candidates. *J Am Coll Cardiol*. 2015; 65(8): 805–815, doi: [10.1016/j.jacc.2014.11.053](https://doi.org/10.1016/j.jacc.2014.11.053).
54. Ariotti S, Adamo M, Costa F, et al. Is bare-metal stent implantation still justifiable in high bleeding risk patients undergoing percutaneous coronary intervention?: a pre-specified analysis from the ZEUS trial. *JACC Cardiovasc Interv*. 2016; 9(5): 426–436, doi: [10.1016/j.jcin.2015.11.015](https://doi.org/10.1016/j.jcin.2015.11.015), indexed in Pubmed: 26965932.
55. Urban P, Meredith IT, Abizaid A, et al. Polymer-free drug-coated coronary stents in patients at high bleeding risk. *N Engl J Med*. 2015; 373(21): 2038–2047, doi: [10.1056/NEJMoa1503943](https://doi.org/10.1056/NEJMoa1503943), indexed in Pubmed: 26466021.
56. Windecker S, Latib A, Kedhi E, et al. Polymer-based or polymer-free stents in patients at high bleeding risk. *N Engl J Med*. 2020; 382(13): 1208–1218, doi: [10.1056/nejmoa1910021](https://doi.org/10.1056/nejmoa1910021), indexed in Pubmed: 32050061.
57. Mehran R, Baber U, Sharma SK, et al. Ticagrelor with or without aspirin in high-risk patients after PCI. *N Engl J Med*. 2019; 381(21): 2032–2042, doi: [10.1056/NEJMoa1908419](https://doi.org/10.1056/NEJMoa1908419), indexed in Pubmed: 31556978.
58. Jeger RV, Eccleshall S, Wan Ahmad WA, et al. Drug-coated balloons for coronary artery disease: Third report of the international DBC consensus group. *JACC Cardiovasc Interv*. 2020; 13(12): 1391–1402, doi: [10.1016/j.jcin.2020.02.043](https://doi.org/10.1016/j.jcin.2020.02.043), indexed in Pubmed: 32473887.
59. Latib A, Colombo A, Castriota F, et al. A randomized multicenter study comparing a paclitaxel drug-eluting balloon with a paclitaxel-eluting stent in small coronary vessels: the BELLO (Balloon Elution and Late Loss Optimization) study. *J Am Coll Cardiol*. 2012; 60(24): 2473–2480, doi: [10.1016/j.jacc.2012.09.020](https://doi.org/10.1016/j.jacc.2012.09.020), indexed in Pubmed: 23158530.
60. Mathey DG, Wendig I, Boxberger M, et al. Treatment of bifurcation lesions with a drug-eluting balloon: the PEPCAD V (Paclitaxel Eluting PTCA Balloon in Coronary Artery Disease) trial. *EuroIntervention*. 2011; 7 Suppl K: K61–K65, doi: [10.4244/EIJV7SKA11](https://doi.org/10.4244/EIJV7SKA11), indexed in Pubmed: 22027730.
61. Scheller B, Ohlow MA, Ewen S, et al. Bare metal or drug-eluting stent versus drug-coated balloon in non-ST-elevation myocardial infarction: the randomised PEPCAD NSTEMI trial. *EuroIntervention*. 2020; 15(17): 1527–1533, doi: [10.4244/EIJ-D-19-00723](https://doi.org/10.4244/EIJ-D-19-00723), indexed in Pubmed: 31659986.
62. Valgimigli M, Smits PC, Frigoli E, et al. Duration of antiplatelet therapy after complex percutaneous coronary intervention in patients at high bleeding risk: a MASTER DAPT trial sub-analysis. *Eur Heart J*. 2022; 43(33): 3100–3114, doi: [10.1093/eurheartj/ehac284](https://doi.org/10.1093/eurheartj/ehac284), indexed in Pubmed: 35580836.
63. Watanabe H, Domei T, Morimoto T, et al. Effect of 1-month dual antiplatelet therapy followed by clopidogrel vs 12-month dual antiplatelet therapy on cardiovascular and bleeding events in patients receiving PCI: the STOPDAPT-2 randomized clinical trial. *JAMA*. 2019; 321(24): 2414–2427, doi: [10.1001/jama.2019.8145](https://doi.org/10.1001/jama.2019.8145), indexed in Pubmed: 31237644.
64. Byrne RA, Rossello X, Coughlan JJ, et al. 2023 ESC Guidelines for the management of acute coronary syndromes: developed by the task force on the management of acute coronary syndromes of the European Society of Cardiology (ESC). *Eur Heart J*. 2023; 44(38): 3720–3826, doi: [10.1093/eurheartj/ehad191](https://doi.org/10.1093/eurheartj/ehad191), indexed in Pubmed: 37622654.
65. Leszek P, Klotzka A, Bartuś S, et al. A practical approach to the 2022 ESC cardio-oncology guidelines: Comments by a team of experts - cardiologists and oncologists. *Kardiol Pol*. 2023; 81(10): 1047–1063, doi: [10.33963/v.kp.96840](https://doi.org/10.33963/v.kp.96840), indexed in Pubmed: 37660389.