

## **Supplementary material**

---

*Urbanowicz T, Wojtasińska E, Olasińska-Wiśniewska A, et al. Neutrophil to extracellular traps (NET) as an early marker of right ventricular dilatation in patients with left ventricular assist device (LVAD). Pol Heart J. 2024.*

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

**Table S1.** Demographical and clinical characteristics of the presented groups. Group 1 (no changes in RV diameter) and group 2 (RV dilatation)

Parameters	Group 1 (no changes in RV diameter) n=18	Group 2 (RV dilatation) n = 7	P-value
Demographical			
Age, years (median [Q1–Q3])	61 (56–63)	51 (39–60)	0.10
BMI (median [Q1–Q3])	28.5 (27.6–28.7)	27.8 (25.7–29.9)	1.00
Clinical			
Ischemic cardiomyopathy, n (%)	10 (56)	3 (43)	0.60
Dilated cardiomyopathy, n (%)	5 (28)	4 (57)	0.27
Pulmonary hypertension, n (%)	12 (67)	4 (57)	0.69
Arterial hypertension, n (%)	5 (28)	3 (43)	0.50
Atrial fibrillation, n (%)	6 (33)	4 (57)	0.30
Diabetes mellitus, n (%)	6 (33)	1 (14)	0.37
ICD implantation, n (%)	16 (89)	6 (86)	0.87

Abbreviations: BMI, body mass index; ICD, implantable cardioverter-defibrillator; RV, right ventricle

**Table S2.** Echocardiographic results collected preoperatively and during follow-up in both subgroups

Parameters	Group 1 (no changes in RV diameter) n = 18	Group 2 (RV dilatation) n = 7	P-value
Pre-implantation echocardiographic results			
LVEDd, mm (median [Q1–Q3])	66 (60–71)	70 (64–78)	0.24
RVEDd, mm (median [Q1–Q3])	34 (32–37)	35 (25–37)	0.61
IVS, mm (median [Q1–Q3])	10 (9–10)	11 (10–12)	0.26
LAD, mm (median [Q1–Q3])	47 (44–49)	48 (43–56)	0.88
LVEF, % (median [Q1–Q3])	18 (15–22)	23 (15–28)	0.47
Mitral valve regurgitation, grade (median [Q1–Q3])	1.0 (1.0–1.0)	1. (1.0–1.8)	0.46
Tricuspid valve regurgitation, grade (median [Q1–Q3])	1.0 (1.0–1.8)	1.0 (1.0–1.0)	0.57
TAPSE, mm (median [Q1–Q3])	15 (14–16)	14 (13–15)	0.06
Follow-up — echocardiographic results			
LVEDd, mm (median [Q1–Q3])	66 (63–71)	69 (53–75)	0.98
RVEDd, mm (median [Q1–Q3])	31 (28–32)	40 (35–41)	0.01 <sup>a</sup>
RV diameter change, mm (median [Q1–Q3])	-3 (-5 to -2)	3 (3–6)	<0.001 <sup>a</sup>
IVS, mm (median [Q1–Q3])	10 (9–11)	11 (10–11)	0.46
LAD, mm (median [Q1–Q3])	48 (48–50)	46 (40–50)	0.47
LVEF, % (median [Q1–Q3])	20 (20–21)	20 (15–24)	0.72
Mitral valve regurgitation, grade (median [Q1–Q3])	1.0 (1.0–1.8)	1.0 (1.0–1.0)	0.88
Tricuspid valve regurgitation, grade (median [Q1–Q3])	1.0 (1.0–1.0)	1.0 (1.0–1.0)	0.87
TAPSE, mm (median [Q1–Q3])	14 (14–16)	14 (14–15)	0.61

<sup>a</sup>Statistically significant

Abbreviations: IVS, intraventricular septum; LAD, left atrium diameter; LVEDd, left ventricular end diastolic diameter; LVEF, left ventricular ejection fraction; RV, right ventricle; RVEDd, right ventricular end-diastolic diameter; TAPSE, tricuspid annular systolic excursion

**Table S3.** LVAD characteristics, laboratory results in the follow-up and changes in citH3 plasma concentrations in both subgroups

Parameters	Group 1 (no changes in RV diameter) n = 18	Group 2 (RV dilatation) n = 7	P-value
LVAD function parameters			
Speed (median [Q1–Q3])	5200 (5125– 5200)	5000 (4888– 5125)	0.32
PI (median [Q1–Q3])	2.1 (1.9–2.3)	2.1 (2.0–2.3)	1.00
Generated cardiac index, mm (median [Q1–Q3])	4.350 (4.225– 4.475)	4.000 (3.900– 4.150)	0.09
Whole blood count analysis			
WBC, K/ul (median [Q1–Q3])	7.74 (7.34–8.11)	8.68 (6.79–9.76)	0.77
Neutrophil, K/ul (median [Q1–Q3])	5.56 (4.73–6.51)	4.80 (4.76–8.07)	0.86
Monocyte, K/ul (median [Q1–Q3])	0.58 (0.46–0.62)	0.45 (0.43–0.51)	0.21
Lymphocyte, K/ul (median [Q1–Q3])	1.18 (0.95–1.71)	1.35 (1.25–1.36)	0.59
LUC, K/ul (median [Q1–Q3])	0.13 (0.12–0.22)	0.17 (0.16–0.20)	0.39
Hb, mmol/l (median [Q1–Q3])	8.40 (7.15–9.45)	8.30 (6.30–8.80)	0.54
Hematocrit, % (median [Q1–Q3])	40.0 (35.3–46.0)	42.0 (31.0–43.0)	0.71
Platelets, K/ul (median [Q1–Q3])	200 (158–271)	178 (166–199)	0.68
MCV, fl (median [Q1–Q3])	88.0 (86.3–90.8)	95 (89.0–97.0)	0.33
MCHC, g/dl (median [Q1–Q3])	20.5 (20.3–20.9)	20.5 (20.0–21.0)	0.77
Biochemical laboratory tests			
LDH, U/mg (median [Q1–Q3])	211 (203–232)	176 (168–196)	0.14
NT pro-BNP, pg/ml (median [Q1– Q3])	1299 (921–2157)	598 (556–719)	0.13
CRP, mg/l (median [Q1–Q3])	3 (2–5)	4 (1–7)	0.88
ALT, IU/l (median [Q1–Q3])	37 (28–42)	18 (16–23)	0.09
AST, IU/l (median [Q1–Q3])	26 (23–35)	21 (21–29)	0.36
citH3 plasma concentrations			
Preoperatively, pg/ml (median [Q1– Q3])	2255 (1533– 2375)	2633 (2288– 2699)	0.19

In the follow-up, pg/ml (median [Q1–Q3])	637 (501–770)	1935 (1099–3396)	0.003*
--	---------------	------------------	--------

Abbreviations: ALT, alanine aminotransferase; AST, aspartate aminotransferase; citH3, citrullinated histone 3; CRP, C-reactive protein; LDH, lactate dehydrogenase; LUC, large unstained cell count; LVAD, left ventricular assist device; MCHC, mean corpuscular hemoglobin concentration; MCV, mean corpuscular volume; PI, pulsation index; WBC, white blood cell count

**Table S4.** Correlations between citH3 and echocardiographic parameters

Preoperative correlations					
LVEDd	RVEDd	IVS	LAD	LVEF	
r = 0.64 <i>P</i> = 0.06	r = 0.35 <i>P</i> = 0.35	r = 0.27 <i>P</i> = 0.38	r = 0.55 <i>P</i> = 0.13	r = 0.71 <i>P</i> = 0.03*	
Postoperative correlations					
LVEDd	RVEDd	RVED change	IVS	LAD	LVEF
r = 0.013 <i>P</i> = 0.95	r = 0.26 <i>P</i> = 0.22	r = 0.47 <i>P</i> = 0.01*	r = 0.28 <i>P</i> = 0.36	r = 0.08 <i>P</i> = 0.98	r = 0.20 <i>P</i> = 0.48

Abbreviations: IVS, interventricular septum; LAD, left atrium diameter; LVEDd, left ventricular end diastolic diameter; LVEF, left ventricular ejection fraction; RVEDd, right ventricular end diastolic diameter