

Supplementary material

Oleksiak A, Kępka C, Rucińska K, et al. Iron level and vulnerable coronary atherosclerotic plaques. Pol Heart J. 2024.

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

Table S1. Analysis of the predictors of low-attenuation plaque.

Parameter	Univariable ^a		Multivariable	
	OR (95% CI)	P-value	OR (95% CI)	P-value
Male sex	2.50 (1.40–4.46)	0.002	2.41 (1.24–4.70)	0.009
Iron	1.02 (1.00–1.03)	<0.0001	1.02 (1.01–1.03)	0.0002
Hemoglobin	1.23 (1.01–1.49)	0.04	0.89 (0.70–1.14)	0.4
RBC	1.69 (0.97–2.97)	0.07		
hsCRP	0.84 (0.55–1.28)	0.2		
Age	1.00 (0.99–1.01)	0.5		
BMI	1.00 (0.94–1.06)	1.0		
Hypertension	1.35 (0.82–2.24)	0.1		
Dyslipidemia ^b	1.73 (0.92–3.24)	0.08		
Diabetes	1.19 (0.66–2.13)	0.6		

Family history of CAD	1.69 (0.88–3.22)	0.1
Smoking — current	1.09 (0.53–2.22)	0.8
Smoking — in the past	1.40 (0.71–2.76)	0.3

^aData presented for parameters significantly associated or with a trend in univariable analysis.

^bDyslipidemia diagnosis in the medical records at the day of CTCA

Abbreviations: BMI, body mass index; CAD, coronary artery disease; CI, confidence interval; hsCRP, high-sensitivity C-reactive protein; OR, odds ratio; RBC, red blood count