

Supplementary material

Warmiński G, Sadowski KA, Kalińczuk Ł, et al. Artificial intelligence analysis of ECG signals to predict arrhythmia recurrence after ablation of atrial fibrillation. Pol Heart J. 2025.

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

Table S1. Baseline demographic and clinical characteristics

	All (n = 250, 100%)	Arrhythmia recurrence (n = 108, 43.2%)	Preserved sinus rhythm (n = 142, 56.8%)	P-value
Age, years, median (IQR)	61.0 (49.0–67.3)	60.0 (47.25–68.0)	61.5 (53.8–67.0)	0.64
BMI kg/m ² , median (IQR)	27.5(24.5–30.0)	27.8 (25.6–30.3)	27.3 (25.1–29.7)	0.30
Female, n (%)	75 (30)	33 (30.6)	42 (29.6)	0.89
Paroxysmal AF, n (%)	151 (60.4)	51 (47.2)	100 (70.4)	<0.001
Valvular heart disease, n (%)	12 (4.8)	9 (8.3)	3 (2.1)	0.03
Cardiomyopathy, n (%)	43 (17.2)	20 (18.5)	23 (16.2)	0.74
Hypertension, n (%)	169 (67.6)	76 (70.4)	93 (65.5)	0.50
Diabetes, n (%)	21 (8.4)	6 (5.6%)	15 (10.6)	0,18
LVEF%, median (IQR)	60.0 (55.0–65.0)	61.0 (58.3–65.0)	60.0 (55.0–65.0)	0.32
Increased left ventricular wall thickness, n (%)	141 (56.4)	69 (63.9)	72 (50.7)	0.040
Left atrial volume indexed, cm ³ /m ² , median (IQR)	50.2 (40.1–64.6)	58.4 (43.4–71.5)	45.1 (37.7–59.1)	<0.001

Abbreviations: AF, atrial fibrillation; BMI, body mass index; IQR, interquartile range; LVEF, left ventricular ejection fraction