

Supplementary material

Iwańczyk S, Borger M, Kamiński M, et al. Inflammatory response in patients with coronary artery ectasia and coronary artery disease. Kardiol Pol. 2019; 77: 713-715.
doi:10.33963/KP.14812

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

Table S1 Results of the study group

Parameter	CAE group (n = 27)	CAD group (n = 27)	Control group (n = 27)	P value		
				CAE vs CAD	CAE vs control	CAD vs control
MPV, fl	8.6 (7.9- 9.7)	8.8 (8.3- 9.6)	7.8 (7.3- 8.7)	0.52	0.02	0.004
Creatinine, mmol/l	92.3 (73.6- 99.3)	91.4 (82.3- 98.7)	76.7 (68.0- 87.2)	0.97	0.02	0.01
hs-CRP, mg/l	2.2 (1.25- 4.0)	3.2 (1.9- 3.2)	1 (0.8-1.6)	0.77	0.03	0.08
HDL-C, mmol/l	1.3 (1.2- 1.7)	1.3 (1.2- 1.7)	1.7 (1.4- 2.1)	1	0.001	0.001
TG/HDL-C ratio	0.93 (0.72- 1.72)	1.04 (0.59- 1.75)	0.62 (0.42- 1.00)	0.79	0.01	0.05

LDL-C/HDL-C ratio	2.03 (1.40-2.30)	1.30 (1.00-1.97)	1.15 (0.89-2.05)	0.03	0.06	0.81
Monocytes/HDL-L-C ratio	0.39 (0.25-0.49)	0.33 (0.24-0.39)	0.26 (0.18-0.34)	0.23	0.01	0.05
Endocan, ng/l	621 (552-1339)	617 (566-760)	591 (483-1434)	0.44	0.5	0.82

Data are presented as median (interquartile range).

Abbreviations: CAD, coronary artery disease; CAE, coronary artery ectasia; HDL-C, high-density lipoprotein cholesterol; hs-CRP, high-sensitivity C-reactive protein; LDL-C, low-density lipoprotein cholesterol; MPV, mean platelet volume; TG, triglycerides