

## **Supplementary material**

---

*Sławiński G, Dorniak K, Lewicka E. Implantable cardioverter-defibrillators in cardiac amyloidosis: a grey zone requiring an individual approach. Kardiol Pol. 2020; 78: 934-936. doi:10.33963/KP.15435*

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

**Supplementary video 1.** Magnetic resonance imaging. 2-chamber routine balanced steady state free precession cine sequence, showing myocardial thickening and reduced left ventricular long axis systolic foreshortening. (Siemens Aera 1.5 T, Erlangen, Germany).

**Supplementary video 2.** Magnetic resonance imaging. 4-chamber routine balanced steady state free precession cine sequence, showing myocardial thickening and reduced left ventricular long axis systolic foreshortening. (Siemens Aera 1.5 T, Erlangen, Germany).

**Supplementary video 3.** Transthoracic echocardiography: left atrial enlargement, sparkling myocardial appearance, preserved left ventricular ejection fraction (apical 2-chamber view) (Vivid S70, GE Healthcare, Chicago, Illinois, USA).

**Supplementary video 4.** Transthoracic echocardiography: left atrial enlargement, biventricular wall thickening with sparkling myocardial appearance, preserved left ventricular ejection fraction (apical 4-chamber view) (Vivid S70, GE Healthcare, Chicago, Illinois, USA).