

Supplementary material

Wiliński J, Skwarek A, Borek R, et al. Subcostal echocardiographic assessment of tricuspid annular kick (SEATAK) — a novel independent predictor of 30-day mortality in patients with acute pulmonary embolism. Kardiol Pol. 2022.

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

Table S1. Clinical characteristics of excluded patients

	Included patients	Excluded patients	
n	164	19	
PE	82 (50%)	PE 7 (36.84%) No PE 8 (42.11%) No data 4 (21.05%)	
Gender [male]	75 (45.73%)	8 (42.11%)	0.76
Age [years]	70 (59.75–80)	75 (67 - 81.5)	0.26
BMI [kg/m ²]	27.36 ± 5.95	26.46 ± 4.09	0.48
Arterial hypertension	59 (35.98%)	4 (21.05%)	0.31
Hyperlipidemia	43 (26.22%)	2 (10.53%)	0.17
Diabetes mellitus	19 (11.59%)	1 (5.26%)	0.70
CAD	31 (18.9%)	3 (15.79%)	1
CHF	29 (17.68%)	2 (10.53%)	0.75
AF (present or prior)	16 (9.76%)	2 (10.53%)	1
History of stroke	3 (1.83%)	0 (0%)	1
Chronic lung disease	11 (6.71%)	0 (0%)	0.61
Active malignancy	35 (21.34%)	5 (26.32%)	0.79
Acute infection	25 (15.24%)	1 (5.26%)	0.32
HR [bpm]	82 (70–100)	81 (67–91)	0.59
SBP [mm Hg]	126 (114–138)	120 (100.5–128)	0.15
DBP [mm Hg]	75 (70–81)	73 (61–75)	0.16
Wells rule - original version [points]	3 (1.5–4.5)	4.5 (3.38–4.5)	0.49
Revised Geneva rule – original version [points]	5 (4–6)	4.5 (4–5.75)	0.75
PESI	94 (80–118)	97 (88.5–135.5)	0.22
sPESI	1 (0–2)	1.5 (1 – 2)	0.37
EF %	55 (46–63)	53 (45–66)	0.97
RVTD [mm]	39 (36–43)	42 (36–46)	0.39
LVEDd [mm]	46 (41–52)	43.5 (40–51)	0.28