

Supplementary material

Oleksiak A, Kepka C, Rucińska K, et al. HDL cholesterol, triglycerides and characteristics of coronary atherosclerosis in patients with newly diagnosed significant coronary artery disease by CTCA. Kardiol Pol. 2022.

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.

Table S1. The lipids tests results in the study population

Parameter (mg/dl)	Value (n = 300)
Total cholesterol	156.0 (132.0–188.0)
LDL-C	90.0 (68.0–119.5)
HDL-C	51.0 (44.0–62.0)
Triglycerides	113.0 (85.0–162.0)

Values are median (IQR)

Abbreviations: LDL-C, low-density lipoprotein cholesterol; HDL-C, high-density lipoprotein cholesterol

Table S2. The laboratory tests results according to statin treatment categories

Parameter (mg/dl)	Statin-treated (n = 120)	Statin-untreated (n = 180)	P-value
Total cholesterol	153 (129–179)	165 (140–198)	0.008
LDL-C	86 (66–108)	102 (72–131)	0.007
HDL-C	51 (44–61)	54 (44–64)	0.4
Triglycerides	112 (90–167)	111 (82–157)	0.3

Values are median (IQR)

Abbreviations: LDL-C, low-density lipoprotein cholesterol; HDL-C, high-density lipoprotein cholesterol