

Supplementary material

Kaczmarek D, Matuszewska-Brycht O, Kobierecki M, et al. Prediction of major adverse cardiac events in patients with multivessel coronary artery disease with an interpretable classification tree model based on coronary computed tomography angiography. Kardiol Pol. 2023.

Please note that the journal is not responsible for the scientific accuracy or functionality of any supplementary material submitted by the authors. Any queries (except missing content) should be directed to the corresponding author of the article.




Figure S1. Assessing the optimal tree depth using 5-fold cross validation

Table S1. Characteristics of the first study group — clinical and demographic data

Age, mean (SD)	69.42 (8.28)
Gender	
Men, n (%)	74 (69.8)
Women, n (%)	32 (30.2)
BMI, kg/m ² , mean (SD)	27.91 (4.44)
CCS scale	

I, n (%)	16 (15.1)
II, n (%)	47 (44.3)
III, n (%)	40 (37.8)
IV, n (%)	3 (2.8)
NYHA scale	
I, n (%)	13 (12.3)
II, n (%)	75 (70.7)
III, n (%)	17 (16.1)
IV, n (%)	1 (0.9)
HF, n (%)	
HFpEF, n (% of HF)	56 (52.8)
HFmrEF, n (% of HF)	43 (76.7)
HFrEF, n (% of HF)	8 (14.3)
	5 (9.0)
Smoking	
Never, n (%)	48 (45.3)
In the past, n (%)	38 (35.8)
Current, n (%)	20 (18.9)
Hypertension, n (%)	
	98 (92.5)
Diabetes mellitus	
Present diabetes mellitus, n (%)	38 (35.8)
Impaired fasting glucose, n (%)	3 (2.8)
Impaired glucose tolerance, n (%)	2 (2.0)
Alcohol abuse, n (%)	
	1 (0.9)
Dyslipidemia, n (%)	
	106 (100)
CKD, n (%)	
	20 (18.9)
COPD, n (%)	
	6 (5.7)

Abbreviations: BMI, body mass index; CCS, Canadian Cardiovascular Society angina grade; CKD, chronic kidney disease defined as $GFR < 60 \text{ ml/min/1.73m}^2$; COPD, chronic obstructive pulmonary disease; HF, heart failure; HFmrEF, HF with mildly reduced ejection fraction; HFpEF, HF with

preserved ejection fraction; HFrEF, HF with reduced ejection fraction; n, number of patients; NYHA, New York Heart Association Functional Classification; SD, standard deviation

Table S2. Characteristics of the first study group — biochemical parameters

Parameter	Median (1 st quartile–3 rd quartile)
Morphology	
RBC, mln/ μ l	4.51 (4.2–4.86)
WBC, thousand/ μ l	7.32 (6.09–8.52)
Hgb, g/dl	14.0 (12.8–14.78)
PLT, thousand/ μ l	213.0 (182.25–252.0)
Lipid profile	
Total cholesterol, mmol/l	4.15 (3.71–4.95)
HDL, mmol/l	1.21 (1.05–1.52)
LDL, mmol/l	2.26 (1.8–3.02)
TG, mmol/l	1.22 (0.93–1.85)
Others	
Sodium, mmol/l	139.7 (138.02–140.78)
Potassium, mmol/l	4.32 (4.12–4.56)
eGFR, ml/min./1.73 m ²	77.8 (62.32–89.78)
Creatine, μ mol/l	81.75 (72.12–97.58)
TSH, μ IU/ml	1.22 (0.73–2.1)
TnT, ng/L	12.5 (9.0–17.0)

CK-MB mass, ng/ml	2.8 (2.2–3.78)
NT-proBNP, pg/ml	250.0 (100.0–753.8)
Uric acid, $\mu\text{mol/l}$	360.5 (298.48–438.78)

Abbreviations: CK-MB, creatine kinase-myoglobin binding; eGFR, estimated glomerular filtration rate; HDL, high-density lipoprotein; Hgb, hemoglobin; LDL, low-density lipoprotein; NT-proBNP, N-terminal pro-B-type natriuretic peptide; PLT, platelet count; RBC, red blood cells; TG, triglycerides; TnT, troponin T; TSH, thyroid-stimulating hormone; WBC, white blood cells

Table S3. Factors affecting MACE — univariate logistic regression

	Odds ratio	–95% CI	+95% CI	<i>P</i> -value
Male gender	2.08	0.64	6.79	0.22
Age	1.42	0.88	2.29	0.14
Hyperglycemia	2.13	0.71	6.37	0.17
LDL level	0.89	0.54	1.46	0.65
Elevated LDL	0.71	0.17	2.89	0.63
HDL level	1.24	0.77	1.99	0.36
Decreased HDL	1.39	0.36	5.30	0.62
Total cholesterol	0.84	0.50	1.41	0.52
Elevated total cholesterol	0.95	0.31	2.91	0.93
TG level	0.49	0.24	0.99	0.049
Elevated TG level	0.21	0.04	0.98	0.048
Positive family history of CVD	0.69	0.24	1.98	0.49
Arterial hypertension	0.91	0.58	1.4286	0.70

Impaired fasting glucose	8.84	0.76	102.62	0.08
Type 2 DM treated with oral medicines	0.71	0.21	2.37	0.58
Type 2 DM treated with insulin	1.25	0.31	5.01	0.75
Presence of heart failure	3.6	1.20	10.71	0.02
Reduced EF	2.87	0.44	18.43	0.26
Mildly reduced EF	3.09	0.78	12.18	0.10
Preserved EF	0.29	0.09	0.95	0.04
BMI level	0.55	0.32	0.94	0.03
Overweight or obesity (BMI >25)	0.56	0.21	1.55	0.27
CRP	1.11	0.71	1.74	0.61
WBC	1.20	0.75	1.92	0.43
RBC	0.71	0.43	1.17	0.18
Hemoglobin	0.72	0.45	1.17	0.19
PLT	1.01	0.63	1.63	0.94
Creatinine	2.81	1.23	6.37	0.01
eGFR	0.55	0.35	0.89	0.01
TSH	1.36	0.82	2.24	0.22
TnT	4.52	1.81	11.28	0.001
CKMB-mass	1.48	0.94	2.30	0.08
NT-proBNP	1.77	1.13	2.76	0.01

Uric acid	1.42	0.88	2.28	0.14
Current smoking	0.66	0.17	2.52	0.55
History of COVID-19	3.37	1.04	10.90	0.04
History of MI	6.85	2.12	22.13	0.001
History of ischemic stroke	2.62	0.68	9.97	0.15
CKD	2.76	0.93	8.17	0.07
Peripheral thrombosis	2.21	0.72	6.76	0.16
COPD	4.55	0.84	24.43	0.07
Calcium score	1.01	0.63	1.62	0.95
LM significant narrowing in CCTA	16.22	5.10	51.57	<0.001
LAD significant narrowing in CCTA	3.6	1.20	10.71	0.02
LCx significant narrowing in CCTA	1.14	0.39	3.31	0.80
RCA significant narrowing in CCTA	1.24	0.47	3.27	0.66

Abbreviations: BMI, body mass index; CCTA, coronary computed tomography angiography; CI, confidence interval; CKD, chronic kidney disease; CK-MB, creatine kinase-myoglobin binding; COPD, chronic obstructive pulmonary disease; CRP, C-reactive protein; CVD, cardiovascular disease; DM, diabetes mellitus; EF, ejection fraction; eGFR, estimated glomerular filtration rate; HDL, high-density lipoprotein; LAD, left anterior descending artery; LCx, left circumflex coronary artery; LDL, low-density lipoprotein; LM, left main coronary artery; MI, myocardial infarction; NT-proBNP, N-terminal pro-B-type natriuretic peptide; PLT, platelet count; RBC, red blood cells; RCA, right coronary artery; TG, triglycerides; TnT, troponin T; TSH, thyroid-stimulating hormone; WBC, white blood cells