



Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

journal homepage: <http://www.elsevier.com/locate/rpor>



## Case report

# Delayed reconstruction of the upper digestive tract in a patient following total pharyngolaryngectomy with resection of the cervical oesophagus

Mateusz Szewczyk<sup>a,\*</sup>, Jakub Pazdrowski<sup>a</sup>, Paweł Golusiński<sup>a,b</sup>, Wojciech Golusiński<sup>a</sup>

<sup>a</sup> Department of Head and Neck Surgery, Poznan University of Medical Sciences Greater Poland Cancer Centre, Garbary 15, 61-866 Poznan, Poland

<sup>b</sup> Department of Biology and Environmental Studies, Poznan University of Medical Sciences, Dluga 1/2, 61-848 Poznan, Poland



### ARTICLE INFO

#### Article history:

Received 23 April 2014

Received in revised form

27 August 2014

Accepted 20 November 2014

#### Keywords:

Reconstruction

Pharyngolaryngectomy

Free flap

Head and Neck

Delayed

### ABSTRACT

Carcinoma of the hypopharynx is an uncommon disease, with an annual incidence of approximately 1 in 100,000. Post-cricoid carcinoma is more common in women and is not usually associated with tobacco and alcohol abuse. Reconstruction of large pharyngeal defects following surgery for squamous cell carcinoma is complex and often requires microvascular free tissue transfer to achieve the best oncological and functional outcomes. The most common complications of such procedures include fistulas and strictures of the neopharynx. Here, we describe a case of a female patient admitted to the Head and Neck Department at our hospital to undergo delayed reconstruction following pharyngolaryngectomy and removal of the cervical oesophagus. Several complications occurred during post-operative care, including stricture and skin dehiscence. At present, the patient is able to swallow saliva and is currently being prepared to return to a normal diet.

© 2014 Greater Poland Cancer Centre. Published by Elsevier Sp. z o.o. All rights reserved.

## 1. Introduction

The incidence of hypopharyngeal cancer is estimated at 1 in 100,000 and it accounts for 5% of all head and neck malignancies.<sup>1</sup> The pyriform sinus is involved in 70% of cases, the post-cricoid region in 20%, and the posterior wall of the pharynx in 10–15%.<sup>2,3</sup> Post-cricoid cancer is most common in Northern Europe, North America, India and Japan.<sup>4,5</sup> Unlike all

other head and neck tumours, which occur more frequently in men, post-cricoid cancer is more common in females and is not usually associated with tobacco and alcohol abuse.<sup>6,7</sup> Regional metastases occur in 20–40% of cases.<sup>8,9</sup>

Due to nonspecific symptoms, hypopharyngeal cancer is typically diagnosed at late stages, with invasion of surrounding anatomical structures and high risk of distant metastases. Early stage tumours are usually treated with radiochemotherapy.<sup>10</sup> Advanced tumours that have invaded

\* Corresponding author. Tel.: +48 609540393.

E-mail address: [mateuszszewczyk@yahoo.pl](mailto:mateuszszewczyk@yahoo.pl) (M. Szewczyk).

<http://dx.doi.org/10.1016/j.rpor.2014.11.009>

1507-1367/© 2014 Greater Poland Cancer Centre. Published by Elsevier Sp. z o.o. All rights reserved.

surrounding tissues and recurrent tumours both require extensive surgical procedures with one-step reconstruction. It is important to resect hypopharyngeal cancer with a 3 cm surgical margin; as a result, total pharyngolaryngectomy with cervical oesophagectomy is often necessary.<sup>11</sup>

Reconstruction of the upper aerodigestive tract after oncological resection is one of the most challenging and difficult procedures in head and neck surgery. The options for hypopharyngeal reconstruction have evolved over time, starting from multistep reconstruction, which had a high failure rate, to one-step reconstructions in which the functional outcome is better. Several factors are crucial in successful hypopharynx reconstruction: easy access to donor tissue with minimal donor site morbidity, a low rate of fistulas and strictures of the neopharynx, and good tolerance to post-operative radiotherapy.<sup>12</sup>

To date, many hypopharyngeal reconstructive techniques have been introduced, including the use of different types of flaps (local, myocutaneous, free fasciocutaneous, and jejunal flaps), and gastric pull-up. Nevertheless, no consensus has been reached on which technique is most efficient in certain defects.

The pectoralis major myocutaneous flap is a reliable flap with excellent vascularisation; it has the added benefit of providing muscular covering for the carotid artery after neck dissection. However, the flap is too big to be rolled into a tube-like shape to cover the circumferential defect, especially in women and obese patients. Moreover, published data indicate that a high percentage of fistulas occur with this flap and females are sometimes unwilling to accept the donor site scarring that occurs.<sup>13</sup> The main use of the pectoralis major flap is in salvage surgery after a free flap failure and in elderly patients with other comorbidities that preclude them from multi-hour procedures.<sup>14</sup>

The free jejunal flap is often used in circumferential defects of the hypopharynx and the rate of post-operative fistulas is low. In addition, due to segmental vascular supply, a 20 cm long flap on one vascular arcade is possible. On the other hand, patients often suffer from dysphagia after jejunal reconstruction due to uncontrolled peristalsis.<sup>15</sup> Additionally, the reconstruction makes the patient's voice less powerful and "wet-sounding". Finally, jejunal flap grafting requires laparotomy, which is associated with additional donor site morbidity.<sup>12</sup>

Fasciocutaneous free flaps are thin and pliable, and can thus be shaped into a tubular form. When Spriano modification is used, the flap can be sutured to prevertebral fascia, thus providing additional lumen to the neopharynx. No donor site morbidities have been observed. Two team work shortens the total time needed to perform the surgery, and the long vascular pedicle facilitates reconstruction. Moreover, the flap's good vascular supply improves tolerance to ischaemia during vascular reconstruction. When using the radial forearm free flap, the Allen test must be performed to assess the deep palmar arterial arch.<sup>16</sup> When anterolateral thigh flaps are used, ultrasound examination of the thigh must be performed to evaluate the vascular perforators.<sup>17</sup> However, fasciocutaneous flaps present a greater risk of post-operative fistulas and strictures, especially in delayed procedures.<sup>18</sup> Moreover, patients with previous neck dissection may have existing ligations of

the external carotid artery, its branches, and internal jugular vein, which limits vascular.<sup>19</sup>

## 2. Case presentation

A 39-year old female patient was admitted to the Head and Neck Department at our institution. Four months prior to admission to our centre, the patient had been diagnosed at another hospital with squamous cell carcinoma of the post-cricoid region, with circumferential infiltration to the hypopharynx and cervical oesophagus. The patient had undergone a pharyngolaryngectomy, with the resection of the cervical oesophagus and bilateral radical neck dissection at the other institution, with a final histopathological report of pT4bNOM0. Postoperative margins were free from tumour. The cephalic end of the patient's oesophagus had been transferred to the left supraclavicular region (Fig. 1). The patient had not been irradiated.

In order to determine if reconstruction was feasible, the patient underwent computed tomography and magnetic resonance imaging scans to assess the defect and to rule out any potential local and regional recurrence. The defect of the upper digestive tract was 7 cm. After thoroughly discussing the case with the patient, the decision was made to reconstruct the neopharynx with a radial forearm free flap (due to the lack of thigh vascular perforators).

After dissecting the hypopharyngeal oesophageal stumps, the radial forearm free flap was removed from the patient's left forearm, shaped into an inverted U-tube and sutured to the previously dissected prevertebral fascia, which created the posterior wall of the neopharynx (Figs. 2 and 3). The free flap was stabilised with a previously inserted Montgomery tube with a feeding probe inside it (Fig. 4). Next, vascular neck anastomosis was performed with the vascular pedicle sutured to facial vessels with 9.0 sutures. The skin was sutured in layers. No perioperative complications were observed.

Thirty days after surgery the Montgomery tube was removed and barium-swallow X-ray examination was performed (Fig. 5). The radiographic examination revealed that the neopharynx was sealed and the patient was allowed to swallow clear liquids. On day 40 after surgery a partial dehis-




**Fig. 1 – The cephalic end of the patient's oesophagus had been transferred to the left supraclavicular region.**




**Fig. 2 – The operating field after dissecting the hypopharyngeal oesophageal stumps.**


cence of the wound occurred with local infection and stricture of the neopharynx (Fig. 6). A gastroscopy was performed and an oesophageal stent was inserted into the neopharynx during the procedure: simultaneously, percutaneous gastrostomy was performed (Fig. 7). The stent was removed six weeks later. Fourteen weeks after reconstruction the patient is able to swallow saliva and is being fed through the gastrostomy. An oral liquid diet will be initiated in 4 weeks. No donor site morbidities on the patient's forearm have been observed.




**Fig. 3 – Radial forearm free flap was removed from the patient's left forearm, shaped into an inverted U-tube and sutured to the previously dissected prevertebral fascia.**



**Fig. 4 – Previously inserted Montgomery tube with a feeding probe inside it.**



**Fig. 5 – Barium-swallow X-ray examination 30 days after the surgery.**



**Fig. 6 – Forty days after surgery a partial dehiscence of the wound occurred with local infection and stricture of the neopharynx.**

### 3. Discussion

Patients with hypopharyngeal cancer typically have nonspecific early symptoms, leading to a delay in seeking treatment. This explains why these patients often require extensive resections that often involve the larynx, hypopharynx, and cervical oesophagus. Following resection, patients usually undergo one-time reconstruction, whose main goal is to enable swallowing. However, surgical procedures involving reconstruction of the hypopharynx and cervical oesophagus are extremely challenging for head and neck surgeons.



**Fig. 7 – Oesophageal stent was inserted into the neopharynx during gastroscopy.**

Mura et al.<sup>20</sup> evaluated long-term reconstruction results in patients after pharyngolaryngectomy and found that free flaps were significantly better than other flaps. In that study, none of the 27 patients who underwent pedicled flap reconstruction returned to a normal diet, whereas 80% of patients with free flap reconstruction were able to do so. In our case, the lack of adequate vascular perforators on the patient's thigh, together with the potential morbidities of laparotomy, led the surgeon to choose radial forearm free flap as the donor tissue for the hypopharyngeal reconstruction. The main limitation of this technique is a relatively high percentage of fistulas, which range from 25 to 35%.<sup>21–23</sup> It is believed that the main cause of post-operative fistulas in this procedure is the use of accessory vertical sutures, which are used to suture the flap to the pre-vertebral fascia.<sup>24</sup> Another frequent complication, which also occurred in our patient, is neopharyngeal stricture (5–40% of all cases).<sup>25</sup> However, the use of a Montgomery tube can help prevent strictures and facilitate reconstruction. In contrast to Murray et al.,<sup>26</sup> we allowed our patient to consume a liquid diet after tube removal. A similar solution has been proposed by Lopez et al.<sup>27</sup> who examined 55 cases of hypopharyngeal reconstruction in which strictures and fistulas developed in 5% and 9% of patients, respectively.

#### 4. Conclusion

The advantages of fasciocutaneous free flaps in hypopharyngeal reconstruction are mainly the reliability and pliability of the flap, the diameter of the vascular pedicle, and the low donor site morbidity. The main complications are fistulas and strictures, which may delay the ability of the patient to return to a normal diet. However, these complications can be minimised by using the Montgomery tube, which allows the surgeon to create a wide lumen in the neopharynx and to protect the sutures from saliva produced by the patient in the early weeks after surgery. We believe that the choice of a reconstruction technique depends on the characteristics of the defect. Finally, it is important to consider the preferences

of both the surgeon and, especially, the patient when selecting a specific approach.

---

#### Conflict of interest

None declared.

---

#### Financial disclosure

None declared.

#### REFERENCES

---

- Patel R, Goldstein D, Brown D, et al. Circumferential pharyngeal reconstruction: history, critical analysis of techniques, and current therapeutic recommendations. *Head Neck* 2010;32:109–20.
- Bahadur S, Thakar A, Mohanti BK, et al. Results of radiotherapy with, or without, salvage surgery versus combined surgery and radiotherapy in advanced carcinoma of the hypopharynx. *J Laryngol Otol* 2002;116:29–32.
- Godballe C, Jorgensen K, Hansen O, et al. Hypopharyngeal cancer: results of treatment based on radiation therapy and salvage surgery. *Laryngoscope* 2002;112:834–8.
- Hatta C, Satoh T, oshino Y, Fujii K, Sawada T. Clinical study of postcricoid carcinoma. *Nippon Jibinkoka Gakkai Kaiho* 1991;94:495–9.
- Harrison DFN. Pathology of hypopharyngeal cancer in relation to surgical management. *J Laryngol Otol* 1970;84:349–67.
- Jones AS, Mcrae RD, Phillips DE, Hamilton J, Field JK, Husband D. The treatment of node negative squamous cell carcinoma of the postcricoid region. *J Laryngol Otol* 1995;109(February):114–9.
- Archibald S, Young JEM. A pharyngo-cervical esophageal reconstruction. *Clin Plast Surg* 2005;32:339–46.
- Stell PM, Maran AGD. Head and neck surgery. 2nd ed. London: Heinemann Medical Books; 1978. p. 169–77.
- Lefebvre JL, Castelain B, De la Torre JD, Delobelle-Deroide A, Vankemmel B. Lymph node invasion in hypopharynx and lateral epilarynx carcinoma: a prognostic factor. *Head Neck Surg* 1987;10:14.
- Jones AS. The management of early hypopharyngeal cancer: primary radiotherapy and salvage surgery. *Clin Otolaryngol* 1992;17:545–9.
- Ho CM, Lam KH, Wei WI, et al. Squamous cell carcinoma of the hypopharynx – analysis of treatment results. *Head Neck* 1993;15:405–12.
- Couch ME. Laryngopharyngectomy with reconstruction. *Otolaryngol Clin North Am* 2002;35:1097–114.
- Murakami Y, Saito S, Ikari T, Haraguchi S, Okada K, Maruyama T. Esophageal reconstruction with a skin-grafted pectoralis major muscle flap. *Arch Otolaryngol* 1982;108:719–22.
- Chu P-Y, Chang S-Y. Reconstruction of the hypopharynx after surgical treatment of squamous cell carcinoma. *J Chin Med Assoc* 2009;72(7):351–5.
- Reece GP, Schusterman MA, Miller MJ, et al. Morbidity and functional outcome of free jejunal transfer reconstruction for circumferential defects of the pharynx and cervical esophagus. *Plast Reconstr Surg* 1995;96:1307–16.
- Khan AS, Henton JM, Adams TS, et al. Evaluation of hand circulation before radial forearm free flap surgery. *Laryngoscope* 2009;119:1679–81.

17. Golusinski P, Luczewski L, Pazdrowski J, et al. The role of colour duplex sonography in preoperative perforator mapping of the anterolateral thigh flap. *Eur Arch Otorhinolaryngol* 2013;July) [Epub ahead of print].
18. Welkoborska H-J, Deichmüllera C, Bauera L, Hinni ML. Reconstruction of large pharyngeal defects with microvascular free flaps and myocutaneous pedicled flaps. *Curr Opin Otolaryngol Head Neck Surg* 2013;21:318–27, <http://dx.doi.org/10.1097/MOO.0b013e3283631ea2>.
19. Wong KK, Higgins KM, Enepekides DJ. Microvascular reconstruction in the vessel depleted neck. *Curr Opin Otolaryngol Head Neck Surg* 2010;18:223–6.
20. Mura F, Bertino G, Occhini A, et al. Advanced carcinoma of the hypopharynx: functional results after circumferential pharyngolaryngectomy with flap reconstruction. *Acta Otorhinolaryngol Ital* 2012;32:154–215.
21. Su CY, Chiang YC. The fabricated radial forearm flap in pharyngolaryngeal surgery: saliva leakage and its prevention. *Br J Surg* 1995;48:212–7.
22. Cho BC, Kim M, Lee JH, Byun JS, Park JS, Baik BS. Pharyngoesophageal reconstruction with a tubed free radial forearm. *J Reconstr Microsurg* 1998;14:535–40.
23. Anthony JP, Singer MI, Deschler DG, Dougherty ET, Reed CG, Kaplan MJ. Long-term functional results after pharyngoesophageal reconstruction with the radial forearm free flap. *Am J Surg* 1994;168:441–5.
24. Azizzadeh B, Yafai S, Rawnsley JD, et al. Radial forearm free flap pharyngoesophageal reconstruction. *Laryngoscope* 2001;111:807–10.
25. Murray DJ, Novak CB, Neligan PC. Fasciocutaneous free flaps in pharyngolaryngoesophageal reconstruction: a critical review of the literature. *J Plast Reconstr Aesthet Surg* 2008;61:1148–56.
26. Murray DJ, Gilbert RW, Vesely MJ, et al. Functional outcomes and donor site morbidity following circumferential pharyngoesophageal reconstruction using an anterolateral thigh flap and salivary bypass tube. *Head Neck* 2007;29: 147–54.
27. Lopez F, Obeso S, Camporro D, et al. Outcomes following pharyngolaryngectomy with fasciocutaneous free flap reconstruction and salivary bypass tube. *Laryngoscope* 2013;123:591–6.