

Supplementary File

Table S1. Tumor control after repeat SRS for vestibular schwannoma (n = 6)

Series author, year, and location	No. of patients	Median follow-up after repeat SRS	Tumor control after Repeat SRS	
			2-year	5-year
Dewan et al, 2008, Rhode Island, USA ³	11	Not reported	77%	77%
Yomo et al, 2009, Marseille, France ⁴	8	5.3 years	100%	100%
Lonneville et al, 2015, Brussels, Belgium ⁵	27	3.8 years (mean)	Not reported	85%
Fu et al, 2017, Tilburg, The Netherlands ⁶	38	6.3 years	100%	100%
Iorio-Morin et al, 2019, International ⁷	76	4.3 years	99%	92%
Rapp et al, 2020, Gainesville, FL, USA	6	8.4 years	84%	84%

Table S2. Literature review of toxicity following repeat SRS for vestibular schwannoma

Series Author, Year, and Location	No. of Patients	Neurological Complications from Repeat SRS		
		Trigeminal Nerve	Facial Nerve	Brainstem Necrosis
Dewan et al, 2008, Rhode Island, USA ³	11	2 (18%)	0	0
Yomo et al, 2009, Marseille, France ⁴	8	0	0	0
Lonneville et al, 2015, Brussels, Belgium ⁵	27	0	0	0
Fu et al, 2017, Tilburg, The Netherlands ⁶	38	2 (5%)	3 (8%)	0
Iorio-Morin et al, 2019, International ⁷	76	6 (8%)	5 (7%)	0
Rapp et al, 2020, Florida, USA	6	1 (17%)	0	0