Supplementary File

GROUP n = 12	21		
Sex	Female	n = 95	79.30%
	Male	n = 26	20.70%
Age	Mean		51.77±11.40
Education	Primary	n = 1	0.9%
	Vocational	n = 25	20.7%
	Secondary	n = 57	47.2%
	Tertiary	n = 38	32.2%
Residence	Rural	n = 38	31.4%
	Urban	n = 83	68.6%
Occupation	Not employed	n = 19	15.8%
•	Pensioner	n = 28	23.1%
	Disability pension	n = 13	10.7%
	Different forms of occupation	n = 61	50.4%
Marital status	Not in a relation	n = 14	12%
	Married	n = 91	75%
	In an informal relation	n = 16	13%

Table S1	. The s	sociodem	ographic	description	n of the	group	in the study	
								_

Table S2. Criteria for inclusion or exclusion from the participation in the study

NCLUSION CRITERIA CHARACTERISTICS OF THE CRITERION						
Breast and colorectal cancer						
Cancer at an earlier stage of development						
In the course of neoadjuvant and adjuvant therapy (radical						
treatment						
18 to 85 years of age						
Conscious consent to participate in the study						
No diagnosed mental disorders						
CHARACTERISTICS OF THE CRITERION						
Other cancers apart from the breast or colorectal cancer						
Cancer at an advanced stage of development						
In the course of a paliative chemiotherapy						
No consent given to participate in the study						
Diagnosed mental disorder						

Table S3. Existence of an experience of a chronic, serious disease in oneself or in a close family member before contracting cancer vis a vis perception of one's illness, the coping strategy adopted, acceptance of the situation and the mental state of the patient

Experience of a previous se	rious dise	ease						
Average rank	Group I N=27	Group II <i>N</i> =10	Group III N=75	Grou p IV <i>N</i> =9	Inter- group. compari son	H Kruskala– Wallisa (df = 3)	p	ε ²
Perception of the disease:								
Duration of the disease	66.63	56.80	59.89	58.06		0.99	0.81	-
Consequences of the disease	64.44	69.25	57.71	68.89		1.95	0.58	-
Freq. of the occur. of sympt.	69.22	49.35	59.45	62.22		2.80	0.42	—
Self-control	59.78	50.15	63.25	58.00		1.38	0.71	—
Monitoring of treatment	59.33	59.65	61.91	59.89		0.14	0.99	-
Feeling of cohesion	56.06	45.05	62.59	80.33		5.53	0.14	-
Emotional representation	59.96	70.90	60.43	57.89		0.92	0.82	_
The essence of the disease:								
No of sympt.since contraction	53.41	56.05	61.75	83.06		5.11	0.16	-
No of symp. related to it	52.17	53.35	63.53	74.89		4.03	0.26	_
Causes of the disease:								
Psychological explanation	57.80	81.05	58.87	66.06		3.98	0.26	_
Action of risk factors	58.83	63.25	61.92	57.33		0.30	0.96	-
Strategies of coping:								
Fear absorption	69.43	79.43	55.29	62.67		6.40	0.09	-
Fighting spirit	68.70	51.05	60.63	52.06		2.73	0.45	_
Hopelessness-Helplessness	66.17	80.20	55.59	69.22		5.92	0.12	_
Positive revaluation	68.54	58.20	58.07	65.94		2.04	0.57	-
Acceptance of the disease	56.20	59.25	63.70	54.83		1.26	0.74	-
Mental state:								
Fear	57.06	87.95	58.07	67.33		7.12	0.07	-
Depression	60.63	71.85	60.34	55.56		1.22	0.75	-
Irritation/aggression	48.50	78.50	64.44	50.78	I < II	7.72	0.05*	0.06

p < 0.05; Legend: group I — patients with no previous experience, group II — patients with experience of previous own disease, group III — patients with experience of the disease in a close family member, group IV — persons with experience of their own disease and a disease in a close family member

Table S4. Type of cancer and patients without previous experience of their own serious, chronic disease or a disease in a close family member and the perception of the contracted disease, coping strategies, acceptance of the disease and their mental state

	Average ra			
	Breast	Colore		
Descention of the dimension	cancer	cancer	U	p
Perception of the disease:				
Duration of the disease	11.23	15.91	57.50	0.13
Consequences of the disease	12.18	15.25	68	0.34
Freq. of the occur. of sympt.	12.14	15.28	67.50	0.32
Self-control	13.27	14.50	80	0.72
Monitoring of treatment	14.36	13.75	84	0.87
Feeling of cohesion	13.18	14.56	79	79
Emotional representation	14.27	13.81	85	0.90
The essence of the disease:				
No of sympt.since contraction	10.41	16.47	48.50	0.05*
No of symp. related to it	10.77	16.22	52.50	0.08
Causes of the disease:				
Psychological explanation	12.91	14.75	76	0.58
Action of risk factors	12.36	15.13	70	0.39
Strategies of coping:				
Fear absorption	15	13.31	77	0.61
Fighting spirit	13.50	14.34	82.50	0.79
Hopelessness-Helplessness	11.55	15.69	61	0.20
Positive revaluation	11.09	16	56	0.12
Acceptance of the disease	17.55	11.56	49	0.06
Mental state:				
Fear	13.27	14.50	80	0.72
Depression	12.36	15.13	70	0.39
Irritation/aggression	14.50	13.66	82.50	0.79

**p* < 0.05

Table S5. Type of cancer and patients with previous experience of their own serious, chronic disease or a disease in a close family member and the perception of the contracted disease, coping strategies, acceptance of the disease and their mental state

	Average rank				
	Breast	Colorectal		U	
Perception of the disease:	cancer	cancer		U	p
Duration of the disease	44.85	51.97	876		0.22
Consequences of the disease	46.33	49.47	963	.50	0.59
Freq. of the occur. of sympt.	45.69	50.54	926		0.40
Self-control	48.10	46.49	997		0.78
Monitoring of treatment	49.99	43.30	885		0.25
Feeling of cohesion	55.84	33.44	540	.50	< 0.001*
Emotional representation	50.69	42.11	844		0.14
The essence of the disease:					
No of sympt.since contraction	46.89	48.53	996	.50	0.78
No of symp. related to it	47.03	48.30	100	4.50	0.83
Causes of the disease:					
Psychological explanation	48	46.66	100	3	0.82
Action of risk factors	41.72	57.24	691	.50	0.01*
Strategies of coping:					
Fear absorption	45.72	50.50	927	.50	0.41
Fighting spirit	45.33	51.16	904		0.31
Hopelessness-Helplessness	45.25	51.29	900		0.30
Positive revaluation	45.10	51.54	891		0.27
Acceptance of the disease	47.96	46.73	1005.50		0.83
Mental state:					
Fear	46.75	48.77	988		0.73
Depression	45.80	50.37	932 0.43		0.43
Irritation/aggression	50.19	42.97	874 0.2		0.20

**p* < 0.05