

Supplementary File

Figure S1. Factors associated with overall survival and hazard ratios

Chemotherapy Race/Factor		Stage	Hazard Ratio*	95% CI		p-value
				Lower CI	Upper CI	
Hispanic	No surgery	I	0.433	0.084	2.239	0.318
		II	0.385	0.127	1.166	0.091
		IV	0.827	0.510	1.256	0.443
	Surgery	I	1.95	1.266	1.611	< 0.001
		II	1.048	0.872	1.26	< 0.001
		IV	1.232	1.082	1.403	< 0.001
Asian	No surgery	I	0.035	n/a	n/a	0.593
		II	0.115	0.011	1.179	0.069
		IV	0.894	0.445	1.795	0.894
	Surgery	I	5.660	1.656	19.344	0.006
		II	1.184	0.743	1.888	0.477
		IV	2.141	0.849	5.400	0.107
Black	No surgery	I	1.647	0.168	16.115	0.032
		II	0.569	0.211	1.532	0.265
		IV	0.512	0.289	0.906	0.021
	Surgery	I	1.402	0.492	3.997	0.528
		II	1.119	0.772	1.621	0.553
		IV	2.940	1.756	4.924	< 0.001
White	No surgery	I	1.998	0.945	4.222	0.070
		II	0.558	0.352	0.885	< 0.001
		IV	0.718	0.554	0.930	0.012
	Surgery	I	4.352	3.040	6.230	< 0.001
		II	1.050	0.895	1.230	0.550
		IV	3.635	2.757	4.793	< 0.001

Radiation Race/Factor		Stage	Hazard ratio**	95% CI		p-value
				Lower CI	Upper CI	
White	No surgery	I	3.112	1.572	4.652	0.002
		II	0.603	0.384	0.822	0.028
		IV	1.025	0.789	1.261	0.852
	Surgery	I	2.637	1.881	3.393	< 0.001
		II	1.059	0.908	1.210	0.466
		IV	1.752	1.348	2.156	< 0.001

*races were compared to White patients